

# Soil arthropod species and their abundance in different chili management practices in freshwater swamps of South Sumatra, Indonesia

*By Marieska Verawaty*

PAPER • OPEN ACCESS

## Soil arthropod species and their abundance in different chili management practices in freshwater swamps of South Sumatra, Indonesia

To cite this article: S Herlinda *et al* 2021 *IOP Conf. Ser.: Earth Environ. Sci.* **713** 012022

View the [article online](#) for updates and enhancements.

You may also like

- [Locomotion of arthropods in aquatic environment and their applications in robotics](#)  
Bokeon Kwak and Joonbum Bae
- [Composition and structure community of arthropods in organic and conventional arabica coffee plantation](#)  
Husni, Jauharlina and Saifullah
- [Diversity of Soil Arthropods in Different Soil Stratification Layers, The National Park of Gede Pangrango Mountain, Cisarua Resort, West Java, Indonesia](#)  
Meityani and A P Dharna



**ECS Toyota Young Investigator Fellowship**

For young professionals and scholars pursuing research in batteries, fuel cells and hydrogen, and future sustainable technologies.

At least one \$50,000 fellowship is available annually.  
More than \$1.4 million awarded since 2015!

 Application deadline: January 31, 2023

**Learn more. Apply today!**

The advertisement features a dark background with a glowing globe and futuristic interface elements. The ECS TOYOTA logo is prominently displayed in the top right corner.

## Soil arthropod species and their abundance in different chili management practices in freshwater swamps of South Sumatra, Indonesia

S Herlinda<sup>1,2\*</sup>, R Fadli<sup>1</sup>, Hasbi<sup>2,3</sup>, C Irsan<sup>1,2</sup>, A Setiawan<sup>4</sup>, Elfita<sup>5</sup>, M Verawaty<sup>4</sup>,  
S Suwandi<sup>1,2</sup>, Suparman<sup>1</sup> and T Karenina<sup>6</sup>

<sup>1</sup> Department of Plant Pests and Diseases, Faculty of Agriculture,  
Universitas Sriwijaya, Indralaya 30662, South Sumatra, Indonesia

<sup>2</sup> Research Center for Sub-optimal Lands (PUR-PLSO), Universitas Sriwijaya,  
Palembang 30139, South Sumatra, Indonesia

<sup>3</sup> Department of Agricultural Engineering Program, Faculty of Agriculture,  
Universitas Sriwijaya, Indralaya 30662, South Sumatra, Indonesia

<sup>4</sup> Department of Biology, Faculty of Mathematics and Natural Sciences,  
Universitas Sriwijaya, Indralaya 30662, Indonesia

<sup>5</sup> Department of Chemistry, Faculty of Mathematics and Natural Sciences,  
Universitas Sriwijaya, Indralaya 30662, Indonesia

<sup>6</sup> Research and Development Agency of South Sumatra Province. Palembang 30136,  
South Sumatra, Indonesia

\*E-mail: sitiherlinda@unsri.ac.id

**Abstract.** The chili management practices can influence the predatory arthropod community. This study aimed to identify soil arthropod species and examine their abundance in different chili management practices in freshwater swamps of South Sumatra. The survey was conducted in three types of chili field, first without mulch and synthetic insecticides, and by fertilizing using manure (EF). The Conventional 1 (C-1) used plastic mulch, synthetic fertilizers, and synthetic insecticides. The Conventional 2 (C-2) used insecticides and synthetic fertilizers but without mulch. The total of all soil arthropod species was found in the different chili management practices of 24 species originating from Insecta, Arachnida, and Diplopoda. The highest number of soil arthropod species was found in EF, while the least number was found in C-1. This survey found species of predatory mites (*Macrocheles dispar*), spiders (*Pardosa birmanica*), and predatory insects (*Coccinella transversalis*) and other important predators such as *Pheropsophus occipitalis* and *Pardosa pseudoannulata*. The abundance of soil arthropods either predators, herbivores or neutral insects was the highest in EF while the lowest one was in the C-1. Overall, different chili management practices affected the number of species and abundance of soil arthropods; the environmentally friendly plot has the highest number of species and the largest abundance.

### 1. Introduction

Freshwater swamps or non-tidal lowlands are wetlands flooded with water from rivers or rain throughout the year [1]. The duration and depth of submerging with water determine the type of swamps. These types of freshwater swamps are shallowly flooded (depth <50 cm for 3 months),



Content from this work may be used under the terms of the [Creative Commons Attribution 3.0 licence](https://creativecommons.org/licenses/by/3.0/). Any further distribution of this work must maintain attribution to the author(s) and the title of the work, journal citation and DOI.

moderately flooded (depth 50-100 cm for 3-6 months), and deeply flooded (depth > 100 cm for 6 months) [2]. In freshwater swamps of South Sumatra, the local farmers grow rice, corn, chili or other adaptive vegetables in the dry season [3]. The rice is generally planted in moderately and deeply flooded swamps [4], whereas in shallowly flooded swamps, they are planted with chili [5] and corn [6].

In South Sumatra, especially in Ogan Ilir District which is the center of chili production, this vast expanse of chili is generally owned individually by many local farmers. The farmers are generally diverse in implementing chili management practices. The farmers having large capital generally use plastic mulch, manure and synthetic fertilizers, and synthetic insecticide spraying. Those having moderate capital generally do not use plastic mulch although they still use fertilizers and spray synthetic insecticides. There are a small number of farmers who start to implement environmentally friendly management using manure and certified seeds and without spraying synthetic insecticides. However, full tillage for growing chili is carried out by almost all farmers in South Sumatra.

The various management practices can affect the soil arthropod community in the agro-ecosystem. For example, full tillage causes species arthropod diversity to decrease significantly when compared to soil insects in the forest [7]. The collembolan population decreases significantly after being sprayed with an insecticide made from active carbofuran or phorate [8]. Likewise, predatory arthropods also decrease in abundance after pesticide applications [9]. The weed that grows at the surface of the soil can increase the abundance and diversity of predatory arthropod species [10]. Soil dwelling spiders are more abundant and have a diversity of species in the plot that adopts organic agriculture compared to conventional plot [11]. This study aimed to identify soil arthropod species and examine the arthropod abundances in different chili management practices in freshwater swamps of South Sumatra.

## 2. Materials and Method

### 2.1. Survey and Arthropod Sampling

The survey was conducted at the chili production center in South Sumatra, namely in Ogan Ilir District. The local farmers generally had chili management practices that were still conventional. Most of them still used synthetic fertilizers and insecticides, as well as full tillage, and some used plastic mulch. This method of managing chili by local farmers in Ogan Ilir District was generally grouped into three characteristics and selected as the sample plot for this study. The first type of plot was an environmentally friendly plot (EF) which was an ideal designed control plot for the management, i.e. did not use plastic mulch, used hybrid seeds, only used manure, sprayed bioinsecticide with active ingredient of *Beauveria bassiana*, and mechanically manual weeding. Bioinsecticide was made following the method of [12]. The bioinsecticide dose used in this EF plot was 2 L ha<sup>-1</sup> and manure of 20 tons ha<sup>-1</sup>. The second was the type of conventional local farmer (C-1) habit characterized by using plastic mulch, applying synthetic insecticides, using self-produced seeds originating from previous harvested fruit, and fertilizing using manure and artificial fertilizers. The third was the conventional type of local farmer (C-2) habit which did not use mulch, applied synthetic insecticides, used self-produced seeds originating from previous harvested fruit, and fertilized using manure and artificial fertilizers. The synthetic insecticide doses that used on C-1 plot (propinex, profenofos, and lamda sihalotrin) and C-2 plot (diphenconazole and diafentiuiron) based on the recommendations of the respective packages. The synthetic fertilizers applied in both fields were also following the recommendations and the dosage for manure of 20 tons ha<sup>-1</sup>.

The soil arthropods were sampled using pitfall traps following the method of [13] and Berlese funnel following the method of [14]. The sampling was carried out when chili plants were 13, 27, 41, 55, 69, 83, 97, 111, 125, and 139 days after planting (DAT). Each type of plot was repeated three times (3 sample sub-plots) with a total area of 1 ha per plot. Each plot was sampled at five observation points. Pitfall traps were installed on the ground for 24 hours, while the soil taken for Berlese funnel was top soil volume of 600 cm<sup>3</sup> (10 x 10 x 6 cm<sup>3</sup>). The obtained arthropoda were put into vials

containing 70% alcohol and were morphologically identified using identification books [15–17] and the number of individuals of each species from each survey location was recorded.

### 2.2. Data Analysis

Species data and the number of individuals were analyzed descriptively, and the data were displayed in tables and graphs.

## 3. Results and Discussions

### 3.1. Soil arthropod species in different chili management practices

The total of all soil arthropod species found in three types of chili management practices (EF, C-1, and C-2) was 24 species (Table 1). On the EF fields the number of arthropod species found the most was 17 species, 13 species on the conventional plot 1 (C-1) and 16 species on the conventional plot 2 (C-2). Of the three types of plot found in various guilds arthropods, i.e. predators, herbivores, and neutral insects, there were no parasitoids found. The soil arthropods found in EF plot were four species of predators, six species of herbivores and seven species of neutral insects. The soil arthropods found in C-1 plot were four species of predators, four species of herbivores, and 6 species of neutral insects. The soil arthropods found in C-2 plot were nine predator species, three species of herbivores, and four species of neutral insects. In this survey, there were interesting findings, namely species of predatory mites (*Macrocheles dispar*), spiders (*Pardosa birmanica*), and predatory insects (*Coccinella transversalis*). *M. dispar* was a predator of soil arthropods, such as *Collembola* [18].

Besides, this survey also found important predators, which were predators of rice pest insects, including *Pheropsophus occipitalis*, *Pardosa pseudoannulata*, and *Pardosa birmanica*. The highest number of soil arthropod species on this environmentally friendly plot (EF) was due to more species being able to settle on the plot, while in C-1 and C2 plot applied with synthetic pesticides there was a decrease in the number of soil arthropod species. The decrease occurred in the number of neutral insect species (dominated by *Collembola*) on the C-1 and C2 plot indicating that chili management practices in the two types of plot were unsuitable for neutral insects in the soil. The decreased number of neutral insect species was one reason for the application of synthetic pesticides [19]. Besides, the lowest number of soil arthropod species on C-1 plot using the plastic mulch resulted from rising soil temperatures unsuitable for the habitat of certain species. According to [20] the soil temperature using plastic mulch can reach 32.5°C.

### 3.2. Soil arthropod abundance in different chili management practices

The abundance of predatory soil arthropods in the environmentally friendly type (EF) was the highest compared to the conventional plot 1 (C-1) and conventional 2 (C-2) (Figure 1). Of the seven predatory arthropod families found (*Carabidae*, *Coccinellidae*, *Staphylinidae*, *Pentatomidae*, *Labiidae*, *Lycosidae*, and *Macrochelidae*), *Lycosidae* (pitfall trap samples) and *Macrochelidae* (Berlese funnel samples) were the most dominant in all types of chili plot, on C-2 plot, besides being dominant with *Lycosidae* and *Macrochelidae*, they were also dominated by *Carabidae* (pitfall trap samples). The abundance of soil predator arthropods that remained high on the EF plot was due to the production inputs that were used safely for the predators, for example, the applied *B. bassiana* did not endanger predator arthropods. The *B. bassiana* was generally specific to the order of *Lepidoptera* [1], [21] or *Homoptera* [22], [23], while the predators in this study did not originate from either order. The abundance of these soil predators was lower on the conventional C-1 and C-2 plot compared to the EF plot due to the intensive spraying of synthetic insecticides on both types of plot. The synthetic insecticides can kill soil arthropods from various orders [19].


**Table 1.** Arthropod species found in three chili fields with different management practices in South Sumatra, Indonesia

Class/Ordo/Families	Species	Guilds	EF		C-1		C-2	
			PT	BF	PT	BF	PT	BF
Insecta/Coleoptera/Carabidae	<i>Dromius piceus</i>	PR	-	-	+	-	+	-
Insecta/Coleoptera/Carabidae	<i>Pheropsophus occipitalis</i>	PR	-	-	-	-	+	-
Insecta/Coleoptera/Carabidae	<i>Pterostichus subovatus</i>	PR	-	-	-	-	+	-
Insecta/Coleoptera/Coccinellidae	<i>Coccinella transversalis</i>	PR	+	-	+	-	+	-
Insecta/Coleoptera/Staphylinidae	<i>Paederus littoralis</i>	PR	-	-	-	-	+	-
Insecta/Hemiptera/Pentatomidae	<i>Andranalus spinidens</i>	PR	+	-	-	-	-	-
Insecta/Dermoptera/Labiidae	<i>Labia</i> sp. A	PR	-	-	-	-	+	-
Arachnida/Araneae/Lycosidae	<i>Pardosa birmanica</i>	PR	+	-	+	-	+	-
Arachnida/Araneae/Lycosidae	<i>Pardosa pseudoannulata</i>	PR	-	-	-	-	+	-
Arachnida/Mesostigmata/Macrochelidae	<i>Macrocheles dispar</i>	PR	-	+	-	+	-	+
Insecta/Orthoptera/Gryllidae	<i>Gryllus bimaculatus</i>	HV	+	-	+	-	+	-
Insecta/Orthoptera/Acrididae	<i>Aiolopus strepens</i>	HV	+	-	+	-	+	-
Insecta/Orthoptera/Acrididae	<i>Atractomorpha crenulata</i>	HV	+	-	+	-	-	-
Insecta/Diptera/Muscidae	<i>Atherigona exigua</i>	HV	+	-	+	-	+	-
Insecta/Hemiptera/Alydidae	<i>Leptocorisa acuta</i>	HV	+	-	-	-	-	-
Arachnida/Acari/Tetranychidae	<i>Tetranychus</i> sp. A	HV	+	-	-	-	-	-
Diplopoda/Spirobolida/Trigoniulidae	<i>Trigonicilus corallinus</i>	NI	+	-	+	-	-	-
Insecta/Hymenoptera/Formicidae	<i>Dolichoderus thoracicus</i>	NI	+	-	+	-	+	-
Insecta/Hymenoptera/Formicidae	<i>Paratrechina longicornis</i>	NI	+	-	-	-	+	-
Insecta/Coleoptera/Scarabidae	<i>Aphodius rufipes</i>	NI	+	-	+	-	+	-
Insecta/Coleoptera/Cuculidae	<i>Cucujus clavipes</i>	NI	+	-	+	-	-	-
Insecta/Collembola/Actaletidae	Collembola A	NI	-	+	-	+	-	-
Insecta/Collembola/Neelidae	Collembola B	NI	-	+	-	-	-	-
Insecta/Collembola/Neelidae	Collembola C	NI	-	-	-	-	-	+
Insecta/Collembola/Paleotullbergiidae	Collembola D	NI	-	-	-	+	-	-
The number of species (S)			17		14		16	


Note: PR predator, HV herbivore, NI neutral insect, + arthropods found, - no arthropods found, BF berlese funnel, PT pitfall traps, EF environmentally friendly plot, C-1 and C-2 conventional plots

The herbivore abundance on the EF plot was the highest compared to C-1 plot and C-2 plot (Figure 2). Of the 5 families of herbivores found (Gryllidae, Acrididae, Muscidae, Alydidae, and Tetranychidae), the most dominant one was Gryllidae found on all types of chili plot. The herbivores were only found from sampling using the pitfall traps, while the sampling using Berlese funnels they were not found. The lowest abundance of herbivores was found on C-1 plot because according to [24] the plastic mulch on the plot caused the herbivore population to decline. Besides, the decrease in abundance was also caused by the synthetic insecticide sprays.


The abundance of neutral insects on EF plot was the highest compared to C-1 and C-2 plots (Figure 3). The lowest abundance of neutral insects was found on C-1 plot from both observations using pitfall traps and Berlese funnel. Of the 6 found families of the neutral insects (Formicidae, Scarabidae, Cuculidae, Actaletidae, Neelidae, and Paleotullbergiidae), the most dominant one was the Formicidae found on all types of chili plot. The Formicidae were added to neutral insect guilds because these families generally acted as feeding on plants and arthropod exudates, fungus culturing in fresh leaves or dead organic matter and partly as scavengers and omnivorous insects [25]. In addition to the Formicidae, the families (Actaletidae and Neelidae) from Collembola were predominantly found on the EF plot. The highest abundance of Collembola on the EF plot was caused by the applied *B. bassiana* which did not decrease its abundance. The lowest abundance of Collembola was on the C-1 plot applied by synthetic and mulched insecticides. The Collembola is sensitive to synthetic insecticides, the especially broad-spectrum [19]. The plastic mulch caused the abundance of Collembola to decrease because according to [24] the plastic mulch causes the soil microclimate to be less suitable for the habitat of soil insects.



**Figure 1.** Abundance of predatory arthropods on EF plot sampled using pitfall traps (A), berlese traps (B), C-1 plot sampled using pitfall trap (C), berlese traps (D), C-2 plot sampled using pitfall traps (E), dan berlese traps (F)



**Figure 2.** Abundance of herbivores sampled using *pitfall traps* on EF plot (A), on C-1 plot (B), and on C-2 plot (C)



**Figure 3.** Abundance of neutral insects on EF plot sampled using *pitfall traps* (A), *berlese traps* (B), C-1 plot sampled using *pitfall trap* (C), *berlese traps* (D), C-2 plot sampled using *pitfall traps* (E), dan *berlese traps* (F)



#### 4. Conclusions

The results showed that the total of all soil arthropod species found in different chilli management practices of 24 species originated from the three arthropod classes (Insecta, Arachnida, and Diplopoda). The highest number of soil arthropod species was found on environmentally friendly plot, while the lowest one was found in C-1 plot with plastic mulch. The survey found interesting findings on the species of predatory mites (*Macrocheles dispar*), spiders (*Pardosa birmanica*), and predatory insects (*Coccinella transversalis*) and other important predators such as *Pheropsophus occipitalis* and *Pardosa pseudoannulata*. The found predatory arthropods were Carabidae, Coccinellidae, Staphylinidae, Pentatomidae, Labiidae, Lycosidae, and Macrochelidae. The abundance of Lycosidae (spiders) and Macrochelidae (mites) was the highest on the EF plot, whereas on the C-1 plot they were the lowest in abundance. The neutral insects families found were the Formicidae, Scarabidae, Cuculidae, Actaletidae, Neelidae, and Paleotullbergiidae, and the families of the Collembola order (Actaletidae and Neelidae) were the most dominant on the EF plot. In contrast, on the C-1 plot they were the lowest abundance. The abundance of soil arthropods-the predators, herbivores, and neutral insects-was the highest on the EF plot, while the lowest one was on the C-1 plot. In consequence, the different chili management practices affect the number of species and the abundance of soil arthropods, and the environmentally friendly plot has the highest number of species and the largest abundance.

#### Acknowledgment

This research was funded by the Applied Research scheme with budget year of 2020 according to Directorate of Research and Community Service, Ministry of Research and Technology/National Research and Innovation Agency, Grant Contract Number: SP DIPA-042.06.1.401516/2020.

#### References

- [1] K. I. Hanif, S. Herlinda, C. Irsan, and Y. Pujiastuti, "The impact of bioinsecticide overdoses of *Beauveria bassiana* on species diversity and abundance of not targeted arthropods in South Sumatra ( Indonesia ) freshwater swamp paddy," *Biodiversitas*, vol. 21, no. 5, pp. 2124–2136, 2020.
- [2] B. Lakitan, A. Alberto, L. Lindiana, and K. Kartika, "The Benefits of Biochar on Rice Growth and Yield in Tropical Riparian Wetland , South Sumatra , Indonesia," vol. 17, pp. 111–126, 2018.
- [3] T. Karenina, S. Herlinda, C. Irsan, and Y. Pujiastuti, "Arboreal Entomophagous Arthropods of Rice Insect Pests Inhabiting Adaptive Vegetables and Refugia in Freshwater Swamps of South Sumatra," *Agrivita J. Agric. Sci.*, vol. 42, no. 2, pp. 1–10, 2020.
- [4] R. P. Ria, B. Lakitan, F. Sulaiman, and K. Kartika, "Cross-ecosystem utilizing primed seeds of upland rice varieties for enriching crop diversity at riparian wetland during dry season," *Biodiversitas*, vol. 21, no. 7, pp. 3008–3017, 2020.
- [5] E. Siaga, B. Lakitan, H. Hasbi, S. M. Bemas, L. I. Widuri, and K. Kartika, "Floating seedbed for preparing rice seedlings under unpredictable flooding occurrence at tropical riparian land," vol. 25, no. 2, pp. 326–336, 2019.
- [6] S. Herlinda, N. Octariati, and S. Suwandi, "Exploring entomopathogenic fungi from South Sumatra ( Indonesia ) soil and their pathogenicity against a new invasive maize pest , *Spodoptera frugiperda*," vol. 21, no. 7, pp. 2955–2965, 2020.
- [7] B. Zodinpuui, "Diversity of soil macroarthropods in shifting cultivation and forest ecosystem of Mizoram , Northeast India," vol. 11, no. 3, pp. 601–611, 2019.
- [8] A. Ghosal and A. Hati, "Impact of some new generation insecticides on soil arthropods in rice maize cropping system," *J. Basic Appl. Zool.*, vol. 80, no. 6, pp. 1–8, 2019.
- [9] C. Sattler, A. T. Gianuca, O. Schweiger, M. Franzén, and J. Settele, "Agriculture , Ecosystems and Environment Pesticides and land cover heterogeneity affect functional group and taxonomic diversity of arthropods in rice agroecosystems," *Agric. Ecosyst. Environ.*, vol.

- 297, no. May 2019, p. 106927, 2020.
- [10] I. Munir, A. Ghaffar, A. Aslam, M. K. Shahzad, and M. Jafir, "Impact of weeds on diversity of soil arthropods in Bt cotton field in Faisalabad Pakistan," vol. 26, no. 1, pp. 119–129, 2025.
- [11] X. He, Y. Qiao, L. Sigsgaard, and X. Wu, "Agriculture , Ecosystems and Environment The spider diversity and plant hopper control potential in the long-term organic paddy fields in sub-tropical area , China," *Agric. Ecosyst. Environ.*, vol. 295, no. June 2019, p. 106921, 2020.
- [12] G. Prabawati, S. Herlinda, and Y. Pujiastuti, "The abundance of canopy arthropods in South Sumatra (Indonesia) freshwater swamp main and ratooned rice applied with bioinsecticides and synthetic insecticide," *Biodiversitas*, vol. 20, no. 10, pp. 2921–2930, 2019.
- [13] S. Herlinda, S. Yudha, and R. Thalib, "Species richness and abundance of spiders inhabiting rice in fresh swamps and tidal lowlands in South Sumatra, Indonesia," *J. ISSAAS*, vol. 24, no. 1, pp. 82–93, 2018.
- [14] V. Nsengimana, B. A. Kaplin, F. Francis, and D. Nsabimana, "A comparative study between sampling methods for soil litter arthropods in conserved tree plots and banana crop plantations in Rwanda," *Int. J. Dev. Sustain.*, vol. 6, no. 8, pp. 900–913, 2017.
- [15] M. D. Pathak and Z. R. Khan, *Insect Pests of Rice*. Manila: The International Rice Research Institute (IRRI), 1994.
- [16] E. A. Heinrichs, F. E. Nwilene, M. J. Stout, B. A. R. Hadi, and T. Freita, *Rice Insect Pests and their Management*. London: Burleigh Dodds Science Publishing, 2016.
- [17] R. Whyte and G. Anderson, *A Field Guide to Spiders of Australia*. Queensland: CSIRO Publishing, 2017.
- [18] H. H. Koehler, "Predatory mites (Gamasina, Mesostigmata)," *Agric. Ecosyst. Environ.*, vol. 74, pp. 395–410, 1999.
- [19] G. K. Frampton and P. J. Van Den Brink, "Collembola and macroarthropod community responses to carbamate , organophosphate and synthetic pyrethroid insecticides : Direct and indirect effects," *Environ. Pollut.*, vol. 147, pp. 14–25, 2007.
- [20] P. Pramanik, K. K. Bandyopadhyay, D. Bhaduri, and R. Bha, "Effect of mulch on soil thermal regimes - A review," *Int. J. Agric. Environ. Biotechnol.*, vol. 8, no. 3, pp. 645–658, 2015.
- [21] D. R. Ayudya, S. Herlinda, and S. Suwandi, "Insecticidal activity of culture filtrates from liquid medium of *Beauveria bassiana* isolates from South Sumatra (Indonesia) wetland soil against larvae of *Spodoptera litura*," *Biodiversitas*, vol. 20, no. 8, pp. 2101–2109, 2019.
- [22] S. J. Lee, J. S. Yu, Y. S. Nai, B. L. Parker, M. Skinner, and J. S. Kim, "Beauveria bassiana sensu lato granules for management of brown planthopper, *Nilaparvata lugens* in rice," *BioControl*, vol. 60, no. 2, pp. 263–270, 2015.
- [23] E. Sumikarsih, S. Herlinda, and Y. Pujiastuti, "Conidial density and viability of *Beauveria bassiana* isolates from Java and Sumatra and their virulence against *Nilaparvata lugens* at different temperatures," *AGRIVITA J. Agric. Sci.*, vol. 41, no. 2, pp. 335–349, 2019.
- [24] A. Nyoike, W. Teresia, and E. Oscar, "Reusing plastic mulch for a second strawberry crop : Effects on arthropod pests , weeds , diseases and strawberry yields," *Florida Entomol.*, vol. 97, no. 3, pp. 928–936, 2020.
- [25] B. Gulcu, S. Hazir, E. E. Lewis, and H. K. Kay, "Evaluation of responses of different ant species (Formicidae) to the scavenger deterrent factor associated with the entomopathogenic nematode-bacterium complex," *Eur. J. Entomol.*, pp. 312–317, 2018.

# Soil arthropod species and their abundance in different chili management practices in freshwater swamps of South Sumatra, Indonesia

---

ORIGINALITY REPORT

---

# 13%

SIMILARITY INDEX

---

PRIMARY SOURCES

---

**1** André Cirilo De S Almeida, Cássio A Dierings, Celio Borella Júnior, Flávio G Jesus, Jose A F Barrigossi. 38 words — 1%  
"Resistance of rice genotypes to *Tibraca limbativentris* (Hemiptera: Pentatomidae)", *Journal of Economic Entomology*, 2019  
[Crossref](#)

---

**2** Gleidson Gimenes Rieff, Tiago Natal-da-Luz, Mathieu Renaud, Henrique M.V.S. Azevedo-Pereira et al. 36 words — 1%  
"Impact of no-tillage versus conventional maize plantation on soil mesofauna with and without the use of a lambda-cyhalothrin based insecticide: A terrestrial model ecosystem experiment", *Applied Soil Ecology*, 2020  
[Crossref](#)

---

**3** Muhammad Mohsin, Haseeb Ahmad, Muhammad Nabeel Nasir, Zain Ul Abideen et al. 35 words — 1%  
"Quantifying the Soil Arthropod Diversity in Urban Forest in Dera Ghazi Khan", *BioMed Research International*, 2022  
[Crossref](#)

---

**4** Benyamin Lakitan, Buyung Hadi, Siti Herlinda, Erna Siaga et al. 31 words — 1%  
"Recognizing farmers' practices and constraints for intensifying rice production at Riparian Wetlands in Indonesia", *NJAS - Wageningen Journal of Life Sciences*, 2018  
[Crossref](#)

---

5 Huayang Zhen, Xu Feng, Muhammad Ahmed Waqas, Monica Quevedo Cascante et al. "Solutions to neutralize greenhouse gas emissions of the rice value chain — A case study in China", Sustainable Production and Consumption, 2023

31 words — 1%

Crossref

---

6 Ramalingam Karthik Raja, Alagarsamy Arun, Mustapha Touray, Sebnem Hazal Gulsen et al. "Antagonists and Defense Mechanisms of Entomopathogenic Nematodes and their Mutualistic Bacteria", Biological Control, 2020

31 words — 1%

Crossref

---

7 So Eun Park, Jong Cheol Kim, Se Jin Lee, Mi Rong Lee et al. "Solid cultures of thrips-pathogenic fungi *Isaria javanica* strains for enhanced conidial productivity and thermotolerance", Journal of Asia-Pacific Entomology, 2018

31 words — 1%

Crossref

---

8 Juliana Marques Ferreira, Filippe Elias de Freitas Soares. "Entomopathogenic fungi hydrolytic enzymes: a new approach to biocontrol?", Journal of Natural Pesticide Research, 2023

30 words — 1%

Crossref

---

9 Finbarr G. Horgan, Ainara Peñalver-Cruz. "Compatibility of Insecticides with Rice Resistance to Planthoppers as Influenced by the Timing and Frequency of Applications", Insects, 2022

28 words — 1%

Crossref

---

10 Naoki Katayama, Miyuki Mashiko, Chieko Koshida, Yuichi Yamaura. "Effects of rice-field abandonment rates on bird communities in mixed farmland-woodland

28 words — 1%

landscapes in Japan", Agriculture, Ecosystems & Environment, 2021

Crossref

---

11 Nikola Jureková, Natália Raschmanová, Dana Miklisová, Ľubomír Kováč. "A comparison of collecting methods in relation to the diversity of Collembola in scree habitats", Subterranean Biology, 2021 27 words — 1%

Crossref

---

12 Endang Warih Minarni, Loekas Soesanto, Agus Suyanto, Rostaman. "Molecular identification of three entomopathogenic fungi infecting the brown plant hopper pest in Indonesia", Egyptian Journal of Biological Pest Control, 2021 25 words — 1%

Crossref

---

13 Ginbert P. Cuaton, Laurence L. Delina. "Two decades of rice research in Indonesia and the Philippines: A systematic review and research agenda for the social sciences", Humanities and Social Sciences Communications, 2022 25 words — 1%

Crossref

---

14 Sing-Shan Wu, Ching-Tzu Tseng, Yu-Hung Yang, Yao-Chia Liu et al. "Potential of a combination of entomopathogenic fungal strains and a non-ionic surfactant to control the fall armyworm (Spodoptera frugiperda)", Journal of Asia-Pacific Entomology, 2022 23 words — 1%

Crossref

---

15 M.A. Kader, M. Senge, M.A. Mojid, K. Ito. "Recent advances in mulching materials and methods for modifying soil environment", Soil and Tillage Research, 2017 22 words — 1%

Crossref

---

16 Doug Waterer, William Hrycan, Theresa Simms. "Potential to double-crop plastic mulch", Canadian Journal of Plant Science, 2008 19 words — < 1%

17 S. Korenko, K. Hamouzová, K. Kysilková, M. Kolářová, T.G. Kloss, K. Takasuka, S. Pekár. 13 words — < 1%  
"Divergence in host utilisation by two spider ectoparasitoids within the genus *Eriostethus* (Ichneumonidae, Pimplinae)", *Zoologischer Anzeiger*, 2018  
Crossref

18 Fátima Gonçalves, Cátia Nunes, Cristina Carlos, Álvaro López et al. 9 words — < 1%  
"Do soil management practices affect the activity density, diversity, and stability of soil arthropods in vineyards?", *Agriculture, Ecosystems & Environment*, 2020  
Crossref

19 Hennie Husniah, Udjiana S. Pasaribu, Yuza Z. Beta, Asep K. Supriatna. 9 words — < 1%  
"Non-cooperative and cooperative solutions of government subsidy on public transportation", *MATEC Web of Conferences*, 2018  
Crossref

20 Keita Goto, Shin Yabuta, Shotaro Tamaru, Peter Ssenyonga, Bore Emanuel, Naoya Katsuhama, Jun-ichi Sakagami. 8 words — < 1%  
"Root hypoxia causes oxidative damage on photosynthetic apparatus and interacts with light stress to trigger abscission of lower position leaves in *Capsicum*", *Scientia Horticulturae*, 2022  
Crossref

EXCLUDE QUOTES OFF

EXCLUDE SOURCES OFF

EXCLUDE BIBLIOGRAPHY OFF

EXCLUDE MATCHES OFF