

Comparison of Antenatal and Postnatal Findings of Body Stalk Anomaly

by Abarham Martadiansyah

Submission date: 10-May-2023 02:06PM (UTC+0700)

Submission ID: 2089294441

File name: COMPARISON_OF_antenatal_AND_POSTNATAL_FINDINGS.pdf (2.39M)

Word count: 1651

Character count: 9215

Comparison of Antenatal and Postnatal Findings of Body Stalk Anomaly

Peby M Lestari¹, Wim T Pangemanan², Ahmad Kurdi Syamsuri³, M. Hatta Ansyori⁴, Nuswil Bernolian⁵, Putri Mirani⁶, Abarham Martadiansyah⁷, Cindy Kesty⁸, Yurizka Sabrina⁹

ABSTRACT

Background: Body stalk anomaly was a rare, sporadic defect in the abdominal wall with the expulsion of the contents of the thoracoabdominal organs. Body stalk anomaly was reported in about one in 7,500 births within 10–14 weeks of gestation. Body stalk anomaly was a rare malformation with a prevalence of about 0.12 cases per 10,000 births (both live and stillbirth).

Aim: To compare the antenatal and postnatal findings of body stalk anomaly.

Case description: A 29-year-old multigravid patient was referred to our hospital due to a congenital anomaly. There were several findings on ultrasonography and magnetic resonance imaging (MRI), namely low-set ears, banana sign, neural tube defects, severe scoliosis, short umbilical cord, fetus attached to the placenta, abdominoschisis, thoracoschisis with some organs out, and defects in both legs. Hysterotomy was performed at 24 weeks of gestation. A male baby weighing 650 gm with a body length of 33 cm and head circumference of 24 cm was born. His Apgar score was 1/1/1 and survived for 30 minutes after birth. The postnatal examination of the baby, performed postmortem, confirmed the antenatal diagnosis.

Synopsis: A case about a comparison between ultrasound and MRI prenatal screening with postoperative findings in the diagnosis of body stalk anomaly.

Conclusion: Antenatal ultrasonography provided an accurate diagnosis of body stalk anomaly, and the results could be a consideration for detection of the defect earlier. Furthermore, it could help in patient counseling about poor outcomes in neonates and termination planning earlier to avoid other additional risks or even complications of delivery.

Keywords: Antenatal, Body stalk anomaly, Diagnosis, Magnetic resonance imaging, Ultrasound.

Journal of South Asian Federation of Obstetrics and Gynaecology (2022): 10.5005/jp-journals-10006-2045

Body stalk anomaly was a rare, sporadic defect in the abdominal wall with the expulsion of the contents of the thoracoabdominal organs.¹ Body stalk anomaly was reported in about one in 7,500 births within 10–14 weeks of gestation.² Body stalk anomaly was a rare malformation with a prevalence of about 0.12 cases per 10,000 births (both live and stillbirth).^{1,3}

The exact etiology of body stalk anomaly was unknown; three possible causes have been hypothesized, namely premature rupture of membranes, vascular compromise, and embryonic dysgenesis.¹

Van Allen et al. proposed that a diagnosis of BSA must meet at least two of the following three criteria, exencephaly/encephalocele with a facial cleft, thoracoschisis and abdominoschisis (midline defect), and limb defect (e.g., clubfoot, polydactyly, oligodactyly, syndactyly, brachydactyly, or amelia).^{2,4}

A 29-year-old multigravid patient was referred to our hospital due to a congenital anomaly. From anamnesis, she denied any history of consuming teratogenic drugs, infection during pregnancy, diabetes mellitus, premature rupture of membrane, previous congenital anomaly, and family history of congenital anomaly. Based on physical examination, the fundal height was palpable at the umbilicus (20 cm) with a normal fetal heart rate. There were several findings on ultrasonography, namely low-set ears, banana sign, neural tube defects, severe scoliosis, short umbilical cord, fetus attached to the placenta, abdominoschisis, thoracoschisis with some organs out, and defects in both legs (Fig. 1).

MRI result was similar to US findings. It showed a single live fetus intrauterine, the fetal position was bent with the head on the left inferolateral, and there were two superior and inferior extremities

^{1–7}Division of Maternal-Fetal Medicine, Department of Obstetrics and Gynecology, Dr Mohammad Hoesin General Hospital, Faculty of Medicine Sriwijaya University, Palembang, South Sumatera, Indonesia

^{8,9}Department of Obstetrics and Gynecology, Dr Mohammad Hoesin General Hospital, Faculty of Medicine Sriwijaya University, Palembang, South Sumatera, Indonesia


Corresponding Author: Cindy Kesty, Department of Obstetrics and Gynecology, Dr Mohammad Hoesin General Hospital, Faculty of Medicine Sriwijaya University, Palembang, South Sumatera, Indonesia, e-mail: cindykestyJL18@gmail.com

How to cite this article: Lestari PM, Pangemanan WT, Syamsuri AK, et al. Comparison of Antenatal and Postnatal Findings of Body Stalk Anomaly. *J South Asian Feder Obst Gynae* 2022;14(3):325–328.

Source of support: Nil


Conflict of interest: None

formed. The head surface of the gyrus appeared wide and smooth (lissencephaly). The ventricles appeared dilated asymmetrically, and there was an inhomogeneous intensity lesion in the left thalamus region which was suspicious for an intracranial mass. The right and left auricles appeared low (low-set ears). There are 2 lungs formed, positioned in the thoracic cavity. There were defects in thoracoabdominal organ followed by heart, intestines, and outside the thoracoabdominal cavity. The placenta was seen on the right anterolateral side, and the umbilical cord appeared short in single entanglement (Fig. 2). Laboratory examination showed normal IgM and IgG toxoplasma, IgM rubella, IgM CMV, IgM HSV I, IgM HSV II,




Figs 1A to E: Ultrasound examination showed (A) Hypotelorism, neural tube defects, a banana sign indicating cerebellar hypoplasia; (B) Thoracoschisis and abdominoschisis; (C) Fetus attached to the placenta and short umbilical cord; (D) Exstrophic bladder; and (E) 3D showing neural tube defects and scoliosis





Figs 2A to D: Magnetic resonance imaging examination showed a single live fetus intrauterine, the position of the fetus was bent with the head on the left inferolateral. There were 2 extremities superior and inferior formed. The head surface of the gyrus appeared wide and smooth (lissencephaly). The ventricles appeared asymmetrically dilated, there was an inhomogeneous intensity lesion in the left thalamus region, and an intracranial mass was suspected. Right and left auricles looked lower (low-set ears). There were 2 lungs formed, positioned in the thoracic cavity. There were thoracoabdominal defects followed by the heart, intestines and outside the thoracoabdominal cavity. The placenta was seen on the right anterolateral side, the umbilical cord appeared short in 1 circular tie



Figs 3A to C: Postoperative male neonate weighing 650 gm was born. There were abdominoschisis, lower limb deformity, scoliosis, and congenital vertical talus

IgG HSVII. However, there was increased IgG Rubella, IgG CMV, IgG HSV I, but they were not significant as an acute infection markers.

We conducted a joint conference with the ethical committee, religious leader, legal adviser, and clinicians from multidiscipline. From the joint conference, we agreed to perform hysterotomy due to the poor fetal prognosis (lethal anomaly). After we did informed consent and choice to the patient and her family, we did hysterotomy at 24 weeks of gestation.

A male baby weighing 650 gm with a body length of 33 cm and head circumference of 24 cm was born. His Apgar score was 1/1/1 and survived for 30 minutes after birth. The postnatal examination of the baby performed postmortem, confirmed the antenatal diagnosis (Fig. 3). There were abdominoschisis with scoliosis and congenital vertical talus. Postmortem pathological analysis could not be performed. In addition, the results of postoperative laboratory examinations showed an increased alpha-fetoprotein 918.56 ng/mL. Although the recurrence rate of body stalk anomaly remained low, we still suggested her for having multivitamin, such as high dose folic acid, vitamin D, and other antioxidants every day at least for 6 months before the next conception for preventing the recurrence of body stalk anomaly in the subsequent pregnancy. Furthermore, she should have routine antenatal care and do some screening in every trimester of her pregnancy with obstetricians.

Antenatal ultrasonography and MRI provided an accurate diagnosis of body stalk anomaly and the results could be a consideration for detection of the defect earlier. However, it

depended on the operator's skill, knowledge, and experience. Furthermore, it could help on patient counseling about poor outcomes on neonates and termination planning earlier to avoid other additional risks or even complications of delivery.

AUTHOR CONTRIBUTIONS

Peby Maulina Lestari and Yurizka Sabrina contributed to the acquisition of data. Wim T Pangemanan, Nuswil Bernolian, and Putri Mirani designed and drafted the article. Ahmad Kurdi Syamsuri and Abarham Martadiansyah were responsible for critical manuscript revision. M Hatta Ansyori and Cindy Kesty were responsible for the final approval of the version to be submitted.

REFERENCES

- Balachandran R, Samal SK. Fetal body stalk malformation-a rare case report. *Int J Reprod Contracept Obstet Gynecol* 2020;10(1):394. DOI: 10.18203/2320-1770.ijrcog20205807.
- Juusela AL, Gimovsky M. When ultrasonography reveals a fetal abdominal wall defect. *OBG Manag* 2021;33(1):34–44. DOI: 10.12788/obgm.0058.
- Singh A, Singh J, Gupta K. Body stalk anomaly: antenatal sonographic diagnosis of this rare entity with review of literature. *J Ultrason* 2017;17(69):133–135. DOI: 10.15557/jou.2017.0020.
- Balçı O, Ercan F, Sarıkaya S, et al. Body stalk anomaly in a 12-week pregnancy. Antenatal diagnosis of a uncommon neural tube defect: anterior encephalocele. 2016. p. 220–222.

Comparison of Antenatal and Postnatal Findings of Body Stalk Anomaly

ORIGINALITY REPORT

15%

SIMILARITY INDEX

6%

INTERNET SOURCES

11%

PUBLICATIONS

2%

STUDENT PAPERS

PRIMARY SOURCES

- 1** Yi Yang, Hong Wang, Zhenpeng Wang, Xuefeng Pan, Ying Chen. "First trimester diagnosis of body stalk anomaly complicated by ectopia cordis", Journal of International Medical Research, 2020 4%
Publication

- 2** Jephthah O. Odiba, Mzwandile A. Mabhala. "The Effectiveness of Glyburide Compared to Insulin in the Management of Gestational Diabetes Mellitus: A Systematic Review", Journal of Diabetes Mellitus, 2015 2%
Publication

- 3** misjournal.net 2%
Internet Source

- 4** Submitted to University of North Texas 1%
Student Paper

- 5** Merz, Eberhard, Bahlmann, Bernaschek, Bollmann, Chaoui, Deutinger, Eichhorn, Ertan, and Flock. "26 Anomalies of the

Gastrointestinal Tract and Anterior Abdominal Wall", Gastroenterologie compact, 2005.

Publication

6	www.scopemed.org Internet Source	1 %
7	vdocuments.mx Internet Source	1 %
8	www.acutecarepathlabs.com Internet Source	1 %
9	www.medicalscg.de Internet Source	1 %
10	Elena Spiller, Laura Salvador, Gianna Bogana, Francesca Neri, Guido Ambrosini, Erich Cosmi, Donato D'Antona. "Body stalk anomaly: Management of two dichorionic–diamniotic pregnancies", The Journal of Maternal-Fetal & Neonatal Medicine, 2009 Publication	1 %
11	Mariana Vide Tavares, Ana Patrícia Domingues, Margarida Tavares, Etelvina Fonseca, Paulo Moura. "Monoamniotic twins discordant for body stalk anomaly", The Journal of Maternal-Fetal & Neonatal Medicine, 2014 Publication	1 %
12	Takayuki Higuchi, Hiroaki Sato, Miho Iida, Seon Hye Kim, Yoshiaki Narimatsu, Mamoru	<1 %

Tanaka. "Early second-trimester diagnosis of body stalk anomaly by fetal magnetic resonance imaging", Japanese Journal of Radiology, 2013

Publication

13

J. M. Smrcek. "Prenatal ultrasound diagnosis and management of body stalk anomaly: analysis of nine singleton and two multiple pregnancies", Ultrasound in Obstetrics and Gynecology, 04/2003

Publication

<1 %

Exclude quotes On

Exclude matches < 2 words

Exclude bibliography On

Comparison of Antenatal and Postnatal Findings of Body Stalk Anomaly

GRADEMARK REPORT

FINAL GRADE

/0

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4
