

Antibacterial activity of Cordyline fruticosa leaf extracts and its endophytic fungi extracts

by Hary Widjajanti

Submission date: 20-Sep-2022 03:09PM (UTC+0700)

Submission ID: 1904379892

File name: ctivity_of_Cordyline_fruticosa_JURNAL_BIODIVERSITAS_DES_2019.pdf (647.97K)

Word count: 5885

Character count: 31144

Antibacterial activity of *Cordyline fruticosa* leaf extracts and its endophytic fungi extracts

ELFITA¹, MARDIYANTO², FITRYA², JUWITA EKA LARASATI², JULINAR¹, HARY WIDJAJANTI³, MUHARNI^{1*}

¹Department of Chemistry, Faculty of Mathematics and Natural Sciences, Universitas Sriwijaya, Jl. Raya Palembang-Prabumulih Km. 32, Indralaya, Ogan Ilir 30662, South Sumatra, Indonesia. Tel.: +62-71 1-580056, *email: muharnimyd@yahoo.co.id

²Department of Pharmacy, Faculty of Mathematics and Natural Sciences, Universitas Sriwijaya, Jl. Raya Palembang-Prabumulih Km. 32, Indralaya, Ogan Ilir 30662, South Sumatra, Indonesia

³Department of Biology, Faculty of Mathematics and Natural Sciences, Universitas Sriwijaya, Jl. Raya Palembang-Prabumulih Km. 32, Indralaya, Ogan Ilir 30662, South Sumatra, Indonesia

Manuscript received: 26 September 2019. Revision accepted: 29 November 2019.

Abstract. Elfita, Mardiyanto, Fitrya, Larasati JE, Julinar, Widjajanti H, Muharni. 2019. Antibacterial activity of *Cordyline fruticosa* leaf extracts and its endophytic fungi extracts. *Biodiversitas* 20: 3804-3812. Endophytic fungi live by forming colonies in plant tissues without harming the host plant. *Cordyline fruticosa* has been used as traditional medicine for the treatment of pathogenic bacterial infections; therefore, its endophytic fungi are expected to have similar activity. In this study, *C. fruticosa* leaves were extracted with gradient solvents and evaluated for their antibacterial activity using the Kirby-Bauer method against Gram-negative (*Salmonella typhi*, *Escherichia coli*) and Gram-positive (*Staphylococcus aureus*, *Bacillus subtilis*) bacteria. The extracts of endophytic fungi from *C. fruticosa* leaves were evaluated for their antibacterial activity, as well. The endophytic fungus with good antibacterial activity was identified by molecular and phylogenetic methods. Chemical compounds from endophytic fungus DB1 were isolated by chromatography and subsequently determined by spectroscopy. The methanol extract of *C. fruticosa* leaves showed strong antibacterial activity equivalent to the endophytic fungus (DB1). Molecular identification and analysis through the phylogenetic tree showed that the DB1 fungus has a high level of similarity to *Neurospora tetrasperma* strain APBDSDF108. The antibacterial compounds (compound 1 and 2) isolated from the endophytic fungus DB1 were identified as 4-hydroxy-5-phenylpenta-1,3-dien-1-yl acetate and ergosterol, respectively.

Keywords: Antibacterial activity, *Cordyline fruticosa*, endophytic fungi, *Neurospora tetrasperma*

INTRODUCTION

The World Health Organization (WHO) has reported that 90% of bacteria strains are resistant to antibiotics of first choice (Nwakanma et al. 2016). The emergence of pathogenic bacterial resistance to commercial antibiotics has become a serious and worrying health problem throughout the world as new antibiotic-resistant organisms emerge and spread globally every day. This situation encourages researchers to find new and efficient antibacterial metabolites. Therefore, there is a growing interest in screening the antibacterial activity of medicinal plants and their endophytic fungi.

Endophytic fungi are an important reservoir of therapeutically active compounds (Liang et al. 2012; Deshmukh et al. 2015; Pinheiro et al. 2017). Some endophytic fungi live in specific uninfected plant tissues, such as in the leaves, stems, roots, fruits, flowers, seeds, or rhizomes, and they do not cause damage to the host plant but have mutualistic symbiotic interactions. Endophytic fungi obtain nutrients, protection, and growing signals from host plants. Several endophytic fungal species have been reported to produce antibacterial, antifungal, antiviral, anti-inflammatory, and antitumor bioactive compounds that can protect host plants (Hussain et al. 2014; Bungtongdee et al. 2018; Praptiwi et al. 2018; Ding et al. 2019).

One plant-based traditional medicine for the treatment of infectious diseases is a decoction of *Cordyline fruticosa* leaves. *Cordyline* genus consists of more than 480 species spread throughout tropical and subtropical regions around the world. Based on ethnobotanical information, *C. fruticosa* is used as a hemostatic and also used to treat diarrhea, sore throat, neck pain, toothache, laryngitis, and mammary gland infections (Fouedjou et al. 2014; Fouedjou et al. 2016). Plants used in traditional medicine around the world are studied for their ability as host for endophytic fungi that have antibacterial potential, and recent developments reported that endophytic fungi are a potential reservoir of bioactive compounds useful for medicinal exploitation (Alshaibani et al. 2017; Liang et al. 2012; Chi et al. 2019).

Endophytic fungi have a tremendous ability to produce chemical compounds as their host plant, and the biological activity of compounds from endophytic fungi are often found to correspond to the biological activity of its host's compounds (Budiono et al. 2019; Fadhilla et al. 2019). In the previous research, two compounds from endophytic fungi associated with sambiloto (*Andrographis paniculata* Nees), i.e., 7-hydroxypiranopyridine-4-one and 1-(3,8-dihydroxy-4,6,6-trimethyl-6H-benzochromene-2-yl)oxypropane-2-one, have stronger antimalarial and antibacterial activity than the sambiloto extracts (Elfita et al. 2011; Elfita et al. 2015). Continuing our important work on

screening for antibacterial activity in medicinal plant extracts and their endophytic fungi, we found that ethyl acetate (EtOAc) extract from endophytic culture of *Neurospora tetrasperma* isolated from medicinal plant *C. fruticosa* showed antibacterial activity equivalent to methanol extract of the host plant. Details of the bioassay, isolation of pure compounds, and identification of their molecular structures are described in this paper.

MATERIALS AND METHODS

Material

Fresh leaves of *Cordyline fruticosa* were collected in November 2018 from Inderalaya, Ogan Ilir, South Sumatra, Indonesia, and identified at the Botany Laboratory, University of Sriwijaya, Indonesia (specimen number VIC 2685).

Extraction of *Cordyline fruticosa* leaves

The dried powder of *C. fruticosa* leaves (50 g) was extracted by maceration in a gradient solvent system, namely n-hexane, ethyl acetate (EtOAc), and methanol, for 24 hours each. Maceration was carried out three times, and each extract was evaporated with a rotary evaporator to obtain concentrated extracts.

Isolation of endophytic fungi

Isolation of endophytic fungi follows the protocols described by Liang et al. (2012) and Kalyanasundaram et al. (2015). Fresh leaves of *C. fruticosa* were rinsed with tap water, and the surface was sterilized by immersion in 70% ethyl alcohol (EtOH) for 2 minutes, followed by immersion in 1% sodium hypochlorite for 1 minute and then rinsed three times with sterile water. The fractions of leaves (0.5 × 0.5 cm) were placed on Petri plates containing potato dextrose agar (PDA) medium supplemented with 150 mg/L of chloramphenicol to inhibit bacterial growth. The plates were incubated at 27 ± 2 °C in the dark for 1–2 weeks. The hyphal tips were transferred to new PDA plates and then subcultured on a fresh PDA plate to obtain pure morphotype colonies.

Cultivation of endophytic fungi strains

All strains of the endophytic fungi were cultivated in potato dextrose broth (PDB). Six pieces of mycelia agar (5 × 5 mm²) were inoculated with ose needles into 200 mL potato dextrose broth (PDB) medium, and then incubated for four weeks at room temperature under static conditions. After the incubation period, mycelia were separated from the broth culture using filter paper. Then, 200 mL of EtOAc was added into the culture medium, and the broth was partitioned; this was repeated three times. Finally, the EtOAc extract was evaporated under reduced pressure using a rotary evaporator to obtain concentrated extracts (Marcellano et al. 2017).

Antibacterial activity of *Cordyline fruticosa* leaf extracts and endophytic fungi extracts

The antibacterial activity of the extracts was evaluated using the agar disc diffusion method against four bacteria,

which were *Salmonella typhi* (IPCCCB.11.669), *Escherichia coli* (ATCC 25922), *Staphylococcus aureus* (ATCC 25923), and *Bacillus subtilis* (ATCC 6633). All of the bacterial suspensions were equivalent to 1.5 × 10⁸ CFU/mL, based on the McFarland 0.5 standard. One hundred µl of the bacteria suspensions at a density of 5 × 10⁵ CFU/mL were inoculated on Petri plates containing 20 ml Mueller–Hinton agar (MHA). Paper discs (6.0 mm) were impregnated with concentrated extracts at 400 µg/disc in EtOH were placed on inoculated agar plates. Tetracycline 30 µg/disc was used as a positive control. Petri plates were incubated at 37 °C for 24 h. After the incubation period, clear zone inhibition was measured and clear zone inhibition of the extracts compared to standard antibiotics as the activity percentage. Antibacterial activity was classified as strong (inhibition zone ≥ 70%), moderate (inhibition zone 50–70%), or weak (inhibition zone < 50%). The minimum inhibitory concentration (MIC) values of the pure compounds were evaluated by dilution in EtOH to obtain series concentrations of 256, 128, 64, 32, 16, 8, and 4 µg/mL. The inhibition zone > 9 mm indicated that the compound had antibacterial activity at the corresponding concentration (Ding et al. 2019).

Molecular identification of endophytic fungi strain

Endophytic fungi that showed antibacterial activity were identified based on molecular analysis at the Research Center for Biology, Indonesian Institute of Sciences, Cibinong, Indonesia. Total DNA was extracted from the endophytic fungi strain following the protocol described by Hermansyah et al. (2017). The internal transcribed spacer (ITS) region of rDNA was amplified with PCR amplification using primer ITS-4 (5' TCC TCC GCT TAT TGA TAT GC 3') and primer ITS-5 (5' GCA AGT AAA AGT CGT AAC AAG G 3'). The PCR products were purified and sequenced using an automated DNA sequencer (ABI Prism 3130 Genetic Analyzer; Applied Biosystems, Foster City, CA, USA). The identified sequences were compared with the Basic Local Alignment Search Tool (BLAST) database (<http://www.ncbi.nlm.nih.gov/BLAST/>) to determine taxon/species. The phylogenetic trees were constructed using the neighbor-joining method with PHYLIP version 3.695, and phylogenetic trees were visualized by using the TreeView program with 1,000 bootstrap replicates (Budiono et al. 2019).

Cultivation and isolation of pure compounds from selected endophytic fungi

The endophytic fungi showed the highest antibacterial activity was re-cultivated to obtain more extract. The fungal strain was cultured in 5 × 200 mL of PDB medium, and after the cultivation period, the mycelium was separated from the liquid media. The liquid media was partitioned with EtOAc three times and evaporated to obtain concentrated EtOAc extract (5.7 g). The mycelium was dried in an oven at 60 °C to constant weight, and the mycelium dry powder was extracted by soxhlet with methanol. The methanol extract was evaporated to obtain concentrated methanol extract (4.9 g)

The concentrated EtOAc extract (5 g) of the broth culture was chromatographed on a silica gel column (70-230 mesh, 50 g) and eluted with a gradient solvent system, n-Hexane-EtOAc (10:0→0:10) and ethyl acetate methanol (EtOAc-MeOH) (10:0→0:10). The concentrated MeOH extract (4 g) of the dried biomass was chromatographed on a silica gel column (70-230 mesh, 60 g) and eluted with a gradient solvent system of n-Hexane - EtOAc (10:0→0:10) and EtOAc - MeOH (10:0→0:10). The chemical structures of the compounds were determined by ultraviolet (UV) spectra (Beck DU-7500 UV-visible scanning spectrophotometer; Beckman Coulter, Brea, CA, USA); infrared (IR) spectra (PerkinElmer Fourier transform IR [FT-IR] spectrometer; PerkinElmer, Waltham, MA, USA); nuclear magnetic resonance (NMR) spectra, including ¹H-NMR, ¹³C-NMR, HMQC, and ¹⁰⁵MB (Agilent DD2 ¹H-500 MHz and ¹³C-125 MHz; Agilent Technologies, Santa Clara, CA, USA); and mass spectroscopy (MS) (LC-MS ESI positive ion; PerkinElmer, Waltham, MA, USA).

RESULTS AND DISCUSSION

Antibacterial activity

Extraction of the dried powder (50 g) of *C. fruticosa* leaves produced concentrated extracts of n-Hexane (4.2 g), EtOAc (2.8), and EtOH (4.6 g). The antibacterial activity of the EtOH extract (Table 1) shows that it had strong antibacterial activity against three of the tested bacteria (*S. typhi*, *E. coli*, and *S. aureus*).

Six endophytic fungi strains were isolated from *C. fruticosa* leaves, but one isolate did not grow well in PDB medium. Five isolates were labeled as DB1-DB5. Each fungus isolate was cultivated in PDB medium for four weeks, and broth cultures were partitioned using EtOAc to

obtain secondary metabolites. Screening for antibacterial activity of these extracts was tested against *S. typhi*, *E. coli*, *S. aureus*, and *B. subtilis* (Table 1).

Phylogenetic analysis

Homology sequences of nucleotides were compared with BLAST (<http://www.ncbi.nlm.nih.gov>). The ClustalW program was used to align nucleotide sequences. Visualization and editing of sequences were carried out with the GeneDoc program. The phylogenetic trees (Figure 1) were constructed using the PHYLIP program and visualized by the TreeView program (Yohandini et al. 2014). The DB1 strain had the highest homology with *Neurospora tetrasperma* strain APBSDFS108. Sequence data of strain DB1 was submitted to GenBank and deposited with the accession number MN341013.

The results of molecular identification in the form of DNA sequencing of Strain DB1 (Contig-DB1) is as follows:

```
ACAAGGTCTCCGTTGGTGAACCAGCGGGAGGGATCAT
TACAGAGTTGCAAAACTTCCCCACAAACCATCGCGAATC
TTACCCGTACGGTTGCCTCGGCGCTGGCGGTCCGGAAG
GCCCTCGGGCCCTCCCGGATCCTCGGGTCTCCCGCTCGC
GGGAGGCTGCCCGCCGGAGTCCCGAACTAACTCTTGA
TATTTTATGTCTCTCTGAGTAAACTTTTAAATAAGTCAA
AACTTTCAACACCGGATCTCTTGGTTCTGGCATCGATGA
AGAACGCAGCGAAATGCGATAAGTAATGTGAATTGCAGA
ATTAGTGAATCATCGAATCTTTGAACGCACATTCGCGCT
CGCCAGTATTTGGCGAGCATGCCTGTTTCGAGCGTCATT
TCAACCATCAAGCTCTGCTTGGGATCCGCGGCT
GTCCGCGGTCCCTCAAAATCAGTGGCGGGCTCGCTAGTC
ACACCGAGCGTAGTAACTCTACATCGCTATGGTTCGCGG
GCGGGTCTTTCGCGTAAAACCCCATTTCTAAGGTTGA
CCTCGGATCAAGTAGGAATACCCGCTGAACCTAAGCAT.
```

Table 1. Antibacterial activity percentage of *C. fruticosa* leaf extracts and its endophytic fungi extracts (at a concentration of 400 µg/disk) compared to a standard antibiotic (tetracycline)

Sample	Extracts	Antibacterial activity ^a			
		<i>S. typhi</i>	<i>B. subtilis</i>	<i>E. coli</i>	<i>S. aureus</i>
<i>C. fruticosa</i> leaves	n-hexane	7.1 ± 0.8 (36.6*)	6.3 ± 0.2 (30.6*)	9.1 ± 1.4 (46.7*)	11.2 ± 1.7 (53.1*)
	Ethyl acetate	9.6 ± 1.1 (49.5*)	11.3 ± 2.3 (54.8**)	12.5 ± 1.9 (64.1**)	13.8 ± 2.2 (65.4**)
	Methanol	15.1 ± 1.4 (77.8***)	12.6 ± 0.9 (61.2**)	16.7 ± 3.2 (85.6***)	17.0 ± 2.6 (80.6***)
Endophytic fungi	DB1	14.2 ± 1.1 (73.2***)	14.7 ± 2.9 (71.4***)	16.0 ± 2.5 (82.0***)	14.3 ± 0.7 (67.8**)
	DB2	11.3 ± 2.3 (58.2**)	11.8 ± 1.2 (57.3**)	10.1 ± 0.7 (51.8**)	9.8 ± 0.8 (46.4*)
	DB3	-	-	7.2 ± 1.1 (36.9*)	-
	DB4	7.7 ± 1.3 (39.7*)	-	8.1 ± 1.3 (41.5*)	-
	DB5	-	7.3 ± 2.4 (35.4*)	-	-
Antibiotic	Tetracycline ^b	19.4 ± 3.2	20.6 ± 2.4	19.5 ± 1.9	21.1 ± 3.3

Note: ^aAntibacterial activity percentage of extracts compared to tetracycline (%); in 30 mm inhibition zone (mm) of extract/ inhibition zone (mm) of tetracycline: *** strong (≥ 70%), **moderate (50-70%), and *weak (< 50%) against *S. typhi*, *B. subtilis*, *E. coli*, and *S. aureus*. ^bpositive controls (30 µg/disk). (-) no inhibition zone




Figure 1. Phylogenetic tree of the DB1 strain

Chemical compounds from endophytic fungi

Neurospora tetrasperma

Separation of concentrated EtOAc extract (5 g) by column chromatography, yielded five subfractions (A–E). Fraction C (1.1 g) was subjected to a silica gel column (70–230 mesh, 20 g) with a solvent system n-Hexane - EtOAc (5:5→0:10) as the eluent and yielded four fractions (C1–C4). Fraction C3 (120 mg) was subjected to a silica gel column (70–230 mesh, 10 g) and eluted with n-Hexane - EtOAc (3:7) to yield compound 1 (32 mg).

Separation of concentrated MeOH extract (4 g), yielded four subfractions (A–D). Subfraction B (0.83 g) was separated over a silica gel (230–400 mesh, 20 g) and eluted with a solvent system of n-Hexane - EtOAc (8:2→3:7) to give four fractions (B1–B4). Subfraction B3 (126 mg) was subjected to a silica gel column (70–230 mesh, 10 g) to give three fractions (B3.1–B3.3). Subfraction B3.2 (58 mg) was purified with re-crystallization to yield compound 2 (white crystals, 41 mg).

The result of column chromatography of the EtOAc extract from the endophytic fungi broth culture was successfully isolated compound 1 that revealed as an

aromatic ester. Column chromatography of the MeOH extract from the biomass successfully isolated compound 2 and based on the spectroscopic data, compound 2 was identified as steroid ergosterol. The MIC values of the pure compounds were shown in Table 2.

Compound 1. White crystals; mp. 111–113°C; UV (MeOH) λ_{max} 216, 248 nm; UV (MeOH + NaOH) λ_{max} 216, 248 nm; IR (KBr) ν_{max} cm^{-1} : 3371.6 (OH), 3064.9 (CH-aromatic), 2922.2, 2852.7 (CH-aliphatic), 1664.9 (C=O esth), 1604.8 (C = C aromatic), 1259.5 (C = O ester); 1H , ^{13}C NMR: see Table 3.; LCMS –ESI m/z = 219.13 [M + H] $^+$.

Table 2. MIC values of tetracycline, compound 1 and 2 against four tested bacteria

Compound	MIC values (46 mL)			
	<i>S. typhi</i>	<i>B. subtilis</i>	<i>E. coli</i>	<i>S. aureus</i>
1	128	128	64	> 256
2	> 256	> 256	> 256	> 256
Tetracycline	4	4	4	4

The inhibition zone > 9 mm indicated that the compound has antibacterial activity at the corresponding concentration

Compound 2. White crystal; mp. 126-128°C ; UV (MeOH) λ_{\max} : no absorption; IR (KBr) ν_{\max} cm⁻¹: 3429.4 (OH), 29586.9 and 2870.1 (CH-aliphatic), 1627.9 (C=C), 1460.1 (bending CH₂ cyclic) and 73.3 for -CH₂(CH₃)₂γ, 1033.85 (cycloalkane); NMR (¹H-500 MHz; ¹³C-125.71 Hz; in CDCl₃) see Table 4.

The ¹H-NMR data (Table 3) showed the presence of three signals in the aromatic region at δ_{H} 7.2–7.7 ppm for five protons. The signals indicated the presence of an aromatic group with mono substitution. Other signals appeared at δ_{H} 7.67 (1H; d; J = 5.85 Hz); 6.27 (1H; dd; J = 2.45; 5.85 Hz); and 6.14 ppm (1H; d; J = 2.45 Hz), which indicate a side chain that has three vinyl protons located at C = C conjugated. The appearance of a signal at δ_{H} 3.81 ppm (2H; s) suggested that the side chain was connected through a methylene group attached directly to the aromatic ring. The signal at δ_{H} 1.25 ppm (3H; s) was a methyl ester group in the side chain.

The C-NMR and HMQC spectra (Table 3 and Figure 2) showed the presence of 11 signals for 13 carbons, two of which were C sp³. The signal of two carbons at δ_{C} 129.1 and 129.2 ppm is carbon equivalent to four carbon methines. The presence of one other methine carbon signal at δ_{C} 127.7 ppm indicated that compound 1 was in the aromatic group with mono substitution. The spectrum also revealed the presence of a carbon carbonyl ester at δ_{C} 179.3 ppm and two oxyarylcarbons at 155.3 and 168.2 ppm. The two oxyaryl carbons consisted of oxyaryl to ester and oxyaryl to hydroxyl groups. These signals indicated the presence of ester side chain conjugates with two double bonds (C = C).

The HMBC spectrum (Table 2 and Figure 2) showed the presence of a ¹H-¹³C correlation between the proton signal at δ_{H} 3.81 ppm (methylene signal) and the aromatic carbon δ_{C} 129.2; 134.6; 115.7 ppm. It indicates that the methylene group in the side chain was attached directly to the aromatic ring. Two bonds also separated the methylene proton with oxyaryl carbon (δ_{C} 168.2) and the three bonds with carbon methine (δ_{C} 115.7 ppm). The oxyarylmethine proton (δ_{H} 7.67 ppm) showed a long-distance correlation with oxyaryl carbon δ_{C} 168.2 ppm. Based on the spectroscopic analysis, compound 1 was identified as 4-hydroxy-5-phenylpenta-1,3-dien-1-yl acetate. The correlation between the structures of compound 1 and HMBC is shown in Figure 3.

The steroid characteristics of compound 2 (Figure 4 and Table 4) can be identified in the H-NMR spectrum by the presence of six methyl signals at δ_{H} 0.5-1.0 ppm. These signals include two methyl singlets found in the steroids skeleton and four methyl doublets in the side chains. The two vinyl protons at δ_{H} 5.38 (1H, d, J = 8.4 Hz) and 5.57 ppm (1H, d, J = 8.4 Hz) were mutually coupled in the steroids skeleton. Furthermore, two vinyl protons appear at δ_{H} 5.19 (1H, m) and 5.22 (1H, m) attached to the side chain. The typical proton signal for steroids is shown by

the presence of an oxygenated methine at δ_{H} 3.63 ppm (1 H, m) at C3.

Table 3. ¹H and ¹³C NMR data of compound 1 (¹H-500 MHz, ¹³C-125 Mhz, in CDCl₃), δ in ppm, J in Hz

No. C	δ_{C} 1	δ_{H} (Σ H, multiplicity, J) 1	HMBC
1	155.3	7.67 (1H; d; J=5.85 Hz)	116.8; 168.2; 179.3
2	116.8	6.27 (1H; dd; J=2.45; 5.85 Hz)	155.3
3	115.7	6.14 (1H; d; J=2.45 Hz)	168.2
4	168.2		
5	40.2	3.81 (2H; s)	129.2; 134.6; 115.7; 168.2
1'	134.6		
2'	129.2	7.23 (1H; m)	127.7; 129.1; 40.2
3'	129.1	7.35 (1H; m)	129.2; 134.6
4'	127.7	7.30 (1H; m)	129.2
5'	129.1	7.35 (1H; m)	129.2; 134.6
6'	129.2	7.23 (1H; m)	127.7; 129.1; 40.2
1''	179.3		
2''	29.8	1.25 (3H; s)	

Table 4. ¹H and ¹³C NMR data of compound 2 (¹H-500 MHz, ¹³C-125 Mhz, in CDCl₃) and ergosterol^b (¹H-400 MHz, ¹³C-100 Mhz, in CDCl₃ + CD₃OD, ppm) (Martinez et al. 2015)

No. C	¹³ C-NMR 2	¹ H-NMR 2	DEPT 2	¹³ C-NMR ^b
1	38.5		CH ₂	38.4
2	32.1		CH ₂	32.0
3	70.6	3.63 (1H, m)	CH	70.4
4	40.9		CH ₂	40.8
5	139.9		C	139.8
6	119.7	5.38 (1H, d, J=8.4 Hz)	CH	119.6
7	116.4	5.57 (1H, d, J=8.4 Hz)	CH	116.3
8	141.5		C	141.3
9	46.4		CH	46.2
10	37.2		C	37.0
11	21.2		CH ₂	21.1
12	39.2		CH ₂	39.1
13	42.9		C	42.8
14	54.7		CH	54.6
15	23.2		CH ₂	23.0
16	28.4		CH ₂	28.3
17	55.8		CH	55.7
18	12.2	0.63 (3H, s)	CH ₃	12.0
19	17.8	0.94 (3H, s)	CH ₃	17.6
20	40.6		CH	40.4
21	21.2	1.04 (3H, d, J=6.6 Hz)	CH ₃	21.1
22	135.7	5.22 (1H, m)	CH	135.6
23	132.1	5.19 (1H, m)	CH	132.0
24	43.0		CH	42.8
25	33.2		CH	33.1
26	19.8	0.83 (3H, d, J=7.15 Hz)	CH ₃	19.6
27	20.1	0.83 (3H, d, J=7.15 Hz)	CH ₃	19.9
28	16.4	0.92 (3H, d, J=6.8 Hz)	CH ₃	16.2




Figure 2. The ^{13}C -NMR (A), HSQC (B), and HMBC (C) spectra of compound 1 (^1H -500 MHz, ^{13}C -125 MHz, in CDCl_3)




Figure 3. Structure of 4-hydroxy-5-phenylpenta-1,3-dien-1-yl acetate and HMBC correlation




Figure 4. The ^1H -NMR (A) and ^{13}C -NMR (B) spectra of compound 2 (^1H -500 MHz, ^{13}C -125 MHz, in CDCl_3)

69




Figure 5. Structure of ergosta-5,7,22-trien-3 β -ol (ergosterol)

The ^{13}C -NMR spectrum showed the presence of 28 carbon signals consisting of six C sp^2 at δ_{C} 116.4, 119.7, 132.1, 135.7, 139.9, and 141.5 ppm. The signals indicate that steroids have two double bonds conjugating in the skeleton and one double bond on the side chain. Four C sp^2 conjugated in the skeleton consisted of two quaternary carbons and two methyl carbons. The spectrum showed the existence of one carbon bound to heteroatoms (OH groups), specifically, at δ_{C} 70.6 ppm. Based on the spectroscopic analysis and compared with data in the literature, compound **2** was ergosta-5,7,22-trien-3 β -ol (ergosterol). The structure of compound **2** was shown in Figure 5.

Discussion

Antibacterial activity of *C. fruticosa* leaf extracts (Table 1) showed that methanol extract had strong antibacterial activity against *S. typhi*, *E. coli*, and *S. aureus*. The antibacterial activity of methanol extract is related to the chemical content of *C. fruticosa* leaves, namely, the flavonoids farrerol, quercetin-3-O-[6"trans-p-Coumaroyl- β -D-glucopyranoside] (helichrysid), Apigenin-8-C-glucoside, quercetin-3-O- β -D-glucopyranoside, and quercetin-3-O- α -L-rhamnopyranosyl-(1-6)- β -D-glucopyranoside (rutin) (Fouedjou et al. 2016). Flavonoids are compounds that have a variety of biological activities, including antibacterial (Cushnie and Lamb 2005). Farrerol has been commonly used in traditional Chinese medicine as an antieczemic and antibacterial agent, and farrerol can inhibit the hemolysis of *S. aureus* by reducing the level of α -toxin production with the MIC value ranged from 4 to 16 $\mu\text{g/mL}$ against 35 *S. aureus* strains (Qiu et al. 2011).

The methanol extract has higher antibacterial activity than the n-hexane and EtOAc extract that may be due to the secondary metabolites in methanol extract that are more varied than other extracts. This is because the gradient fractionation causes non-polar and semi-polar compounds to be extracted into the n-hexane and ethyl acetate fractions. In contrast, the polar compounds with more variations are extracted into the methanol fraction. The synergistic effect of several compounds within extract often occurs in the methanol fraction, because the methanol fraction is more enriched with secondary metabolite compounds than the n-hexane and ethyl acetate fractions. The more variations of secondary metabolite compounds will be able to increase the potential for synergy. Research

conducted by Amin et al. (2015) that combined antibacterial compound quercetin with morin and routine that is not active against *S. aureus* (ATCC 43300) showed that the combination of all three flavonoids could improve antibacterial activity. The structure of the compound influences the synergistic effect, so the researchers combined resistant antibiotics with companion compounds to increase the working power of these antibiotics (Amin et al. 2015).

The antibacterial activity of endophytic fungi extract from *C. fruticosa* leaves showed that the DB1 extract (at a concentration of 400 $\mu\text{g/disc}$) exhibited the high level of growth inhibition (strong: ***) against *S. typhi*, *B. subtilis*, and *E. coli*. The extract of DB2 had moderate (**) inhibitory effects on the growth of *S. typhi*, *E. coli*, and *B. subtilis* and was weak (*) against *S. aureus*, while the other three endophytic extracts (DB3-DB5) showed weak antibacterial activity or no growth inhibition against the tested bacteria.

The antibacterial activity of the methanol extract of *C. fruticosa* leaves was comparable to the antibacterial activity of the endophytic fungal extract DB1 which has strong antibacterial against 3 tested bacteria and moderate antibacterial against one tested bacteria. Plants that have strong antibacterial activity may also be a potential source of antibacterial compounds from their endophytic fungus (Elfita et al., data not shown).

Results of the antibacterial assay showed that compound **1** has good antibacterial activity against *E. coli* (MIC 64 $\mu\text{g/ml}$) and moderate activity against *S. typhi* (MIC 128 $\mu\text{g/ml}$) and *B. subtilis* (MIC 128 $\mu\text{g/ml}$), while DB2 showed no activity against the four tested bacteria. Pure compound with MIC value <100 $\mu\text{g/mL}$ categorized as good antibacterial (Pinheiro et al. (2017).

The antibacterial activity of compound **2**, when compared to the antibacterial activity of several other compounds from the ergostanone group produced by *Colletotrichum* isolated from *Ilex canariensis*, showed the differences in antibacterial activity. The six ergostanone compounds showed good antibacterial activity against *E. coli* and *Bacillus megaterium* at 0.05 $\mu\text{g/disc}$ (5 $\mu\text{g/ml}$) (Deshmukh et al. 2015). Compound **2** and the six ergostanone compounds have the same skeletal structure, but different in the positions of the carbonyl, hydroxyl, and double bond groups, where the double bond and OH groups can affect antibacterial activity (Fouedjou et al. 2014). Compound **1** that isolated from the broth culture, is a new compound. Eight compounds have been reported from *C. fruticosa* leaves such as the steroidal saponins, isoquercitrin, helichrysid, quercetin 3-rutinoside, apigenin 8-C-b-D-glucopyranoside, and farrerol (Fouedjou et al. 2014). However, the good antibacterial activity of compound **1** correlates with the good activity of the host. The results of this study indicate that endophytic fungi isolated from *C. fruticosa* leaves produce different bioactive compounds than those originating from the host plant, they have similar activities. It is known that some endophytic fungi have the ability to produce bioactive compounds that are the same or similar to compounds

produced by their host plants. For instance, taxol is a well-known and widely used tetracyclic diterpenoid compound as an anticancer agent produced from types of yew trees (*Taxus* spp.). Similarly, podophyllotoxin compound is an aryl tetralin lignan compound with anticancer, antiviral, antioxidant, antibacterial, antirheumatic, and immunostimulating properties produced by *Sinopodophyllum* plant (Zhao et al. 2010; Garyali et al. 2013).

However, there are many compounds from endophytic fungi that are different from the compounds produced by host plant but they have strong biological activity. The endophytic fungus *Alternaria* sp. from the medicinal plant *Morinda officinalis* produces 13 compounds that don't have a chemotaxonomic relationship with its host plant. However, these compounds have inhibitory activities against four tumor cell lines and α -glucosidase (Wang et al. 2017; Lee et al. 2017). Calcul et al. (2013) reported the antimarial compounds produced by endophytic fungi from mangroves. Previously, Ravikumar et al. (2011) also reported the antimarial activity and the chemical content of mangrove plants is alkaloids, triterpenes, flavonoids, tannins, catechins, anthraquinones, phenols, sugars, and proteins. All of these classes of compounds do not have a chemotaxonomic relationship to the compounds produced by their endophytic fungi.

Results in this study showed that methanol extract of *C. fruticosa* leaves possess good antibacterial activity, and the endophytic fungus DB1 that identified as *N. tetrasperma* also has the potential as an antibacterial source. Therefore, *C. fruticosa* and the endophytic *N. tetrasperma* have the prospect of being developed as an antibacterial agent.

ACKNOWLEDGEMENTS

The authors thank the Ministry of Research, Technology, and Higher Education of the Republic of Indonesia, which provided research funding through *Penelitian Dasar Unggulan Perguruan Tinggi (PDUPT)* 2019, with contract no. 0060.08/UN9/SB3.LP2M.PT/2019. The authors also thank Dr. Laila Hanum for identification of the plant specimens.

REFERENCES

- Alshaibani MM, MohamadZin N, Jalil J, Sidik NM, Ahmad SJ, Kamal N, Edrada-Ebel R. 2017. Isolation, purification, and characterization of five active diketopiperazine derivatives from endophytic *Streptomyces* SUK 25 with antimicrobial and cytotoxic activities. *J Microbiol Biotechnol* 27 (7): 1249-1256.
- Amin MU, Khurram M, Khattak B, Khan J. 2015. Antibiotic Additive and Synergistic Action of Rutin, Morin and Quercetin Against Methicillin-Resistant *Staphylococcus aureus*. *BMC Compl Altern Med* 15: 59-70.
- Budiono, Elfiti, Muharni, Yohandini H, Widjanti H. 2019. Antioxidant activity of *Syzygium samarangense* L. and their endophytic fungi. *Molekul* 14 (1): 48-55.
- Bungtongdee N, Sopalan K, Laosripaiboon W, Lamtham S. 2018. The chemical composition, antifungal, antioxidant and antimutagenicity properties of bioactive compounds from fungal endophytes associated with Thai orchids. *J Phytopathol* 2019 (167): 56-64.

- Calcul L, Waterman C, Ma WS, Lebar MD, Harter C, Mutka T, Morton L, Maignan P, Olphen AV, Kyle DE, Vrijmoed L, Pang KL, Pearce C, Baker BJ. 2013. Screening mangrove endophytic fungi for antimarial natural products. *Mar Drugs* 11: 5036-5050.
- Chi WC, Pang KL, Chen WL, Wang GJ, Lee TH. 2019. Antimicrobial and iNOS inhibitory activities of the endophytic fungi isolated from the mangrove plant *Acanthus ilicifolius* var. *xiamenensis*. *Bot Stud* 60: 4-11.
- Cushine TPM, Lamb AJ. 2005. Antimicrobial activity of flavonoids. *Int J Antimicrob Agents* 26 (5): 343-356.
- Deshmukh SK, Verekar SA, Bhawe SV. 2015. Endophytic fungi: A reservoir of antibacterials. *Front Microbiol* 5 (715): 1-43.
- Ding Z, Tao T, Wang L, Zhao Y, Huang H, Zhang D, Liu M, Wang Z, Han J. 2019. Bioprospecting of novel and bioactive metabolites from endophytic fungi isolated from rubber tree *Ficus elastic* leaves. *J Microbiol Biotechnol* 29 (5): 731-738.
- Elfiti, Muharni, Munawar, Salmi, Oktasari A. 2011. Senyawa anti-malaria dari jamur endofitik tumbuhan sambiloto (*Andrographis paniculata* Nees). *J Natur Indonesia* 13 (2): 56-129. [Indonesian]
- Elfiti, Munawar, Muharni, Ivantri I. 2015. Chemical constituents from an endophytic fungus *Aspergillus* sp. (Sbd5) isolated from sambiloto (*Andrographis paniculata* Nees). *Microbiol Indonesia* 9 (2): 82-88. [Indonesian]
- Fadhilah, Elfiti, Muharni, Yohandini H, Widjanti H. 2019. Chemical compound isolated from antioxidant active extract of endophytic fungus *Cladosporium tenuissimum* *Svietenia mahagoni* leaf stalks. *Biodiversitas* 20 (9): 2645-2650.
- Fouedjou RT, Nguetefack-Mbuyo EP, Ponou BK, Nguetefack TB, Barboni L, Tapondjou LA. 2016. Antioxidant activities and chemical constituents of extracts from *Cordyline fruticosa* (L.) A. Chev. (Agavaceae) and *Eriobotrya japonica* (Thunb) Lindl. (Rosaceae). *Pharmacologia* 7 (2-3): 103-113.
- Fouedjou RT, Teponno RB, Quassinti L, Bramucci M, Petrelli D, Vitali LA, Fiorini D, Tapondjou LA, Barboni L. 2014. Steroidal saponins from the leaves of *Cordyline fruticosa* (L.) A. Chev. and their cytotoxic and antimicrobial activity. *Phytochem Lett* 7: 62-68.
- Garyali S, Kumar A, Reddy MS. 2013. Taxol production by an endophytic fungus, *Fusarium redolens*, isolated from Himalayan Yew. *J Microbiol Biotechnol* 23 (10): 1372-1380.
- Hermansyah, Adhiyanti N, Julinar, Rahadiyanto KY, Susilawati. 2017. Identification of *Candida* Species By Assimilation and Multiplex-PCR Methode. *J Chem Technol Metallurgy* 52 (6): 1070-1078.
- Hussain H, Kliche-Spory C, Al-Harrasi A, Al-Rawahi A, Abbas G, Green IR, Schulz B, Krohn K, Shah A. 2014. Antimicrobial constituents from three endophytic fungi. *Asian Pac J Trop Med* 7 (Suppl 1): S224-S227.
- Kalyanasundaram I, Nagamuthu J, Muthukumaraswamy S. 2015. Antimicrobial activity of endophytic fungi isolated and identified from salt marsh plant in Vellar Estuary. *J Microbiol Antimicrob* 7 (2): 13-20.
- YK, Bang HJ, Oh JB, Whang WK. 2017. Bioassay-guided isolated compounds from *Morinda officinalis* inhibit Alzheimer's disease pathologies. *Molecules* 22: 1638-1649.
- Liang H, Xing Y, Chen J, Zhang D, Guo S, Wang C. 2012. Antimicrobial activities of endophytic fungi isolated from *Ophiopogon japonicus* (Liliaceae). *BMC Compl Altern Med* 12: 238-244.
- Marcellano JP, Collanto AS, Fuentes RG. 2017. Antibacterial activity of endophytic fungi isolated from the bark of *Cinnamomum mercedoi*. *Pharmacogn J* 9 (3): 405-409.
- Nwakanma C, Njoku EN, Pharamat T. 2016. Antimicrobial activity of secondary metabolites of fungi isolated from leaves of bush mango. *Next Generation Seq Appl* 3 (3): 1-6.
- Pinheiro EAA, Pina JRS, Feitosa AO, Carvalho JM, Borges FC, Marinho PSB, Marinho AMR. Bioprospecting of Antimicrobial activity of extracts of endophytic fungi from *Bauhinia guianensis*. *Rev Argent Microbiol* 49 (1): 3-6.
- Praptiwi, Raunsai M, Wulansari D, Fathoni A, Agusta A. 2018. Antibacterial and antioxidant activities of endophytic fungi extracts of medicinal plants from Central Sulawesi. *J Appl Pharmaceut Sci* 8 (8): 069-074.
- Qiu J, Xiang H, Hu C, Wang Q, Dong J, Li H, Luo M, Wang J, Deng X. 2011. Subinhibitory concentration of farrerol reduces α -toxin expression in *Staphylococcus aureus*. *FEMS Microbiol Lett* 315: 129-133.
- Ravikumar S, Inbaneson SJ, Suganthi P, Gnanadesigan M. 2011. In vitro antiplasmodial activity of ethanolic extracts of mangrove plants from

- south east coast of India against chloroquine-sensitive *Plasmodium falciparum*. Parasitol Res 4: 873-878.
- Wang Y, Liu Hong-Xin, Chen Yu-Chen, Sun Zhang-Hua, Li Hao-Hua, Li Sai-Ni, Yan Ming-Li, Zhang Wei-Min. 2017. Two new metabolites from the endophytic fungus *Alternaria* sp. A744 derived from *Morinda officinalis*. Molecules 22: 765-771.
- Yohandini H, Julinar J, Muharni M. 2015. Isolation and phylogenetic analysis of thermophile community within Tanjung Sakti hot spring, South Sumatera, Indonesia. Hayati J Biosci 22: 143-148. [Indonesian]
- Zhao J, Zhou L, Wang J, Shan T, Zhong L, Liu X, Gao X. Endophytic fungi for producing bioactive compounds originally from their host plants. Curr Res Technol Edu Top Appl Microbial Microbial Biotechnol 2010: 567-576.

Antibacterial activity of Cordyline fruticosa leaf extracts and its endophytic fungi extracts

ORIGINALITY REPORT

28%

SIMILARITY INDEX

21%

INTERNET SOURCES

24%

PUBLICATIONS

11%

STUDENT PAPERS

PRIMARY SOURCES

1	www.ikprress.org Internet Source	1%
2	www.journal.ugm.ac.id Internet Source	1%
3	mail.scialert.net Internet Source	1%
4	digitalcommons.wayne.edu Internet Source	1%
5	naturalproduct.skku.edu Internet Source	1%
6	herbal-medicine.imedpub.com Internet Source	1%
7	ethnobotanica.us Internet Source	1%
8	www.longdom.org Internet Source	1%
9	aip.scitation.org Internet Source	<1%

10

Kute Lachu, Madhu Kamle, Rituraj Borah, Beauty Tiwari, Pradeep Kumar. "Chapter 1 Endophytic Bacteria: Application Against Biotic and Abiotic Stresses and Plant Health Improvements for Sustainable Agriculture", Springer Science and Business Media LLC, 2022

Publication

<1 %

11

eprints.univ-batna2.dz

Internet Source

<1 %

12

www.journal.uad.ac.id

Internet Source

<1 %

13

www.rjpharmacognosy.ir

Internet Source

<1 %

14

123docz.net

Internet Source

<1 %

15

P R W Astana, U Nisa, A Triyono, D Ardiyanto, U Fitriani, Z Zulkarnain, K P Adwaita, F Novianto. "Medicinal plants used by traditional healers for hemorrhoid treatment in Borneo island: Ethnopharmacological study RISTOJA", IOP Conference Series: Earth and Environmental Science, 2021

Publication

<1 %

16

Rufin Marie Kouipou Toghueo. "Bioprospecting endophytic fungi from genus

<1 %

as sources of bioactive metabolites ",
Mycology, 2019
Publication

17 Submitted to University of Mosul <1 %
Student Paper

18 e-journal.unair.ac.id <1 %
Internet Source

19 journal.unj.ac.id <1 %
Internet Source

20 download.atlantis-press.com <1 %
Internet Source

21 www.mysciencework.com <1 %
Internet Source

22 www.scipress.com <1 %
Internet Source

23 Xu, Lijian, Wei Meng, Cong Cao, Jian Wang,
Wenjun Shan, and Qinggui Wang. <1 %
"Antibacterial and Antifungal Compounds
from Marine Fungi", Marine Drugs, 2015.
Publication

24 hdl.handle.net <1 %
Internet Source

25 www.acmicrob.com <1 %
Internet Source

26

Nicholas T. Amponsah, E. Eirian Jones, Hayley J. Ridgway, Marlene V. Jaspers. "Identification, potential inoculum sources and pathogenicity of botryosphaeriaceous species associated with grapevine dieback disease in New Zealand", *European Journal of Plant Pathology*, 2011

Publication

<1 %

27

Submitted to Universitas Islam Indonesia

Student Paper

<1 %

28

Kyoko Sugai, Shin Watanabe, Taiki Kuishi, Shinya Imura, Keiichi Ishigaki, Masatsugu Yokota, Shinichi Yanagawa, Yoshihisa Suyama. "Extremely low genetic diversity of the northern limit populations of *Nypa fruticans* (Arecaceae) on Iriomote Island, Japan", *Conservation Genetics*, 2015

Publication

<1 %

29

Kathrine Blom, Øystein Kalsnes Jørstad, Rowan Thomas Faber, Ingar Stene - Johansen et al. "Primary vitrectomy or intravitreal antibiotics followed by early vitrectomy for acute endophthalmitis: A prospective observational study", *Acta Ophthalmologica*, 2022

Publication

<1 %

30

baadalsg.inflibnet.ac.in

Internet Source

<1 %

31	e-sciencecentral.org Internet Source	<1 %
32	exeley.com Internet Source	<1 %
33	Mirosław Giurg, Katarzyna Piekielska, Magdalena Gębala, Bartosz Ditkowski et al. "Catalytic Oxidative Cyclocondensation of - Aminophenols to 2 - Amino - 3 - phenoxazin - 3 - ones ", Synthetic Communications, 2007 Publication	<1 %
34	journals.pu.edu.pk Internet Source	<1 %
35	rjoas.com Internet Source	<1 %
36	Peng, J.. "The new bioactive diterpenes cyanthiwiggins E-AA from the Jamaican sponge Myrmekioderma styx", Tetrahedron, 20020923 Publication	<1 %
37	journal.bio.unsoed.ac.id Internet Source	<1 %
38	www.scielo.br Internet Source	<1 %
39	Marvin Gohrbandt, André Lipski, James W. Grimshaw, Jessica A. Buttress et al. " Low	<1 %

membrane fluidity triggers lipid phase separation and protein segregation ", Cold Spring Harbor Laboratory, 2021

Publication

40

Seham S. El-hawary, Abeer S. Moawad, Hebatallah S. Bahr, Usama Ramadan Abdelmohsen, Rabab Mohammed. " Natural product diversity from the endophytic fungi of the genus ", RSC Advances, 2020

Publication

<1 %

41

onlinelibrary.wiley.com

Internet Source

<1 %

42

repositorio.uwiener.edu.pe

Internet Source

<1 %

43

www.springerprofessional.de

Internet Source

<1 %

44

Submitted to Universiti Teknologi MARA

Student Paper

<1 %

45

repository.unair.ac.id

Internet Source

<1 %

46

www.acgpubs.org

Internet Source

<1 %

47

Submitted to Gauhati University

Student Paper

<1 %

48

Hanaa M. Sayed, Mahmoud H. Mohamed, Salwa F. Farag, Gamal A. Mohamed, Peter

<1 %

proksch. " A new steroid glycoside and furochromones from L. ", Natural Product Research, 2007

Publication

49

J Gacén, J Maillo. "The thermal memory of acrylic fibres", Journal of the Society of Dyers and Colourists, 2008

Publication

50

Jiazhang Qiu. "Subinhibitory concentrations of farrerol reduce α -toxin expression in *Staphylococcus aureus* : Farrerol reduces α -toxin production by *S. aureus*", FEMS Microbiology Letters, 02/2011

Publication

51

Kamel H. Shaker, Moustafa M. Zohair, Amal Zaki, Heba-tollah M. Sweelam, Warda E. Ashour. "LC-MS and GC-MS -based phytochemical perspectives and antimicrobial effects of Endophytic Fungus *Chaetomium ovatoascomatis* Isolated from *Euphorbia milii*", Research Square Platform LLC, 2022

Publication

52

Nattakorn Kuncharoen, Somboon Tanasupawat. "Chapter 14 Endophytic Actinomycetes: Secondary Metabolites and Genomic Approaches", Springer Science and Business Media LLC, 2022

Publication

<1 %

<1 %

<1 %

<1 %

53	Submitted to Universiti Putra Malaysia Student Paper	<1 %
54	Submitted to University of Strathclyde Student Paper	<1 %
55	data.epo.org Internet Source	<1 %
56	eksakta.ppj.unp.ac.id Internet Source	<1 %
57	jsfk.ffarmasi.unand.ac.id Internet Source	<1 %
58	libres.uncg.edu Internet Source	<1 %
59	openaccess.leidenuniv.nl Internet Source	<1 %
60	pdfs.semanticscholar.org Internet Source	<1 %
61	www.imb.ntou.edu.tw Internet Source	<1 %
62	Submitted to Tshwane University of Technology Student Paper	<1 %
63	Xiaojiang Qin, Xinrong Xu, Xiaomin Hou, Ruifeng Liang et al. "The pharmacological properties and corresponding mechanisms of	<1 %

farrerol: a comprehensive review",
Pharmaceutical Biology, 2021

Publication

64	creativecommons.org Internet Source	<1 %
65	istina.ipmnet.ru Internet Source	<1 %
66	profdoc.um.ac.ir Internet Source	<1 %
67	test.dovepress.com Internet Source	<1 %
68	www.rroj.com Internet Source	<1 %
69	www.thieme-connect.de Internet Source	<1 %
70	Choi, Y.A.. "A novel synthesis of 4,5-diaryl-6-arylamino-2,3-benzo-1,3a,6a-triazapentalenes", Tetrahedron Letters, 20030929 Publication	<1 %
71	Submitted to Macau University of Science and Technology Student Paper	<1 %
72	Truong Van Nguyen Thien, Thi Kieu Loan Vo, Phu Hoang Dang, Ngoc Vinh Huynh et al. "	<1 %

Two new sesquiterpenes from the stems of ", Natural Product Research, 2020

Publication

73

Xiao-Ling Wang. "Two New Sesquiterpenoid Glucosides from the Aerial Parts of *Saussurea involucrate*", *Journal of Integrative Plant Biology*, 5/2007

Publication

<1 %

74

acgpubs.org

Internet Source

<1 %

75

e-journal.biologi.lipi.go.id

Internet Source

<1 %

76

informaticsjournals.com

Internet Source

<1 %

77

peerj.com

Internet Source

<1 %

78

superiorforexsignals.com

Internet Source

<1 %

79

zookeys.pensoft.net

Internet Source

<1 %

80

"Phoma: Diversity, Taxonomy, Bioactivities, and Nanotechnology", Springer Science and Business Media LLC, 2022

Publication

<1 %

81

Carrie Waterman, Laurent Calcul, Jeremy Beau, Wai Sheung Ma et al. "Miniaturized

<1 %

Cultivation of Microbiota for Antimalarial Drug Discovery", Medicinal Research Reviews, 2016

Publication

82

Coruh, Irfan, Arzu Gormez, Sezai Ercisli, and Memnune Sengul. "Total Phenolic Content, Antioxidant, and Antibacterial Activity of *Rumex crispus* Grown Wild in Turkey", *Pharmaceutical Biology*, 2008.

Publication

<1 %

83

Dao Duc Thien, Tran Duc Dai, Trinh Thi Thuy, Nguyen Thi Hoang Anh, Nguyen Thanh Tam. "Ursane Triterpenoids from the Leaves of *Ficus hirta*", *Chemistry of Natural Compounds*, 2021

Publication

<1 %

84

Maurus Landolt, Sophie Stroheker, Valentin Queloz, Anja Gall, Thomas N. Sieber. "Does water availability influence the abundance of species of the *Phialocephala fortinii* s.l. – *Acephala applanata* complex (PAC) in roots of pubescent oak (*Quercus pubescens*) and Scots pine (*Pinus sylvestris*)?", *Fungal Ecology*, 2020

Publication

<1 %

85

Submitted to Universitas Warmadewa

Student Paper

<1 %

86

Vasiliki Saroglou, Anastasia Karioti, Ana Rancic, Kostas Dimas, Catherine Koukoulitsa,

<1 %

Maria Zervou, Helen Skaltsa. " Sesquiterpene Lactones from and Their Antibacterial and Cytotoxic Activities. Prediction of Their Pharmacokinetic Profile ", Journal of Natural Products, 2010

Publication

87

garuda.ristekdikti.go.id

Internet Source

<1 %

88

online.osim.ro

Internet Source

<1 %

89

updatepublishing.com

Internet Source

<1 %

90

www.eurjchem.com

Internet Source

<1 %

91

A. U. Turker, C. Usta. "Biological screening of some Turkish medicinal plant extracts for antimicrobial and toxicity activities", Natural Product Research, 2010

Publication

<1 %

92

F. Alali, M. Hudaib, T. Aburjai, K. Khairallah, N. Al-Hadidi. " GC-MS Analysis and Antimicrobial Activity of the Essential Oil from the Stem of the Jordanian Toothbrush Tree . ", Pharmaceutical Biology, 2008

Publication

<1 %

93

Fernanda Fraga Campos, Policarpo A Sales Junior, Alvaro José Romanha, Márcio SS

<1 %

Araújo et al. "Bioactive endophytic fungi isolated from *Caesalpinia echinata* Lam. (Brazilwood) and identification of beauvericin as a trypanocidal metabolite from *Fusarium* sp.", *Memórias do Instituto Oswaldo Cruz*, 2015

Publication

94

Kathawut Sopalun, Siriluck lamtham. "Isolation and screening of extracellular enzymatic activity of endophytic fungi isolated from Thai orchids", *South African Journal of Botany*, 2020

Publication

95

Maria Michela Salvatore, Anna Andolfi, Rosario Nicoletti. "The Genus *Cladosporium*: A Rich Source of Diverse and Bioactive Natural Compounds", *Molecules*, 2021

Publication

96

Miaomiao Cai, Mengyun Liu, Pengxiao Chen, Hong Liu, Yifei Wang, Depo Yang, Zhimin Zhao, Ping Ding. "Iridoids with anti-inflammatory effect from the aerial parts of *Morinda officinalis* How", *Fitoterapia*, 2021

Publication

97

Nattawadee Bungtongdee, Kathawut Sopalun, Wanpen Laosripaiboon, Siriluck lamtham. "The chemical composition, antifungal, antioxidant and antimutagenicity properties

<1 %

<1 %

<1 %

<1 %

of bioactive compounds from fungal endophytes associated with Thai orchids", *Journal of Phytopathology*, 2019

Publication

98

Nedeljko T. Manojlovic, Milan Novakovic, Vladeta Stevovic, Slavica Solujic. "Antimicrobial Metabolites from Three Serbian .", *Pharmaceutical Biology*, 2008

Publication

99

Sundaram Ravikumar, Samuel Jacob Inbaneson, Palavesam Suganthi, Murugesan Gnanadesigan. "In vitro antiplasmodial activity of ethanolic extracts of mangrove plants from South East coast of India against chloroquine-sensitive Plasmodium falciparum", *Parasitology Research*, 2010

Publication

100

Submitted to University of Yangon

Student Paper

101

pericles.pericles-prod.literatumonline.com

Internet Source

102

research.edgehill.ac.uk

Internet Source

103

storage.googleapis.com

Internet Source

104

www.bioseek.eu

Internet Source

<1 %

<1 %

<1 %

<1 %

<1 %

<1 %

<1 %

- 105 www.dovepress.com Internet Source <1 %
-
- 106 www.hindawi.com Internet Source <1 %
-
- 107 Emmanuel Jean Teinkela Mbosso, Silvère Ngouela, Jules Clément Assob Nguedia, Véronique Penlap Beng et al. " Spathoside, a cerebroside and other antibacterial constituents of the stem bark of ", Natural Product Research, 2008
Publication <1 %
-
- 108 Rabia Tanvir, Aqeel Javeed, Aamir Ghafoor Bajwa. "Endophyte bioprospecting in South Asian medicinal plants: an attractive resource for biopharmaceuticals", Applied Microbiology and Biotechnology, 2017
Publication <1 %
-
- 109 Regina Geris, Matheus A. Pinho, Elisangela F. Boffo, Thomas J. Simpson. "Naturally Occurring Partially Reduced Perylenequinones from Fungi", Journal of Natural Products, 2022
Publication <1 %
-
- 110 Romuald T. Fouedjou, Rémy B. Teponno, Luana Quassinti, Massimo Bramucci et al. "Steroidal saponins from the leaves of Cordyline fruticosa (L.) A. Chev. and their

cytotoxic and antimicrobial activity",
Phytochemistry Letters, 2014

Publication

111 Shuai-Cheng Wu, Fei Liu, Kui Zhu, Jian-Zhong Shen. " Natural Products That Target Virulence Factors in Antibiotic-Resistant ", Journal of Agricultural and Food Chemistry, 2019

Publication

112 "13th European Congress of Clinical Microbiology and Infectious Diseases", Clinical Microbiology and Infection, 2003

Publication

113 A Huda, R Ichwani, C T Handoko, M D Bustan, B Yudono, F Gulo. " Comparative photocatalytic performances towards acid yellow 17 (AY17) and direct blue 71 (DB71) degradation using Sn O flower-like structure ", Journal of Physics: Conference Series, 2019

Publication

114 G A Gowthami, Subhankar Das, Yalpi Karthik, I K Manjula. "Study of antibacterial, anti-proliferative and pro-apoptotic potential of the cell extracts of endophytic fungi and bacteria isolated from Pajanelia longifolia (Willd.) K. Schuman", Plant Science Today, 2021

Publication

115

Yoshiteru Murata, Ayako Sano, Yachiyo Ueda,
Tomo Inomata et al. "Molecular epidemiology
of canine histoplasmosis in Japan", *Medical
Mycology*, 2007

Publication

<1 %

Exclude quotes On

Exclude matches Off

Exclude bibliography Off