

# Antibacterial Activity of Traditional Medicine *Scurrula atropurpurea* (BL) DANS and their Endophytic Fungi

*by* Hary Widjajanti

---

**Submission date:** 20-Sep-2022 03:12PM (UTC+0700)

**Submission ID:** 1904380767

**File name:** ional\_Medicine\_Jurnal\_Key\_Engineering\_Materials\_24\_April\_2020.pdf (557.49K)

**Word count:** 4398

**Character count:** 21945

3

## Antibacterial Activity of Traditional Medicine *Scurrula atropurpurea* (BL) DANS and their Endophytic Fungi

Elfita Elfita<sup>1,a\*</sup>, Muharni Muharni<sup>1,b</sup>, Mardiyanto Mardiyanto<sup>2,c</sup>, Fitriya Fitriya<sup>2,d</sup>, Feti Fera<sup>2,e</sup> and Hary Widjajanti<sup>3,f</sup>

2

<sup>1</sup>Department of Chemistry, Faculty of Mathematics and Natural Sciences, University of Sriwijaya, Indralaya, Ogan Ilir, South Sumatra, 30662 Indonesia

<sup>2</sup>Department of Pharmacy, Faculty of Mathematics and Natural Sciences, University of Sriwijaya, South Sumatra, 30662 Indonesia

<sup>3</sup>Department of Biology, Faculty of Mathematics and Natural Sciences, University of Sriwijaya, Indralaya, Ogan Ilir, South Sumatra, 30662 Indonesia

\*elfita69@gmail.com, <sup>b</sup>muharnimyd@yahoo.co.id, <sup>c</sup>dianto72@yahoo.com, <sup>d</sup>fitriya\_apt@yahoo.com, <sup>e</sup>fetifera99@yahoo.com, <sup>f</sup>haryunsri@yahoo.com

**Keywords:** Antibacterial activity, *Scurrula atropurpurea*, endophytic fungi, *Neopestalotiopsis surinamensis*

19

1

**Abstract.** *Scurrula atropurpurea* known as benalu is a medicinal plant that has been used for the treatment of various diseases such as antibacterial. Plants with ethnomedicine history use to cure pathogenic bacterial infections and their endophytic fungi is a promising source of antibacterial compounds. This study aimed to compare the antibacterial activity of *S. atropurpurea* leaves and their endophytic fungi. The secondary metabolites were isolated from the leaves of *S. atropurpurea* and their endophytic fungi by the chromatography method. The antibacterial activity test was carried out by Kirby Bauer method against *Salmonella typhi* (IPCCCB.11.669) and *Escherichia coli* (ATCC 25922) as Gram (-) and *Bacillus subtilis* (ATCC 6633) and *Staphylococcus aureus* (ATCC 25923) as Gram (+). The antibacterial compound from *S. atropurpurea* was determined by spectroscopy analysis as Quercetin-3-O- $\alpha$ -L-rhamnopyranoside, while the antibacterial compound from endophytic fungi (strain BB1) as a lactone. Phylogenetic tree of strain BB1 has the highest homology with *Neopestalotiopsis surinamensis* strain CBS 450.74.

### Introduction

8

Antibiotics are medications used to prevent and treat bacterial infections. Antibiotic resistance occurs when bacteria change in response to the use of these medications. The resistance causes increased mortality to very dangerous levels in the whole of the world. The list of infections is increasing like pneumonia, tuberculosis, blood poisoning, and gonorrhoea. This infection is difficult to treat because antibiotics are less effective [1].

Antibiotic resistance is a global health problem [2, 3]. Research has shown that microbes have developed resistance to antibiotics through various molecular mechanisms (through various molecular mechanisms). This global problem causes an increase in research-oriented to the discovery of new antibiotic compounds. These compounds can be sourced from medicinal plants [4-6] or endophytic fungi that live symbiosis in the tissue of the medicinal plants. Endophytic fungi, especially those isolated from medicinal plants that have traditionally been used to cure infectious diseases, has potential as a source of new antibiotics [7].

Endophytic fungi are microorganisms that live in all healthy plant tissues (roots, fruits, stems, seeds, and leaves) without causing disease or morphological changes in the plant life cycle. These microbes live symbiosis mutually beneficial with their host plants and can jointly produce specific secondary metabolites [8,9]. The latest investigation has reported the potential of endophytic fungi in producing bioactive compounds such as taxol (anticancer), podophyllotoxin (antitumor), diorthothrome A and diorthothrome B (antimalarials), and Cytochalasin H (antibacterial). Endophytic fungi are a good source of producing new antibiotics [10-13]. In this paper reported the antibacterial activity and chemical content of *S. atropurpurea* and its endophytic fungi.

11

## Experimental Section

**Sampling and identification.** *Scurrula atropurpurea* leaves fresh are taken from around the Sriwijaya University campus, Indralaya, Ogan Ilir district, South Sumatra. Plant identification is carried out at the Botany Laboratory, FMIPA, Sriwijaya University.

**Extraction of *S. atropurpurea*.** *Scurrula atropurpurea* leaves fresh (650 g) were cleaned, cut into small pieces, then dried at room temperature until the weight is constant, then milled so that it is obtained dry powder weighed 312 g. The dry powder (300 g) was macerated using solvents with polarity increases, namely hexane, ethyl acetate, and methanol.

**Kirby Bauer method.** Antibacterial activity test using the Kirby Bauer method with four tested bacteria. Determination of the antibacterial activity of the compounds tested was carried out by comparing inhibitory zones in each concentration with inhibitory zones in three standard antibiotics. Antibacterial activity was expressed as inhibition percentage of a sample of antibiotics that are classified as strong (inhibition of  $\geq 70\%$ ), moderate (inhibition 50-70%), and weak (inhibition  $< 50\%$ ) [14].

**Exploration and structure elucidation of antibacterial compound from *S. atropurpurea* leaves.** Isolation of pure compound from the antibacterial active fraction, carried out by column chromatography, using silica gel G60 (70-230 mesh) as stationary phase, column length 45 cm, and diameter 2 cm. The extracts (5 g) were preabsorbed with silica gel (1:1), then eluted using gradient elution ( $\pm 30$  drops/minute). Eluates were collected in vials (10 mL) and used to monitor the eluates by TLC. Fractions that provide the same stain pattern were combined into one column fraction. Fractions that have potential stains are then purified. A pure compound from isolation results tested for antibacterial activity at a concentration of 400  $\mu\text{g}/\text{disk}$ . Furthermore, the molecular structure was identified by spectroscopic methods, including  $^1\text{H-NMR}$ ,  $^{13}\text{C-NMR}$ , HMQC, and HMBC.

**Isolation of endophytic fungi from *S. atropurpurea* leaves.** Isolation of endophytic fungal strains follows the method described in with slight modifications [15].

**Cultivation and antibacterial activity test of endophytic fungi extracts.** The cultivation of endophytic fungal strains was carried out by the method described in Elfita et al. [16]. Each fungal strain was cultured in 5 x 300 mL of PDB liquid medium. The extracts were tested the antibacterial activity with the Kirby Bauer method. Endophytic fungi extract with the highest antibacterial activity was continued to the separation of pure compound.

**Identification of endophytic fungi.** Endophytic fungi that provide the highest antibacterial activity was identified molecularly and analyzed by the construction of phylogenetic trees [14].

**Exploration of antibacterial compound from endophytic fungi and structure elucidation.** The active extract of the endophytic fungi was separated using column chromatographed with silica gel as a stationary phase (ratio 1:10). Samples that have been prepared by preabsorption was injected into the column and was eluted using eluent with polarity increases. Eluates were stored in bottles every 10 ml, and analyzed by thin layer chromatographed for grouped into the column fractions. Each column fraction was evaporated with a rotary evaporator and further separated and purified by re-chromatography and recrystallization techniques to obtain pure compound. The molecular structure of the compound was determined by spectroscopic analysis includes UV, IR,  $^1\text{H-NMR}$ ,  $^{13}\text{C-NMR}$ , HMQC, and HMBC.

## Results and Discussion

**Exploration of antibacterial compounds from *S. atropurpurea*.** Extraction of dried powder of *S. atropurpurea* leaves (300 g) in a gradient using n-hexane, ethyl acetate, and methanol produced concentrated extracts of 21.13, 29.87, and 30.24 g, respectively. Each fraction was tested for antibacterial activity against four bacteria, namely *S. typhi*, *B. subtilis*, *E. coli*, and *S. aureus*. Antibacterial activity of *S. atropurpurea* extracts compare to standard antibiotics showed to Table 1. These antibiotics can be used as a standard if they provide inhibitory zones with susceptible until intermediate categories when tested on an inoculum equivalent to McFarlan 0.5. If the antibiotic inhibition zone is categorized as resistant, it cannot be used as a standard [17].

**Table 1.** Antibacterial activity of fraction from *S. atropurpurea* leaves.

Sample	Fraction (400 µg /disk)	Test bacteria	Inhibition zone (mm)	Antibacterial activity percentage of ethyl acetate extracts compare to standard antibiotics (%)			
				Inhibition zone of Tetracycline (mm)	Inhibition zone of Streptomycin (mm)	Inhibition zone of Penicillin (mm)	
<i>S. atropurpurea</i> leaves	n-hexane	<i>S. typhi</i>	0 ± 0	0	0	0	
		<i>B. subtilis</i>	0 ± 0	0	0	0	
		<i>E. coli</i>	0 ± 0	0	0	0	
	Ethyl acetate	<i>S. aureus</i>	6.3 ± 0.61	30.4*	33.3*	37.1*	
		<i>S. typhi</i>	0 ± 0	0	0	0	
		<i>B. subtilis</i>	0 ± 0	0	0	0	
		<i>E. coli</i>	10.1 ± 2.11	52.6**	54.3**	56.4**	
		<i>S. aureus</i>	11.6 ± 1.64	56.0**	61.4**	68.2**	
		Methanol	<i>S. typhi</i>	7.2 ± 0.98	36.4*	36.7*	40.7*
			<i>B. subtilis</i>	11.3 ± 2.30	57.9**	61.4**	60.1**
	<i>E. coli</i>		13.6 ± 1.86	70.8***	73.1***	76.0***	
			<i>S. aureus</i>	14.9 ± 2.11	72.0***	78.8***	87.6***

Note: \*\*\* strong: inhibition of ≥ 70%    \*\* moderate: inhibition 50-70%    \* weak: inhibition < 50%

The methanol extract showed the highest antibacterial activity against *E. coli* and *S. aureus*. Lim et al. (2016) reported that several *Scurrula* plant members, such as *S. atropurpurea* had good antibacterial activities against bacteria of Gram-positive and negative [18].

The methanol extract (5 g) was carried out to column chromatography over silica gel 60 (70-230 mesh). A total of 81 eluates were collected. The same fraction was combined to obtain six fractions (F1-F6). The antibacterial activity of column fractions shows that F-2 was more active compared to other column fractions. The F2 fraction showed a major compound under a UV lamp at λ 254 nm, and it was further to the purification to yield compound 1 as a yellow crystal (96 mg). The results of the antibacterial activity test on 4 test bacteria showed that compound 1 (at a concentration 400 µg/disk) had a moderate category against *E. coli*.

Phytochemical tests showed the compound 1 as a flavonoid group. The <sup>1</sup>H-NMR spectrum revealed 11 signals with 13 numbers of protons. The presence of five aromatic protons attached to rings A and B of flavonoids. The appearance of five proton signals in a chemical shift around 3-5 ppm indicated that compound 1 has a glucoside group. These signals in the area often appear overlapping. The characteristics of the glucoside group are the presence of anomeric protons (H1) with multiplicity of doublet in δ<sub>H</sub> 6-4 ppm. The small coupling value of H1 and H2 (1.3 Hz) indicated the glucoside is an anomeric α. The type of rhamnosyl group attached at the compound is the appearance of a methyl (3H, d) signal at δ<sub>H</sub> around 1 ppm (J = ±7 Hz).

The <sup>13</sup>C-NMR spectrum showed the presence of 21 carbon signals, 15 carbon for the flavonol skeleton, and six for the rhamnosyl group. This compound 1 as flavonol rhamnoside indicated the presence of carbonyl carbon signals at δ<sub>C</sub> 179.7 ppm. The spectrum of compound 1 revealed seven

oxyaryl (= C-O) carbon atoms at  $\delta_c$  136-166 ppm. The signals indicated the presence of four hydroxyl groups attached to flavonoids, two hydroxyl groups in the A and C rings, respectively. The hydroxyl group in the C ring is substituted with the rhamnosyl group.

Based on spectrum analysis and after compared with literature [19] (Table 2), it was concluded that compound **1** as quercetin-3-O- $\alpha$ -L-rhamnopyranoside. The molecular formula  $C_{21}H_{20}O_{11}$ , with DBE = 12. The HMBC correlation and  $\delta$ -assignment of compound **1** showed in Fig. 1.

**Table 2.**  $^1H$  and  $^{13}C$  NMR data of compound **1** recorded at  $^1H$ -500 MHz,  $^{13}C$ -125 Mhz, in  $CD_3OD$  dan  $1''$  recorded at  $^1H$ -400 MHz,  $^{13}C$ -100 Mhz, in DMSO, ppm [19]

No. C	$\delta_c$ (ppm) 1	DEPT 1	$\delta_H$ (ppm), $\Sigma H$ , mult, J (Hz) 1	HMBC 1	COSY 1	$\delta_c$ (ppm) 1'	$\delta_H$ (ppm) 1'
2	159.4	C				157.9	
3	136.3	C				134.9	
4	179.7	C				178.3	
5	163.3	C				161.8	
6	99.9	CH	6.20 (d, 2.0)	163.3; 166.0; 94.8; 106.0	6.37	98.4	6.21 br,s
7	166.0	C				164.5	
7	94.8	CH	6.37 (d, J=2.0)	158.6; 166.0; 99.9; 106.0	6.20	93.4	6.33 br,s
9	158.6	C				157.1	
10	106.0	C				104.6	
1'	122.9	C				121.6	
2''	117.0	CH	7.33 (d, J=2.6)	123.0; 149.9; 159.4; 146.5		115.6	7.32 m
3''	146.5	C				145.0	
4''	149.9	C				148.4	
5''	116.4	CH	6.91 (d, J=8.4)	146.5; 149.9; 123.0	7.30	115.0	6.91 (d, J=8.4)
6''	123.0	CH	7.30 (dd, J=8.4; 2.6)	117.0; 149.9; 159.4	6.91	121.5	7.32 m
1''	103.6	CH	5.35 (d, J=1.3)	136.3; 72.1	4.22	102.2	5.30 br, s
2''	71.9	CH	4.22 (m)	73.3;	5.35; 3.75	70.8	4.22 m
3''	73.3	CH	3.34 (m)	72.2		70.7	3.20-3.76 m
4''	72.2	CH	3.42 (m)	73.3	0.94	70.7	3.20-3.76 m
5''	72.1	CH	3.75 (dd, J=3.25; 6.50)	73.3	4.22; 3.44	70.6	3.20-3.76 m
6''	17.7	CH <sub>3</sub>	0.94 (d, J=6.50)	72.2	3.42	16.4	0.93 (d, J=5.4)

In this study, it was revealed that the antibacterial activity decreased from the methanol fraction to pure compound. This might be due to the synergistic effect in the methanol fraction, which strengthens its antibacterial activity. Amin et al. (2015) have reported the antibacterial activity of flavonoid compounds, namely quercetin, morine, and rutin. The quercetin, morine, and rutin (at 500  $\mu g$  / disk) against *S. aureus* (ATCC 43300) showed the inhibition zones  $13.5 \pm 0.21$ ; 0.0; and 0.0 mm, respectively [20]. When the three compounds are combined, namely quercetin + morin + rutin, the inhibition zone increases to  $16.5 \pm 0.21$  mm. Inhibition zones also increase when flavonoids are combined with resistant antibiotics. Flavonoids, alone or in combination, also damage bacterial cell membranes [20].




Figure 1. The HMBC correlation and  $\delta$ -assignment of compound 1

53  
Antibacterial activity test of endophytic fungi extracts from *S. atropurpurea* leaves. Four strains of endophytic fungi were isolated from *S. atropurpurea* leaves, namely BB1-BB4. Each isolate has been cultivated in 5x300 mL of PDB medium for 6 weeks. The medium has been partitioned with ethyl acetate and evaporated to remove the solvent. The crude extract of each endophytic fungus is BB1 = 2.1 g; BB2 = 2.4 g; BB3 = 1.6 g; and BB4 = 1.9 g. The inhibition zone of all ethyl acetate extracts from endophytic fungi against four test bacteria compared to standard antibiotics is shown in Table 3.

Antibacterial activity of endophytic fungi extracts (Table 3) showed that BB1 isolates had strong activity against four bacteria (*S. typhi*, *B. subtilis*, *E. coli* and *S. aureus*), but has moderate activity against *S. aureus* by comparing tetracycline. Isolate BB1 generally has broad-spectrum. Table 3 reveals that BB4 provides moderate activity against four test bacteria. Two other fungal isolates (BB2 and BB3) have weak to inactive activity.

17  
Table 3. Antibacterial activity of endophytic fungi extracts from *S. atropurpurea* leaves

Sample	Extracts (400 $\mu$ g/disk)	Test bacteria	Inhibition zone (mm)	Antibacterial activity percentage of ethyl acetate extracts compare to standard antibiotics (%)		
				Inhibition zone of Tetracycline (mm)	Inhibition zone of Streptomycin (mm)	Inhibition zone of Penicillin (mm)
				<i>S. typhi</i> (19.8)	<i>S. typhi</i> (19.6)	<i>S. typhi</i> (17.7)
				<i>B. subtilis</i> (19.5)	<i>B. subtilis</i> (18.4)	<i>B. subtilis</i> (18.8)
				<i>E. coli</i> (19.2)	<i>E. coli</i> (18.6)	<i>E. coli</i> (17.9)
				<i>S. aureus</i> (20.7)	<i>S. aureus</i> (18.9)	<i>S. aureus</i> (17.0)
	BB 1	<i>S. typhi</i>	14.7 $\pm$ 1.80	74.2***	75.0***	83.1***
		<i>B. subtilis</i>	14.0 $\pm$ 2.25	71.8***	76.1***	74.5***
		<i>E. coli</i>	16.3 $\pm$ 1.69	84.9***	87.6***	91.1***
		<i>S. aureus</i>	13.8 $\pm$ 1.08	66.7**	73.0***	81.2***
	BB2	<i>S. typhi</i>	0 $\pm$ 0	0	0	0
		<i>B. subtilis</i>	0 $\pm$ 0	0	0	0
		<i>E. coli</i>	8.3 $\pm$ 0.65	43.2*	44.6*	46.4*
		<i>S. aureus</i>	7.2 $\pm$ 1.30	34.8*	38.1*	42.4*
Endophytic fungi	BB3	<i>S. typhi</i>	0 $\pm$ 0	0	0	0
		<i>B. subtilis</i>	8.1 $\pm$ 1.05	41.5*	44.0*	43.1*
		<i>E. coli</i>	0 $\pm$ 0	0	0	0
		<i>S. aureus</i>	10.7 $\pm$ 2.02	51.7**	56.6**	62.9**
	BB4	<i>S. typhi</i>	11.1 $\pm$ 1.24	56.1**	56.6**	62.7**
		<i>B. subtilis</i>	13.0 $\pm$ 1.08	66.7**	70.6***	69.1**
		<i>E. coli</i>	10.7 $\pm$ 1.27	55.7**	57.5**	59.8**
		<i>S. aureus</i>	9.8 $\pm$ 1.41	47.3*	51.8**	57.6**

Note: \*\*\* strong: inhibition of  $\geq$  70% \*\* moderate: inhibition 50-70% \* weak: inhibition  $<$  50%

4 **Identification of endophytic fungal isolates.** Identification of endophytic fungal isolate (BB1) was carried out molecularly based on partial genetic analysis at locus *Internal Transcribed Spacer (ITS) ribosomal DNA* fungi [14].

The sequencing data are then trimming and assembling using the SeqTrace program. The sequence data (contig) that has been assembled is then carried out by BLAST with a database that has been registered with the NCBI/National Center for Biotechnology Information (<http://ncbi.nlm.nih.gov/BLAST/>). This aims to determine taxa or species that have the greatest homology/similarity molecularly. The visualization uses the Treeview v1.4.3 program to design the phylogeny tree as shown in Fig 2. Based on the phylogeny tree, the endophytic fungus BB1 has highest homology with *Neopestalotiopsis surinamensis* strain CBS 450.74

**Isolation of secondary metabolite from BB1.** Ethyl acetate extract of BB1 (2 g) was separated by column chromatography using silica gel 60 (70-230 mesh). A total of 56 eluates were collected consisting of five fractions (F1-F5). Fraction F4 was purified by re-chromatography to obtain compound 2 (32 mg). Antibacterial activity test of compound 2 (at 400 µg/disk) against four bacteria showed that this compound had a moderate category of *E. coli* and *S. aureus*.

Compound 2 as white crystals. UV (MeOH)  $\lambda_{\max}$  270 nm; U<sub>33</sub> (MeOH + NaOH)  $\lambda_{\max}$  313 nm (bathochromic shift); IR (KBr)  $\nu_{\max}$   $\text{cm}^{-1}$ : 3265 (O-H), 3175 (C-H aromatic), 2922; 2851 (C-H aliphatic), 1661 (C=O), 1470 (C=C aromatic), and 6027 (C-O ester). The <sup>1</sup>H-NMR spectrum showed the presence of three signals at  $\delta_{\text{H}}$  4.42 (2H, s), 6.50 (1H, s), and 7.96 ppm (1H, s). This signal indicates that compound 2 has one methylene  $\text{sp}^3$  group and two methine  $\text{sp}^2$  groups. The <sup>13</sup>C-NMR spectrum exhibited the presence of six carbons namely one carbon  $\text{sp}^3$  and five carbon  $\text{sp}^2$ . The presence of two oxyaryl (= C-O) carbon atoms exhibited at  $\delta_{\text{C}}$  141.2 and 170.3 ppm. Three signals of carbonyl cyclic-ester at  $\delta_{\text{C}}$  176.9 ppm were assigned to lactone, which was indicated by the presence of oxyaryl carbon at  $\delta_{\text{C}}$  141.2 ppm. Further identification of carbon and protons signals was determined based on 2D NMR spectrum (Table 4). Based on the spectroscopic analysis and compared with literature [21], it was concluded that compound 2 as 5-hydroxy-4-hydroxymethyl-2H-pyran-2-one. Compound 2 has the molecular formula C<sub>6</sub>H<sub>6</sub>O<sub>4</sub>, with DBE = 4. Fig. 3 showed the HMBC correlation and  $\delta$ -assignment of compound 2.





Figure 2. The phylogenetic tree of BB1 with 1000 bootstrap replicates

**Table 4.** <sup>1</sup>H and <sup>13</sup>C NMR data of compound **2** recorded at <sup>1</sup>H-500 MHz, <sup>13</sup>C-125 Mhz, in CD<sub>3</sub>OD and 2\* recorded at <sup>1</sup>H-500 MHz, <sup>13</sup>C-125 Mhz, in CD<sub>3</sub>OD [21]

No. C	$\delta_c$ (ppm) <b>2</b>	$\delta_H$ (ppm), $\Sigma H$ , mult, $J$ (Hz) <b>2</b>	HMBC <b>2</b>	$\delta_c$ (ppm) <b>2*</b>	$\delta_H$ (ppm) <b>2*</b>
2	176.9			176.8	<b>63</b>
3	110.8	6.50 (1H; s)	61.1; 147.2; 170.3; 176.9	110.7	<b>6.49</b> (1H; s)
4	147.2			147.3	
5	170.3			170.4	
6	141.2	7.96 (1H; s)	147.2; 170.3; 176.9	141.0	<b>7.94</b> (1H; s)
7	61.1	4.42 (2H; s)	110.8; 170.3	61.2	<b>4.40</b> (2H; s)



**Figure 3.** The HMBC correlation and  $\delta$ -assignment of compound **2**

Previous studies have found an esterification derivative of compound **2** from *Andrographis paniculata* leaves, namely **3** (5-hydroxy-2-oxo-2H pyran-4-yl) methyl acetate). The inhibition zone in the antibacterial activity test of the compound is lower than that of compound **2**. The esterification of free hydroxyl groups might cause a decrease in antibacterial activity [22]. Thus it is known that free hydroxyl groups affect the antibacterial activity of a compound.

### Summary

Endophytic fungi extract of *Neopestalotiopsis surinamensis* has a higher antibacterial activity than host. Compound **1** had a moderate category against *E. coli*, and compound **2** had moderate activity against *E. coli* and *S. aureus*. The results of this study recommend that further research should be needed to combine compound **1** and **2** as additives to resistant antibiotics to determine their increased antibacterial activity.

### Acknowledgments

The authors thank to The Ministry of Research Technology and Higher Education (Kemenristekdikti) Republik Indonesia through Hibah Penelitian Dasar Unggulan Perguruan Tinggi (PDUPT) 2018-2019.



## References

- [1] E.A.A. Pinheiro, J.R.S. Pina, A.O. Feitosa, J.M. Carvalho, F.C. Borges, P.S.B. Marinho, A.M.R. Marinho, Bioprospecting of antimicrobial activity of extracts of endophytic fungi from *Bauhinia guianensis*, Rev. Argent. Microbiol. 49 (2017) 3-6.
- [2] M.R. Habib, M.M. Rahman, A. Mannan, A.H. Zulfiker, M.E. Uddin, M.A. Sayeed, Evaluation of antioxidant, cytotoxic, antibacterial potential and phytochemical screening of chloroform extract of *Phyllanthus acidus*, Int. J. Appl Biol. Pharm. 2 (2011) 420-427.
- [3] S.K. Biswas, A. Chowdhury, J. Das, U.K. Karmakar, M.C. Shill, S.Z. Raihan, Assessment of cytotoxicity, antibacterial activity and phytochemical screening of ethanol extract of *Phyllanthus Acidus* L. (Family: Euphorbiaceae) Bark, J. Appl. Pharm. Sci. 1 (2011) 112-114.
- [4] V. Kothari, S. Seshadri, *In vitro* antibacterial activity in seed extracts of *Manilkara zapota*, *Annona squamosa*, and *Tamarindus indica*, Biol. Res. 43 (2010) 165-168.
- [5] M. Kaneria, S. Chanda, Evaluation of antioxidant and antimicrobial properties of *Manilkara zapota* L. (chiku) leaves by sequential soxhlet extraction method, Asian Pac. J. Trop. Biomed. (2012) S1526-S1533.
- [6] S. Bhargavi, B. Kanakaiah, D.K. Sowmya, B. Ravi, S. Nama, An evaluation of the antibacterial activity of root extracts of *Manilkara Zapota* against *Staphylococcus Aureus* and *Escherichia Coli*, Int. J. Phytopharm. 4 (2013) 171-173.
- [7] J.P. Marcellano, A.S. Collanto, R.G. Fuentes, Antibacterial activity of endophytic fungi isolated from the bark of *Cinnamomum mercadoi*, Pharmacogn. J. 9 (2017) 405-409.
- [8] S.S. Sandhu, S. Kumar, R.P. Aharwal, Isolation and identification of endophytic fungi from *Ricinus communis* Linn and their antibacterial activity, Int. J. Res. Pharm. Chem. 4 (2014) 611-618.
- [9] M. Rajamanikyam, V. Vadlapudi, R. Amanchy, S.M. Upadhyayula, Endophytic fungi as novel resources of natural therapeutics, Braz. Arch. Biol. Technol. 60 (2017) 1-26.
- [10] P. Zhang, P. Zhou, L. Yu, An Endophytic taxol-producing fungus from *Taxus media*, *Cladosporium cladosporioides* MD2, Curr. Microbiol. 59 (2009) 227-232.
- [11] J. Zhao, L. Zhou, J. Wang, T. Shan, L. Zhong, X. Liu, X. Gao, Endophytic fungi for producing bioactive compounds originally from their host plants. Current Research, Technology and Education Topics in Applied Microbiology and Microbial Biotechnology, A. Mendez-Vilas (Ed). (2010) 567-576.
- [12] L. Calcul, C. Waterman, W.S. Ma, M.D. Lebar, C. Harter, T. Mutka, L. rton, P. Maignan, A.V. Olphen, D.E. Kyle, L. Vrijmoed, K.L. Pang, C. Pearce, B.J. Baker, Screening mangrove endophytic fungi for antimalarial natural products, Mar. Drugs. 11 (2013) 5036-5050.
- [13] S.K. Deshmukh, M.K. Gupta, V. Prakash, S. Saxena, Endophytic fungi: A source of potential antifungal compounds, J. Fungi. 4 (2018) 2-42.
- [14] Elfita, M. Muharni, M. Mardiyanto, F. Fitrya, E. Nurmawati, R. Simangunsong, Antibacterial activity of *Ageratum conyzoides* and their endophytic fungi extracts. Microbiology and Biotechnology Letter (2019) (In Press).
- [15] H. Liang, Y. Xing, J. Chen, D. Zhang, S. Guo, C. Wang, Antimicrobial activities of endophytic fungi isolated from *Ophiopogon japonicus* (Liliaceae), BMC. Complement Altern. Med. 12 (2012) 238-243.
- [16] E. Elfita, M. Munawar, M. Muharni, G. Pratiwi, R. Rahmadania, A benzoyl compound isolated from endophytic fungi of kandis gajah (*Garcinia griffithii*) and asam kandis (*Garcinia cowa*), Makara J. Sci. 20 (2016) 18-23.

- 
- [17] <sup>32</sup> J.P. Harley, L.M. Prescott LM, Laboratory Exercises in Microbiology, Fifth ed., The McGraw-Hill Companies, 2002.
- [18] <sup>5</sup> Y.C. Lim, R. Rajabalaya, S.H.F. Lee, K.U. Tennakoon, Q.V. Le, A. Idris, I.N. Zulkipli, N. Keasberry, S.R. David, Parasitic mistletoes of the genera *scurrula* and *viscum*: from bench to bedside, *Molecules* 21 (2016) 1-34.
- [19] <sup>24</sup> A. Gohar, S.R. Gedara, H.N. Baraka, New acylated flavonol glycoside from *Ceratonia siliqua* L. seeds, *J. Med. Plant. Res.* 3 (2009) 424-428.
- [20] <sup>16</sup> M.U. Amin, M. Khurram, B. Khattak, J. Khan, Antibiotic additive and synergistic action of rutin, morin and quercetin against methicillin resistant *Staphylococcus aureus*, *BMC. Complement Altern. Med.* 15 (2015) 59-71.
- [21] <sup>31</sup> E. Elfita, M. Munawar, M. Muharni, M.A. Sudrajat, Identification of new lactone derivatives isolated from *Trichoderma* sp., an endophytic fungus of brotowali (*Tinospora crispa*). *Hayati J. Biosci.* 21 (2014) 15-20.
- [22] <sup>1</sup> E. Elfita, M. Munawar, M. Muharni, I. Ivantri, Chemical constituents from an endophytic fungus *Aspergillus* Sp (Sbd5) isolated from sambiloto (*Andrographis paniculata* Nees), *Microbiol. Indones.* 9 (2015) 82-88.

# Antibacterial Activity of Traditional Medicine *Scurrula atropurpurea* (BL) DANS and their Endophytic Fungi

## ORIGINALITY REPORT

**29%**  
SIMILARITY INDEX

**24%**  
INTERNET SOURCES

**22%**  
PUBLICATIONS

**10%**  
STUDENT PAPERS

## PRIMARY SOURCES

**1** [scitepress.org](https://scitepress.org) 1%  
Internet Source

**2** [innovareacademics.in](https://innovareacademics.in) 1%  
Internet Source

**3** [www.abe.pl](http://www.abe.pl) 1%  
Internet Source

**4** Toga Pangihotan Napitupulu, Indriati Ramadhani, Atit Kanti, I Made Sudiana. "Diversity, phosphate solubilizing, and IAA production of culturable fungi associated with healthy and wilt banana", Archives of Phytopathology and Plant Protection, 2021 1%  
Publication

**5** Babra Moyo, Nikita Tawanda Tavengwa, Ntakadzeni Edwin Madala. "Diverse chemical modifications of the chlorogenic acid composition of *Viscum combreticola* Engl.: A premise for the state of readiness against excessive sunlight exposure", Journal of 1%

# Photochemistry and Photobiology B: Biology, 2022

Publication

---

6	<a href="http://www.ethnoleaflets.com">www.ethnoleaflets.com</a> Internet Source	1 %
7	Submitted to Universidad de Cádiz Student Paper	1 %
8	<a href="http://www.science.gov">www.science.gov</a> Internet Source	1 %
9	Submitted to Landmark University Student Paper	1 %
10	Rajeve Brooks, Andrea Goldson-Barnaby, Dennis Bailey. " Nutritional and Medicinal Properties of L. (Jimbilin) ", International Journal of Fruit Science, 2020 Publication	1 %
11	<a href="http://www.researchgate.net">www.researchgate.net</a> Internet Source	1 %
12	Dhandapani Ganapathy, Johannes R. Reiner, Guillermo Valdomir, Soundararasu Senthilkumar, Lutz F. Tietze. "Enantioselective Total Synthesis and Structure Confirmation of the Natural Dimeric Tetrahydroxanthone Dicerandrol C", Chemistry - A European Journal, 2017 Publication	1 %

---

- |    |  |     |
|----|--|-----|
| 13 | Garima Yadav, Mukesh Meena.<br>"Bioprospecting of endophytes in medicinal plants of Thar Desert: An attractive resource for biopharmaceuticals", Biotechnology Reports, 2021<br>Publication  | 1 % |
| 14 | <a href="http://prp.hec.gov.pk">prp.hec.gov.pk</a><br>Internet Source  | 1 % |
| 15 | <a href="http://www.scialert.net">www.scialert.net</a><br>Internet Source  | 1 % |
| 16 | Jonghoon Shin, Vasantha-Srinivasan Prabhakaran, Kwang-sun Kim. "The multi-faceted potential of plant-derived metabolites as antimicrobial agents against multidrug-resistant pathogens", Microbial Pathogenesis, 2018<br>Publication | 1 % |
| 17 | <a href="http://jeb.co.in">jeb.co.in</a><br>Internet Source  | 1 % |
| 18 | Hu, Jiang, Yan Song, Hui Li, Xia Mao, and Xiao-Dong Shi. "Diterpene glycosides from the ethanol extract of <i>Diplopterygium rufopilosum</i> ", Journal of Asian Natural Products Research, 2015.<br>Publication                     | 1 % |
| 19 | <a href="http://iopscience.iop.org">iopscience.iop.org</a><br>Internet Source  | 1 % |

20

coek.info  
Internet Source

1 %

21

Flávia Cristina Seabra Pires, Ana Paula de Souza e Silva, Ivonete Quaresma da Silva, Joicy Corrêa de Oliveira et al. "Application of supercritical technology in the production of dietary supplement based on plant extracts", Elsevier BV, 2021

Publication

1 %

22

Rosa Elvira Sánchez-Fernández, Rosalía Sánchez-Fuentes, Hiram Rangel-Sánchez, Simón Hernández-Ortega et al. "Antifungal and antioomycete activities and modes of action of isobenzofuranones isolated from the endophytic fungus Hypoxylon anthochroum strain Gseg1", Pesticide Biochemistry and Physiology, 2020

Publication

1 %

23

Éder de Vilhena Araújo, João Guilherme de Moraes Pontes, Stephanie Nemesio da Silva, Luciana da Silva Amaral et al. "The chemical warfare involved in endophytic microorganisms-plant associations", Elsevier BV, 2020

Publication

1 %

24

Saliha Besbas, Soumia Mouffouk, Hamada Haba, Laurence Marcourt, Jean-Luc Wolfender, Mohammed Benkhaled. "Chemical

1 %

composition, antioxidant, antihemolytic and anti-inflammatory activities of *Ononis mitissima* L.", *Phytochemistry Letters*, 2020

Publication

---

25

Aarthi Venugopalan, Smita Srivastava.  
"Endophytes as in vitro production platforms of high value plant secondary metabolites",  
*Biotechnology Advances*, 2015

Publication

---

1 %

26

[www.banglajol.info](http://www.banglajol.info)

Internet Source

---

<1 %

27

Khalid Mahmood Zia, Nadia Akram, Shazia Tabasum, Aqdas Noreen, Muhammad Usman Akbar. "Processing of bio-based polymers",  
*Elsevier BV*, 2021

Publication

---

<1 %

28

[arpnjournals.org](http://arpnjournals.org)

Internet Source

---

<1 %

29

[garuda.kemdikbud.go.id](http://garuda.kemdikbud.go.id)

Internet Source

---

<1 %

30

[www.bioscience.org](http://www.bioscience.org)

Internet Source

---

<1 %

31

[pms.unsri.ac.id](http://pms.unsri.ac.id)

Internet Source

---

<1 %

32

E. Elakkiya, Manickam Matheswaran.  
"Comparison of anodic metabolisms in

<1 %

bioelectricity production during treatment of dairy wastewater in Microbial Fuel Cell", Bioresource Technology, 2013

Publication

33

Hila Asham, Shahab Bohlooli, Donya Dostkamel, Sadaf Rezvanpoor, Saghi Sepehri. "Design, Synthesis, and Biological Screening for Cytotoxic Activity of Monastrol Analogues", Polycyclic Aromatic Compounds, 2021

Publication

<1 %

34

[biomedres.us](http://biomedres.us)

Internet Source

<1 %

35

Submitted to National University of Ireland, Galway

Student Paper

<1 %

36

Pushpak Mizar, Rekha Arya, Truc Kim, Soyoung Cha et al. " Total Synthesis of Xanthoangelol B and Its Various Fragments: Toward Inhibition of Virulence Factor Production of ", Journal of Medicinal Chemistry, 2018

Publication

<1 %

37

[jurnal.uns.ac.id](http://jurnal.uns.ac.id)

Internet Source

<1 %

38

[scholarcommons.usf.edu](http://scholarcommons.usf.edu)

Internet Source

<1 %



39

[123dok.com](http://123dok.com)

Internet Source

&lt;1 %

40

Marie, Benjamin, Jaison Arivalagan, Lionel Dubost, Sophie Berland, Arul Marie, and Frédéric Marin. "Unveiling the Evolution of Bivalve Nacre Proteins by Shell Proteomics of Unionoidae", *Key Engineering Materials*, 2016.

Publication

&lt;1 %

41

[scholarworks.bwise.kr](http://scholarworks.bwise.kr)

Internet Source

&lt;1 %

42

"Two New 5-Hydroxy-2-pyrone Derivatives Isolated from a Marine-derived Fungus *Aspergillus flavus*", *The Journal of Antibiotics*, 04/2008

Publication

&lt;1 %

43

[baadalsg.inflibnet.ac.in](http://baadalsg.inflibnet.ac.in)

Internet Source

&lt;1 %

44

[tjnpr.org](http://tjnpr.org)

Internet Source

&lt;1 %

45

K.N. Soujanya, R. Siva, P. Mohana Kumara, Amitava Srimany et al. "Camptothecin-producing endophytic bacteria from *Pyrenacantha volubilis* Hook. (Icacinaeae): A possible role of a plasmid in the production of camptothecin", *Phytomedicine*, 2017

Publication

&lt;1 %

46	M Y Farisa, K E Namaskara, M B Yusuf, Desrina. " Antibacterial Potention of Extract of Rotifers Fed with Different Microalgae to Control ", IOP Conference Series: Earth and Environmental Science, 2019 Publication	<1 %
47	Submitted to Universiti Sains Malaysia Student Paper	<1 %
48	<a href="http://media.neliti.com">media.neliti.com</a> Internet Source	<1 %
49	<a href="http://abisys.in">abisys.in</a> Internet Source	<1 %
50	<a href="http://akt.febi.uin-alauddin.ac.id">akt.febi.uin-alauddin.ac.id</a> Internet Source	<1 %
51	<a href="http://dokumen.pub">dokumen.pub</a> Internet Source	<1 %
52	<a href="http://eksakta.ppj.unp.ac.id">eksakta.ppj.unp.ac.id</a> Internet Source	<1 %
53	<a href="http://etd.repository.ugm.ac.id">etd.repository.ugm.ac.id</a> Internet Source	<1 %
54	<a href="http://ijpbs.com">ijpbs.com</a> Internet Source	<1 %
55	<a href="http://jebas.org">jebas.org</a> Internet Source	<1 %

56

[oamjms.eu](http://oamjms.eu)

Internet Source

&lt;1 %

57

[pubs.rsc.org](http://pubs.rsc.org)

Internet Source

&lt;1 %

58

[ujcontent.uj.ac.za](http://ujcontent.uj.ac.za)

Internet Source

&lt;1 %

59

[www.freepatentsonline.com](http://www.freepatentsonline.com)

Internet Source

&lt;1 %

60

Yao-Kouassi, Philomène A., Abdulmagid Alabdul Magid, Bernard Richard, Agathe Martinez, Marie-José Jacquier, Catherine Caron, Elisabeth Le Magrex Debar, Sophie C. Gangloff, Antoine A. Coffy, and Monique Zèches-Hanrot. "Isoflavonoid Glycosides from the Roots of *Baphia bancoensis*", *Journal of Natural Products*, 2008.

Publication

&lt;1 %

61

N. Venkateswarulu, P. VeeraBramha Chari, S.K. Thaslim basha, C. Nagaraju, T. Vijaya. "Isolation and purification of (E)-3- (2, 3-dihydroxyphenyl) acrylic acid from endophytic fungi *Fusarium equiseti* EF-32 and its anti-candidal and anticancer activities", *Biocatalysis and Agricultural Biotechnology*, 2017

Publication

&lt;1 %

62

Samiksha Sharma, Kishan Lal Tiwari, Shailesh Kumar Jadhav. "Diversity of fungal endophytes in *Typha latifolia* (L.) and their lead biosorption activity", Euro-Mediterranean Journal for Environmental Integration, 2017

Publication

---

<1 %

63

Sonda, S.. "Design and synthesis of orally active benzamide derivatives as potent serotonin 4 receptor agonist", Bioorganic & Medicinal Chemistry, 20030915

Publication

---

<1 %

---

Exclude quotes      On

Exclude matches      Off

Exclude bibliography      Off