

Potency of endophytic fungi from *Nauclea orientalis* L. as antioxidant producer

by Hary Widjajanti

Submission date: 21-Sep-2022 12:15PM (UTC+0700)

Submission ID: 1905177422

File name: 9_POTENCY_OF_ENDOPHYTIC_FUNGI_JURNAL_BERKALA_HAYATI_DES_2021.pdf (467.32K)

Word count: 4319

Character count: 23504

Potency of endophytic fungi from *Nauclea orientalis* L. as antioxidant producer

Elisa Nurnawati, Hary Widjajanti*, Vivi Hendra Sutandar, Maulida Harwati, Eka Amelia, Siti Alharzsa, Nina Tanzerina

33
Biology Department, Faculty of Mathematics and Natural Sciences, Sriwijaya University, Indonesia

Abstract

Nauclea orientalis L. is widely distributed in Indonesia. Secondary metabolites such as flavonoids, polyphenols, and saponins are abundant in the bark of *N. orientalis* L. These compounds have the potential to act as antioxidants. Endophytic fungi, through genetic transfer and coevolution, can produce the same metabolites as their host plant. As a result, understanding the potential of endophytic fungi from *N. orientalis* L. to produce antioxidant compounds that can be developed is critical. According to the findings of this study, twelve isolates have the potential to produce secondary metabolites with antioxidant properties. Three isolates had high antioxidant activity: DB2 was identified as *Aspergillus minisclerotigenes* with an IC₅₀ of 21 g/mL containing tannins, terpenes, and flavonoids, AB3 as *Colletotrichum perseae* with an IC₅₀ of 31 g/mL containing tannins and terpenoids, and AB1 as *Diaphorthe tulliensis* with an IC₅₀ of 48 g/mL containing tannins. The secondary metabolite group has the potential to be developed into an antioxidant agent.

27
words: Antioxidant, *Aspergillus minisclerotigenes*, *Colletotrichum perseae*, *Diaphorthe tulliensis*, Endophytic Fungi, *Nauclea orientalis* L.

Received: June 4, 2021 Revised: November 8, 2021 Accepted: November 9, 2021

Introduction

Based on culture and experience, plants have been widely used as ingredients in traditional medicine. It is due to the presence of active compounds that can be the main component of the drug. In Indonesia, *Nauclea orientalis* L. is popularly referred to as a genpol plant. In contrast, in Central Sulawesi, this plant is known as lokinda, and in Kalimantan and Sumatra, it is known as the bengkal (Tuheteru et al., 2014). Flavonoids, polyphenols, and saponins are abundant in the bark of *N. orientalis* L. (Saefudin and Basri, 2016). According to research by Wali et al. (2018), the stem base of *N. orientalis* L. contains 17 secondary metabolites. Six compounds at the middle of the stem and 19 at the ends. These secondary metabolites include phenols, phenolic acids, phenolic essential oils, terpenes, and tannins. Squalene (terpene), geraniol (terpene), stigmasta (terpene), and hexadecanoic acid (fatty acids) are some of the active compounds found in three parts of bark. Several secondary metabolites have antioxidant and anti-bacterial properties.

Endophytic fungi live intracellularly in healthy plant tissues and form symbiotic relationships with their hosts (Stone et al., 2000). Endophytic fungi produce a variety of secondary metabolites with diverse biological activities, including antimicrobial, anticancer, antioxidant, antituberculosis, antiparasitic, antiviral, immunomodulatory, and insecticide properties (Kaul et al., 2012). Through genetic transfer mechanisms,

46
endophytic fungi can produce the same secondary metabolites as their hosts (Rachman et al., 2018). According to Dao et al. (2015), an extract from the stem of *N. orientalis* L. had a significant ability to reduce DPPH free radicals as well as lipid peroxidase inhibitor activity. Alkaloids, terpenoids, simple phenolics, coumarin glucosides, anthraquinones, and lignans were among the compounds isolated.

Extraction methods with specific solvents are used to separate secondary metabolites from fungi. Many compounds found in polar solvents like methanol or ethanol have the potential to act as antioxidants (Do et al., 2013; Hameed et al., 2017). Smith et al. (2015) discovered that methanol extract of fungi was more effective as an antioxidant. Furthermore, the solvent most commonly used to extract the active compound is ethyl acetate, which is semipolar (Santoso et al., 2012). Because the use of large plant parts can reduce the number of plants, endophytic fungi can be used as an appropriate replacement for plant parts. As a result, this research looks into the antioxidant potential of metabolites from the endophytic fungus *N. orientalis* L.

Methods

Sampling

Bengkal (*Nauclea orientalis* L.) was taken in Kampung Sungai Pedada, Keramasan, Kertapati District, Palembang City, South Sumatra. Geographically, the sampling location is S 3°1'42.68352" and E 104.720194". The bengkal plant that were used as samples were the root, leaves and bark taken from the branching of the stem.

* Corresponding Author:

Hary Widjajanti

31
Biology Department, Faculty of Mathematics and Natural Sciences, Sriwijaya University, Jalan Palembang-Prabumulih, Km 32 Inderalaya Ogan Ilir 30662.

Phone/Fax :0711-580056

E-mail: haryunsri@yahoo.com

Isolation endophytic fungi

Each sample was washed with running water and cut into 2x1 cm pieces. All samples' surfaces were sterilized for 1 minute with 5% sodium hypochlorite (NaOCl) and then for 1 minute with 70% alcohol. Finally, the sample was rinsed twice with sterile distilled water for one minute each and dried on sterile filter paper. Each sample was aseptically placed on solid PDA (Potato Dextrose Agar) media in a petri dish and incubated at room temperature until fungal growth was observed. Purification was performed on fungal colonies that had grown and displayed various morphological characteristics (Kusumawardhani et al., 2015).

Cultivation and secondary metabolites extraction of endophytic fungi

Fungal isolates were inoculated on 500 mL PDB (Potato Dextrose Broth) medium and incubated at room temperature for 35 days under static conditions. Cultivation is carried out in duplicate. The fungal biomass was separated from the medium after incubation. For 24 h, the filtrate was extracted liquid-liquid (partition) with ethyl acetate (1:1). Finally, the solution is separated from the solvent by a rotary evaporator. The biomass was dried in a 40°C oven for 3 h before being weighed. The biomass was then ground and extracted for two days with methanol. The mixture was then filtered, and the solution was removed, yielding a thick extract (Hasiani et al., 2015; Tan et al., 2018).

Qualitative test of antioxidant activity

Each extract and ascorbic acid (control) were spotted on the TLC plate as much as ±5 µL extract and with ±1.5 cm gap between each isolates. To determine antioxidant activity was conducted by spraying TLC plate with DPPH 0.1 mM. Isolates selection were done with comparing the closest color spots of the isolates to yellow spot with purple background of ascorbic acid.

Antioxidant activity test with DPPH method

Variation of extract concentrations were made by solvent dilution with concentrations of 200, 100, 50, 25, 12.5, 6.25 and 0 µg/mL. Next, 0.1 mL concentration of sample solution was added with 3.8 mL of DPPH 0.1 mM solution. Then, the mixture was homogenized and let for 30 min in dark place. Absorbance then measured with spectrophotometer UV-Vis in λmaks 517nm. Standard antioxidant which was ascorbic acid as control positive with same treatment to the samples (Pratiwi et al., 2014). The percentage of inhibition of DPPH free radical activity was calculated using the equation (Mu'nisa et al., 2012):

$$\text{inhibition percentage} = \frac{A_k - A_s}{A_k} \times 100$$

A_k: absorbance control Positive control: ascorbic acid

A_s: absorbance sample Negative control: ethanol

IC₅₀ value then determine using the linear regression equation (Hasanah et al., 2017):

$$Y = a + bX$$

$$Y = 50$$

$$X = \text{sample concentration}$$

$$a = \text{gradients}$$

$$b = \text{constants}$$

TLC Analysis

Extracts were spotted on the TLC plate and eluted with n-hexane and ethyl acetate (1:2). The plate sprayed with H₂SO₄ 10%. Chromatogram pattern on the plate observed under UV lights. Based on Rusnaeni et al. (2016), R_f was calculated using the equation:

$$R_f = \frac{\text{compound distance from spot starting points}}{\text{solvent distance from spot starting points}}$$

Characterization and identification of fungi

Endophytic fungal isolates with high IC₅₀ values were inoculated on solid PDA, Malt Extract Agar (MEA), and Czapek Dox Agar and incubated for seven days at room temperature. The morphological characteristics of the fungus were examined based on the macroscopic and microscopic characters of each isolate and then identified based on Barnett and Hunter (1998). Molecular identification was performed by extracting DNA, amplifying the ITS region using ITS1 and ITS4 primers, then sequencing at Macrogen, Korea. Sequencing results were edited and merged using Chromas. MEGA X was used for the phylogenetic tree reconstruction.

Results

Isolation endophytic fungi

Based on the isolation and purification results, 12 fungi isolates were collected from the roots, bark, and leaves of *N. orientalis* L. (Table 1). Because the fungal isolates had different morphologies, the species were also different. Endophytic fungi are known to live in plant tissues, so that all isolates can be obtained from all plant organs.

Table 1. Endophytic fungal isolate isolated from *Nauclea orientalis* L.

No.	Sample	Code	Number of isolates
1.	Root	AB1	4
		AB2	
		AB3	
		AB4	
2.	Leaf	DB1	4
		DB2	
		DB3	
		DB4	
3.	Stem	BB1	4
		BB2	
		BB3	
		BB4	
Total			12

Cultivation and secondary metabolites extraction

Fungal secondary metabolites were successfully extracted from the filtrate and biomass (Table 2). Table 2 shows no correlation between fungal biomass and the

weight of ethyl acetate extract and methanol extract. Because the fungal isolates differed in species (Table 1), their growth patterns and metabolism differed as well.

Table 2. Result of Cultivation of Endophytic Fungi from Bengkulu *Nauclea orientalis* L.

No	Isolate Codes	Dried Biomass Weight (g)	Ethyl Acetate Extract Weight(g)	Methanol Extract Weight (g)
1.	BB4	3.77	0.43	0.36
2.	DB3	2.7	0.31	0.36
3.	AB2	4.3	0.22	0.68
4.	AB1	4.25	0.34	0.37
5.	BB3	3	0.76	0.81
6.	DB4	1.8	0.57	0.36
7.	AB4	3.03	0.55	1.65
8.	AB3	2.59	0.23	1.35
9.	DB1	5.03	0.33	1.44
10.	DB2	2.31	0.31	0.56
11.	BB2	2.56	0.31	0.38
12.	BB1	2.26	0.18	1.54

Antioxidant Activity of Secondary Metabolites Extract of Endophytic Fungi

The ability of fungal isolates as antioxidant producers were analyzed by two methods, qualitative and quantitative methods. Qualitative analysis needs to be done to select fungal isolates that potentially produce antioxidant compounds before proceeding with quantitative analysis using a spectrophotometer to obtain the IC_{50} value. Based on Figure 1, it can be seen that all extracts produced a yellow color on a purple background with different color intensities. This result indicated that the fungal isolates could produce antioxidant compounds qualitatively.

The weight of the extracts, both ethyl acetate and methanol, was unrelated to the antioxidant activities of the compounds contained in them. Table 2 shows that the isolate with the highest extract weight, isolate BB3 for ethyl acetate extract, has low antioxidant ability (Table 3). In contrast, the methanol extract of biomass isolate AB4 demonstrated a positive correlation between high antioxidant capacity and extract weight. This condition indicates that the class of active compounds is critical in secondary metabolite activity.

Qualitative analysis was conducted using spectrophotometry. The result of the spectrophotometric method is the percentage of inhibition. Then, we analyzed the percentage of inhibition to obtain the IC_{50} value. IC_{50} will show which extract had strong and very strong antioxidant activity.

According to the quantitative analysis in Table 3, the ethyl acetate extract of three isolates, AB1, AB3, and DB2, had very strong antioxidant activity when compared to ascorbic acid as a control. Isolate AB4 has




Figure 1. TLC profile extract of endophytic fungi (*N. orientalis* L.) eluted with ethyl acetate: n-hexane (2:1). (A. filtrate, B. Biomass, 1. BB4, 2. DB3, 3. AB2, 4. AB1, 5. BB1, 6. DB4, 7. AB4, 8. AB3, 9. DB1, 10. DB2, 11. BB2, 12. BB1, 13. Ethyl acetate(A) methanol(B), 14. Ascorbic acid.)


strong antioxidant abilities in both methanol and ethyl acetate extracts, whereas isolates BB2 and BB4 have strong antioxidant abilities only in ethyl acetate extracts. Except for isolate AB4, all biomass extracts of fungal isolates had weak antioxidant abilities - inactive based on the IC_{50} value. This condition is most likely caused by improper extraction of secondary metabolite compounds that act as antioxidants.

Thin layer chromatography secondary metabolites extract of endophytic fungi (*Nauclea orientalis* L.)

Based on Table 3, crude extract of BB4, AB1, AB4, AB3 and DB2 was analysed with TLC. Separation of the compound are conducted to know compound types with ability as antioxidant. Figure 2 showed several different color pattern on top of the TLC plate. The color differences are affected by different types of compound inside each extract hence when separated by eluted TLC it will separated.

Table 3. IC₅₀ Value Secondary Metabolite Extracts of Endophytic Fungi

No	Isolate Codes	Filtrate		Biomass	
		IC ₅₀ (µg/ml)	Activity	IC ₅₀ (µg/ml)	Activity
1.	BB4	75	Strong	1.08 x 10 ³	Inactive
2.	DB3	119	Moderate	1.58 x 10 ⁹	Inactive
3.	AB2	189	Weak	314	Inactive
4.	AB1	48	Very Strong	1.35 x 10 ⁹	Inactive
5.	BB3	220	Very Weak	3.61 x 10 ⁵	Inactive
6.	DB4	127	Moderate	776	Very Weak
7.	AB4	54	Strong	97	Strong
8.	AB3	32	Very Strong	885	Very Weak
9.	DB1	587	Very Weak	5,40 x 10 ³	Inactive
10.	DB2	21	Very Strong	667	Very Weak
11.	BB2	88	Strong	106	Moderate
12.	BB1	206	Very Weak	9.20 x 10 ⁴	Inactive
13.	Ascorbic acid (control)	17	Very Strong	17	Very Strong

**Figure 2.** Chromatogram of crude extract of endophytic fungi. (a. BB4, b. AB1, c. AB4, d. AB4 biomass, e. AB3, f. DB2, g. BB2. 1. Tannin, 2. Flavonoid, 3. Terpenoid, 4. Alkaloid)**Characterization and identification of potential fungi**

Based on the antioxidant activity test, three isolates were known to possess very strong potential. Fungal isolates were characterized at the macroscopic and microscopic levels. Figure 3 depicts the results of the description.

On PDA medium, a greyish white colony of AB1 isolate with a diameter of 4.5 cm, reverse colored in brown and white on the edges. Growth of an AB1 isolate on CDA, with delicate mycelium. Microscopic characteristics of isolate AB1 had several features that are thin and hyaline septate hyphae. A 100x magnification examination reveals the presence of hyaline and cylindrical conidia (Fig. 3A).

Colony AB3 isolate on PDA medium was white, with a diameter of 3.2 cm and a reverse color of white (Fig.

3B). On PDA, AB3 growth was moderate, but on CDA and MEA, the medium around the colony is yellow. The microscopic characteristics of isolate AB3 included hyaline septate hyphae and thin hyphae. Many conidia found with shape are cylindrical and hyaline.

On PDA medium, the colony color of DB2 isolate is green with a white shade and white reversed. On PDA, the colony diameter was 4.5 cm (Fig. 3C). Growth of DB2 isolates on CDA and MEA was slower than growth on PDA. Microscopic features of isolate DB2 at 100x magnification include hyaline septate hyphae, conidia visible and colored in green, conidia and conidiophores abundant and visible and conidia shaped in an ellipse. All isolates were incubated at room temperature for seven days. The identification of the three fungi isolates

was advanced using molecular data with ITS1 and ITS4 primers. BLAST analysis (Table 4) revealed that isolates of fungi AB1, AB3, and DB2 were recognized as

member of the genera *Diaporthe*, *Colletotrichum*, and *Aspergillus*, with an identity percentage greater than 98 percent

Table 4. BLAST analysis of ITS region fungal isolate

No	Fungal isolate	References species from Genbank	Identity (%)
1	AB1	<i>Diaporthe tulliensis</i> BRIP 62248a	99.65
		<i>Diaporthe tectonae</i> MFLUCC 12-0777	99.09
2	AB3	<i>Colletotrichum aenigma</i> ICMP 18608	100.00
		<i>Colletotrichum perseae</i> CBS 141365	99.65
3	DB2	<i>Aspergillus minisclerotigenes</i> strain CBS 116735	98.94
		<i>Aspergillus aflatoxiformans</i> isolate DTO 228-62	98.34




Figure 3. Fungal isolates with the potential to produce very strong antioxidant compounds. (A. AB1, B. AB3, C. DB2, Colonies on PDA incubated for 7 days. a. top, b. reverse. 1. Conidia, 2. Septate hyphae, 3. Conidiogenous cell, 4. Vesicle, 5. Conidiophore, 6. Chain of conidia.)

Figure 4 depicts the construction of a phylogenetic tree based on the ITS region of isolate fungi with very strong antioxidant activity. According to Figure 4, there are three large clusters, I, II, and III. The bootstrap analysis for these three clusters was 100 percent, indicating that this cluster was stable. Cluster I contains isolate AB3 which is a members of the genus *Colletotrichum*. In cluster I, isolate AB3 was closely related to *C. perseae* CBS 141365 and had a 99.65%

identity based on BLAST analysis (Table 4). Isolate AB1 was assigned to cluster II, the genus *Diaporthe*, and was found to be related to *D. tulliensis* BRIP 62248a. Cluster III belongs to the genus *Aspergillus*, which contains isolate DB2. *A. minisclerotigenes* CBS116735 is related to isolate DB2. Cluster III is a member of the *Aspergillus* section *Flavi*. The three isolates' morphological data revealed a match with each related species.




Figure 4. The phylogenetic tree of endophytic fungal isolates with reference strain based on the Maximum Likelihood algorithm with Kimura 2 parameters. The numbers in each branch indicate bootstrap

Discussion

Endophytic fungi from *N. orientalis* L. have been isolated in a total of 12 isolates. Differences in endophytic fungi colonizing plant tissues can be influenced by the plant tissues' ability to provide the fungi with the nutritional needs they require. Furthermore, it is influenced by fungi's ability to adapt to the environmental conditions of their host plants. Fungi [33](#) the nutrients supplied by the host plant as an energy source for growth and development (Sopialena et al., 2018). Because of this condition, the fungal isolates found in each plant differ (Murdiyah, 2017)

Each isolate of endophytic fungi differed in mycelium development, extract weight, and dry weight of biomass (Table 2). These variation occur because each isolate has a unique genome and physiological properties. Secondary metabolite production is controlled by specific genes in each species or group of fungi. Fungi in the Kingdom Ascomycota have more genes than other groups, such as polyketide synthase (PKS) and non-ribosomal protein synthase (NRPS). Secondary metabolites are typically species-specific and arise from

primary metabolic intermediates (Alurappa et al., 2018; Collemare et al., 2008).

The class of secondary metabolite produced by each species differs due to the secondary metabolites that are unique to each fungal species (Figure 2). The group of compounds produced by each fungus determines its antioxidant activity, so the difference in IC₅₀ value in Table 3 represents antioxidant ability. Six fungi, with high- very high antioxidant activity by Molyneux (2004) standards, were able to produce tannins and the majority of flavonoids. Both of the [35](#) groups of compounds are found in polyphenols, which are known to have antioxidant properties (Liu et al., 2009). Phenolic and flavonoids have antioxidants properties for anti-allergy [19](#) anti-inflammation (Huyut et al., 2017). According to Diaz et al. (2012), bioactive extracts of herbs with high phenolic compound and flavonoid content have very strong antioxidant activity. Tannin is a secondary metabolite that has astringent, antidiarrheal, antibacterial, and antioxidant properties (Desmiaty et al. 2008).

Terpenoid compounds are found in AB4 isolate, [6](#) which has strong antioxidant activity. Terpeneoids can act as antioxidants by scavenging free radicals and inhibiting lipid peroxidation (Graßmann, 2005).

Diaporthe tulliensis AB1 contains alkaloids as one of its constituents. Nuraini et al. (2019) discovered a class of endophytic fungi alkaloid compounds with antioxidant properties. *Diaporthe* sp. has been reported to produce phenolic compounds that are antioxidants (Tanapichatsakul et al., 2017), only in this study, there were no phenolic compounds but tannins and flavonoids. Further research is needed in using methanol as a solvent that potential to bind antioxidant compounds as well as optimization of the potential isolates.

Acknowledgement

The publication of this article was funded by DIPA of Public Service Agency of Universitas Sriwijaya 2020.SP DIPA-023.17.2.677515/2020, revision 01, on March 16, 2020. In accordance with the Rector's Decree Number : 0684/UN9/SK.BUK.KP/2020, on July 15, 2020.

References

- Alurappa R., Chowdappa S., Narayanaswamy R., Sinniah U.R., Mohanty S.K., Swamy M.K. 2018. Endophytic Fungi and Bioactive Metabolites Production: An Update. In: Patra J., Das G., Shin HS. (eds) *Microbial Biotechnology*. Springer, Singapore. https://doi.org/10.1007/978-981-10-7140-9_21.
- Barnett, H.L., Hunter, B.B. 1998. *Illustrated General of Imperfect America* : Burger Publishing Company. vi + 240 hlm.
- Collemare, J., Billard, A., Böhnert, H. U., Lebrun, M.-H. 2008. Biosynthesis of secondary metabolites in the rice blast fungus *Magnaporthe grisea*: the role of hybrid PKS-NRPS in pathogenicity. *Mycology Resources*. 112, 207–215. doi: 10.1016/j.mycres.2007.08.003
- Dao, P. T. A., Quan, T. L., Mai, N. T. T. 2015. Constituents of the Stem of *Nauclea orientalis*. *Natural Products Communications* 10(11):1901-1903.
- Desmiaty, Y., Ratih, H. Dewi, M.A., Agustín, R. 2008. Penentuan Jumlah Tanin Total pada Daun Jati Belanda (*Guazuma ulmifolia* Lamk) dan Daun Sambang Darah (*Excoecaria bicolor* Hassk.) Secara Kolorimetri dengan Pereaksi Biru Prusia. *Ortocarpus* 8: 106-109.
- Diaz, P., Jeong, S. C. Lee, S., Khoo, C., Koyyalamudi, S. R. Antioxidant And Anti-Inflammatory Activities Of Selected Medicinal Plants And Fungi Containing Phenolic And Flavonoid Compounds. *Chinese Medicine* 7(1):26-34.
- Do, Q.D., Angkawijaya, A.E., Tran-Nguyen, P.L., Huynh, L.H., Soetaredjo, F.E., Ismadi, S., Ju, Y.H. 2014. Effect of extraction solvent on total phenol content, total flavonoid content, and antioxidant activity of *Limnophila aromatica*. *Journal of Food and Drug Analysis*. 22(3): 296 – 302. <http://dx.doi.org/10.1016/j.jfda.2013.11.001>
- Graßmann, J. 2005. Terpenoids as Plants Antioxidants. *Vitamin and Hormon* 72:505-535.
- Hameed, A., Hussain, S.A., Yang, J.H., Ijaz, M.U., Liu, Q., Suleria, H.A.R., Song, Y., 2017. Antioxidants potential of the filamentous fungi (*Mucor circinelloides*). *Nutrient*. 9(10), 1101; <https://doi.org/10.3390/nu9101101>
- Hasanah, M., Maharani, B., Munarsih, E. 2017. Daya Antioksidan Ekstrak Dan Fraksi Daun Kopi Robusta (*Coffea robusta*) Terhadap Pereaksi Dpph (2,2-difenil-1-pikrilhidrazil). *Indonesian Journal of Pharmaceutical Science and Technology* 4(2):42-49.
- Hasiani, V. V., Ahmad, I., Rijai, L. 2015. Isolasi Jamur Endofit Dan Produksi Metabolit Sekunder Antioksidan Dari Daun Pacar (*Lawsonia inermis* L.). *Jurnal Sains dan Kesehatan* 1(4):146-153.
- Huyut, Z., Beydemir, S., Gulcin, I. 2017. Antioxidant and Antiradical Properties of Selected Flavonoids and Phenolic Compounds. *Biochemistry Research International* 2017:1-10.
- Kaul, S., Gupta, M. A., Dhar, M. K. 2012. Endophytic Fungi from Medicinal Plants: A Treasure Hunt for Bioactive Metabolites. *Phytochemistry Reviews* 11(4):487-505.
- Kusumawardani, Y., Liliek, S., Abdul, C. 2015. Potensi Antagonis Jamur Endofit Pada Tanaman Lada (*Piper nigrum* L.) Terhadap Jamur *Phytophthora capsici* Leionian Penyebab Penyakit Busuk Pangkal Batang. *Jurnal Hama dan Penyakit Tumbuhan*. 3(1) : 21-29.
- Liu S.C., Lin, J.T., Wang, C.K. 2005. Antioxidant properties of various solvent extracts from lychee (*Litchi chinensis* sonn.) flowers. *Food Chemistry*. 114:577-581.
- Mu'nisa, A., Wresdiyati, T., Kusunorin, N., Manalu, W. 2012. Aktivitas Antioksidan Ekstrak Daun Cengkeh. *Jurnal Veteriner* 13(3): 272-277.
- Murdiyah, S. 2017. Fungi Endofit Pada Berbagai Tanaman Berkhasiat Obat Di Kawasan Hutan Evergreen Taman Nasional Baluran Dan Potensi Pengembangan Sebagai Petunjuk Parktikum Mata Kuliah Mikologi. *Jurnal Pendidikan Biologi Indonesia* 3(1):64-71.
- Molyneux, P. 2004. The Use Of The Stable Free Radical Diphenylpicrylhydrazyl (DPPH) For Estimating Antioxidant Acti. *Journal of Science Technology* 26(2):211-219.
- Nuraini, F.R., Setyaningsih, R., Susilowati, A., 2019. Antioxidant Activity of Bioactive Compound Produced by Endophytic Fungi Isolated from Endemic Plant of South Kalimantan *Mangifera casturi* Kosterm. *International Conference on Biology and Applied Science (ICOBAS) AIP Conf. Proc.* 2120,080013-1–080013-9; <https://doi.org/10.1063/1.5115751> Published by AIP Publishing. 978-0-7354-1860-8/\$30.00
- Pratiwi, D. R., Maria, B., Partomuan, S. 2014. Lelutung Tokak (*Tabernaemontana macrocarpa* Jack) Sebagai Sumber Zat Bioaktif Antioksidan dan Antikanker. *Ilmu Kefarmasian Indonesia* 12(2):267-272.
- Rachman, F., Mubarik, N. R., Simanjuntak, P. 2018. Aktivitas Antioksidan Ekstrak Kapang Endofit *Ch.Gm.B3* Asal Ranting Kayu Manis (*Cinnamomum burmanni*). *Jurnal Bioteknologi dan Biosains Indonesia* 5(2):204-213.
- Rusnaeni, Sinaga, D. I., Lanuru, F., Payungallo, I. M., Ulfiani, I. I. 2016. Identifikasi Asam Mefenamat Dalam Jamu Rematik Yang Beredar di Distrik Heram Kota Jayapura, Papua. *Jurnal Pharmacy* 13(1):84-91.
- Saefudin., Basri, E. 2016. Potensi Antioksidan dan Sifat Sitotoksik Ekstrak Kulit Kayu Sembilan Jenis Tumbuhan di Taman Nasional Lore Lindu. *Jurnal Penelitian Hasil Hutan*. 34(2): 147-155.
- Santoso, J., Anwarayah, S., Rumiantin, R. O., Putri, A. P., Ukhty, N., Yoshie-Stark, Y. 2012. Phenol content, antioxidant activity and fibers profile of four tropical seagrasses from Indonesia. *Journal of Coastal Development* 15(2): 189-196.
- Smith, H., Doyle, S., Murphy, R. 2015. Filamentous fungi as source of natural antioxidants. *Food Chemistry*. 185:389 – 397.
- Sopialena, S., Suyadi, S., Sahil M, Nurdiana J. 2018. The diversity of endophytic fungi associated with *Piper nigrum* in the tropical areas: a recent study from Kutai Kartanegara, Indonesia. *Biodiversitas* 19: 2028-2034.
- Stone, J. K., Bacon, C. W., White, J. 2000. An Overview of Endophytic Microbes: Endophytism Defined, p 3-29. in Bacon, C. W. dan White, J. F. 2000. *Microbial Endophytes*. New York: Marcel Dekker Inc. 500 pp.
- Tan, X. M., Zhou, Y. Q., Zhou, X. L., Xia, X. H., Wei, Y., Li, L. H., Tang, H. Z., Yu, L. Y. 2018. Diversity And Bioactive Potential Of Culturable Fungal Endophytes Of *Diosma versipellis*; A Rare Medicinal Plant Endemic To China. *Science Report* 8(1):1-9.
- Tanapichatsakul, C., Monggoot, S., Gentekaki, E., Pripdeevech, P. 2017. Antibacterial and Antioxidant Metabolites of *Diaporthe* spp. Isolated from Flowers of *Melodorum fruticosum*. *Current Microbiology*. 75:476-483.
- Tuheteru, F. D., Kusmana, C., Mansur, I., Iskandar. 2014. Karakteristik Buah Dan Mutu Morfo-Fisiologis Benih Lonkida (*Nauclea orientalis* L.) Dari Habitat Alami Di Sulawesi Tenggara. *Jurnal Pemuliaan Tanaman Hutan* 8(3):152-160.
- Wali, M., Tuharea, M. S., Uar, N. I. 2018. Senyawa Kimia Kayu Marsegu (*Nauclea orientalis* L.). *Jurnal Agribisnis Perikanan* 11(2):70-74.

Potency of endophytic fungi from *Nauclea orientalis* L. as antioxidant producer

ORIGINALITY REPORT

18%

SIMILARITY INDEX

15%

INTERNET SOURCES

14%

PUBLICATIONS

9%

STUDENT PAPERS

PRIMARY SOURCES

1	Submitted to Universitas Brawijaya Student Paper	2%
2	www.atlantis-press.com Internet Source	1%
3	eprints.umm.ac.id Internet Source	1%
4	Suruchi Gupta, Sanjana Kaul, Baljinder Singh, Ram A. Vishwakarma, Manoj K. Dhar. "Production of Gentsyl Alcohol from <i>Phoma herbarum</i> Endophytic in <i>Curcuma longa</i> L. and Its Antagonistic Activity Towards Leaf Spot Pathogen <i>Colletotrichum gloeosporioides</i> ", <i>Applied Biochemistry and Biotechnology</i> , 2016 Publication	1%
5	Submitted to University of Nigeria Student Paper	1%
6	www.tandfonline.com Internet Source	1%

7	Diana Castro - Vazquez, Jessica Nayelli Sánchez - Carranza, Laura Alvarez, Karina Eurídice Juárez - Mercado et al. "Methyl benzoate and cinnamate analogs as modulators of DNA methylation in hepatocellular carcinoma", Chemical Biology & Drug Design, 2022 Publication	1 %
8	repository.upi.edu Internet Source	1 %
9	repository.uinjkt.ac.id Internet Source	1 %
10	www.myfoodresearch.com Internet Source	1 %
11	digilib.uin-suka.ac.id Internet Source	<1 %
12	Salni, Hanifa Marisa, Lili Arista Repi. "Antioxidant Activities Bioactive Compound of Ethyl Acetate Extracts from rose myrtle Leaves (Ait.) Hassk.) ", IOP Conference Series: Materials Science and Engineering, 2020 Publication	<1 %
13	academic.oup.com Internet Source	<1 %
14	ujcontent.uj.ac.za Internet Source	<1 %

15	www.ifrj.upm.edu.my Internet Source	<1 %
16	journal.unhas.ac.id Internet Source	<1 %
17	ir.lib.uth.gr Internet Source	<1 %
18	repositorio.unicamp.br Internet Source	<1 %
19	Pei Ni Chuah, Dhalini Nyanasegaram, Ke-Xin Yu, Rasny Mohamed Razik et al. " Comparative conventional extraction methods of ethanolic extracts of leaves on antioxidant activity and toxicity ", British Food Journal, 2020 Publication	<1 %
20	id.123dok.com Internet Source	<1 %
21	www.scitepress.org Internet Source	<1 %
22	Submitted to Manipal University Student Paper	<1 %
23	ejurnal.untag-smd.ac.id Internet Source	<1 %
24	hdl.handle.net Internet Source	<1 %

25	repository.ub.ac.id Internet Source	<1 %
26	repository.usd.ac.id Internet Source	<1 %
27	<p>Yushiro Fujii, Kenji Satake, Shingo Watada, Tung-Cheng Ho. "Re-examination of Slip Distribution of the 2004 Sumatra–Andaman Earthquake (Mw 9.2) by the Inversion of Tsunami Data Using Green’s Functions Corrected for Compressible Seawater Over the Elastic Earth", Pure and Applied Geophysics, 2021</p> Publication	<1 %
28	istrc-ab.org Internet Source	<1 %
29	<p>Ahsan Hameed, Syed Ammar Hussain, Muhammad Umair Ijaz, Samee Ullah, Zafarullah Muhammad, Hafiz Ansar Rasul Suleria, Yuanda Song. "Antioxidant activity of polyphenolic extracts of filamentous fungus <i>Mucor circinelloides</i> (WJ11): Extraction, characterization and storage stability of food emulsions", Food Bioscience, 2020</p> Publication	<1 %
30	<p>Chunyin Wu, Wei Wang, Xiaoqing Wang, Hamza Shahid, Yuting Yang, Yangwen Wang, Shengkun Wang, Tijiang Shan. "Diversity and</p>	<1 %

communities of culturable endophytic fungi from the root holoparasite *Balanophora polyandra* Griff. and their antibacterial and antioxidant activities", *Annals of Microbiology*, 2022

Publication

31

Hermansyah, Almunady T Panagan, Fatma, Susilawati. "Indigenous Yeast for Bioethanol Production", *Journal of Physics: Conference Series*, 2021

Publication

<1 %

32

Pramulani Mulya Lestari, Supandi, Mawar Nusafitri. "Ethanol Extract Emulgel 96% Brokatul Rice (*Oryza Sativa* L.) as Antioxidant", *IOP Conference Series: Earth and Environmental Science*, 2021

Publication

<1 %

33

biovalentia.mipa.unsri.ac.id

Internet Source

<1 %

34

"Recent Advancement in White Biotechnology Through Fungi", *Springer Science and Business Media LLC*, 2019

Publication

<1 %

35

ejurnal.bppt.go.id

Internet Source

<1 %

36

repository.nwu.ac.za

Internet Source

<1 %

37	repository.unmul.ac.id Internet Source	<1 %
38	"Microbial Biotechnology", Springer Science and Business Media LLC, 2018 Publication	<1 %
39	A Rahmadi, Yusriandi, M H Hanafi, Junaifid, Supriadi, D Setianingrum, A Dina Widyastuti, R Susilawati. "Physical, microbial and pesticide contaminations on fresh vegetable and fruit marketed in Samarinda-Indonesia", IOP Conference Series: Earth and Environmental Science, 2020 Publication	<1 %
40	Kandasamy Saravanakumar, Bhaskaran Sriram, Anbazhagan Sathiyaseelan, Xiaowen Hu et al. "Molecular identification, volatile metabolites profiling, and bioactivities of an indigenous endophytic fungus (<i>Diaporthe</i> sp.)", Process Biochemistry, 2021 Publication	<1 %
41	Melia Trianasta, Warsidah Warsidah, Mega Sari Juane Sofiana. "AKTIVITAS ANTIBAKTERI DARI FUNGI ENDOFIT <i>Caulerpa racemosa</i> ASAL PERAIRAN PULAU LEMUKUTAN (ANTIBACTERIAL ACTIVITY FROM ENDOPHYTIC FUNGI <i>Caulerpa racemosa</i> ORIGIN IN LEMUKUTAN ISLAN WATERS)",	<1 %

Indonesian Journal of Pure and Applied Chemistry, 2021

Publication

42 Weerasak Songoen, Lothar Brecker, Thitaree Yooboon, Vasakorn Bullangpoti, Wanchai Pluempanupat, Johann Schinnerl. "Ursane-type triterpenoids, steroids and phenolics from the stem bark and leaves of *Nauclea orientalis* (L.) L. (Rubiaceae)", *Biochemical Systematics and Ecology*, 2022
Publication

43 discovery.researcher.life
Internet Source

44 ejournal.forda-mof.org
Internet Source

45 epdf.pub
Internet Source

46 media.neliti.com
Internet Source

47 www.hindawi.com
Internet Source

48 "Neotropical Endophytic Fungi", Springer Science and Business Media LLC, 2021
Publication

49 Submitted to Lovely Professional University
Student Paper

Exclude quotes On

Exclude matches Off

Exclude bibliography Off