

IJMS-2017

by Ilmu Kelautan Unsri

Submission date: 10-Apr-2023 12:14PM (UTC+0700)

Submission ID: 2060285096

File name: IJMS-2017.pdf (351.05K)

Word count: 5094

Character count: 26158

Abundance of Phytoplankton in The Coastal Waters of South Sumatera

Riris Aryawati^{1,4*}, Dietrich G. Bengen², Tri Prartono² and Hilda Zulkifli³

¹Marine Sciences Department, Faculty of Mathematics and Natural Sciences, Sriwijaya University
Jl. Raya Palembang-Prabumulih Km.32 Indralaya, Sumatera Selatan, 30662 Indonesia

²Marine Sciences Department, Faculty of Fisheries and Marine Sciences, Bogor Agricultural University
Jl. Agatis, Kampus IPB Dramaga, Bogor, 16680 Indonesia

³Biology Department, Faculty of Mathematics and Natural Sciences, Sriwijaya University
Jl. Raya Palembang-Prabumulih Km.32 Indralaya, Sumatera Selatan, 30662 Indonesia

⁴School of Post Graduate Studies, Bogor Agricultural University, Bogor
Jl. Agatis, Kampus IPB Dramaga, Bogor, 16680 Indonesia
Email: ririsaryawati76@gmail.com

Abstract

Phytoplankton in the ocean has an important role in forming the base of food chain responsible in primary production. Its abundance and number of phytoplankton species will indirectly affect the level of water's fertility. This study aimed to determine the presence of phytoplankton as bio-indicators of water quality in terms of abundance, diversity index, uniformity index, and saprobic coefficient in coastal waters of South Sumatera. The study was conducted at ten stations during high and low tides. Phytoplankton samples were taken vertically using plankton nets, cone-shaped with a diameter of 30 cm, length 100 cm and mesh size 30 μm . The study found 41 genera of phytoplankton, consisted of family Bacillariophyceae (26 genera), Dinophyceae (7 genera) Cyanophyceae (7 genera) and Chlorophyceae (1 genus). The highest number of genera was recorded while low tides in November (24 genera), and the lowest was on May while high tides (16 genera). The highest abundance of phytoplankton was recorded in August during high tides ($2,68 \times 10^7 \text{ cell.m}^{-3}$), and the lowest was in May during high tides ($6,59 \times 10^5 \text{ cell.m}^{-3}$). The diversity index (H'), the uniformity index (E), and the dominance index (D) ranged between 0,64–3; 0,15–0,71 and 0,15–0,83, respectively.

Keywords: phytoplankton, abundance, diversity index, South Sumatera coast.

Introduction

The coast of South Sumatera receives water from large rivers, such as Musi and Banyuasin, and several other small rivers. In this region, from its upstream to downstream, communities use water for various activities such as for factories (fertilizer, rubber, oil), settlements, fishing, aquaculture, agriculture, farming, transportation and various other activities. These activities will increase the input of waste in the form of agricultural, domestic and industrial wastes. This may affect the life of organism in the water, one of them is phytoplankton, the primary producer in the trophic level of food chain in water.

Plankton, especially phytoplankton, has a particular important role in the food chain in aquatic ecosystems and is often used as indicators of stability, fertility and water quality. Phytoplankton can play a role as one of the ecological parameters, aquatic fertility, in describing the condition of a water body (Richardson, 2008).

Increasing activities in the coastal waters of South Sumatera will affect on the distribution abundance and diversity of phytoplankton, which also influence the water quality this region. According to Nybakken (1992) and Nontji (2005), aquatic organisms can be used as indicators of water quality because of their habitat, mobility, and age (relatively long) in inhabiting a water area. Basmi (2010) stated that the determination of aquatic biological index conducted generally from analysis study of phytoplankton community because they have high species diversity and abundance. Some previous phytoplankton studies have been done in this area, however, it was just done in certain month only, and do not represent the seasonal pattern of phytoplankton (Aryawati et al., 2005; Isnaini, 2011; Surbakti et al., 2011). This study aimed to assess the seasonal abundance and diversity of phytoplankton, and to evaluate the water quality of South Sumatera coastal waters through biological analysis study in terms of phytoplankton community as bio-indicator of water quality.

³*) Corresponding author
© Ilmu Kelautan, UNDIP

Materials and Methods

Study sites

The study was conducted in the Banyuasin waters (South Sumatera) at 10 stations during high and low tides (Figure 1).

Phytoplankton sampling

Referring to Aquino *et al.* (2010) and Mulyani *et al.* (2012), phytoplankton sampling was conducted by using plankton nets with mesh size 30 μm , net mouth diameter 30 cm and net length 100 cm long. Sampling was done vertically from the depth of 2 meters to the surface of water. The filtrate, collected in the bucket, then was put into a sample bottle of 250 mL and was added with preservative, 4% formaldehyde (Edler and Elbrachter, 2010). All samples were stored in a cool place and protected from direct light beam to prevent color changing to the solution which can damage the sample.

Analysis of samples

Samples were collected and identified in the laboratory using a high power microscope (Nikon Eclipse E400). The phytoplankton were identified to genus level by using the identification guide of Newell and Newell (1977); Yamaji (1984) and Tomas (1997). Phytoplankton enumeration was done by using the Sedgwick-Rafter Counting Cell on sample fraction, and the results are expressed in cell.m^{-3} . Analysis of phytoplankton abundance refers to the formulation used by APHA (1992). The species diversity of phytoplankton is known by the Shannon and Weaver's equation (1963). Diversity index is a mathematical representation or illustration depicting the structure of communities, and it can ease the analysis of data for the number and kinds of organisms. Diversity shows the number of gene variations or species in a body of water.

Uniformity index is used to determine the distribution pattern of biota. If the value is relatively high, the existence of any biota in the water is in even condition. Dominance index is used to determine the existence of a certain species dominance in the water. Uniformity index ranges between 0 to 1, the smaller the value of uniformity (near zero) indicates that the spread of each member from each species is not even. Conversely, if the uniformity value is greater (close to 1), the population shows uniformity (number of individuals of each genus can be said to be the same or not much different) (Odum, 1998). The uniformity index and dominance index of phytoplankton are calculated using the formula from Odum (1998). The

dominance index values range between 0 to 1, where the smaller value indicates that there are no species that dominate, otherwise, the greater value shows the presence of certain species that dominate (Odum, 1998).

Results and Discussion

Species and abundance of phytoplankton

A species composition can illustrate the diversity or the number of species in a community. This kind of diversity can increase if the community becomes more stable, and otherwise diminished when the environment is unstable or crashed. The identified phytoplankton populations in Banyuasin water were 41 genera. The composition of the genus was greatly found in the family Bacillariophyceae, then Dinophyceae, Cyanophyceae, and Chlorophyceae. The composition was dominated by 26 genera of diatom group, 7 genera of Dinophyceae, 7 genera of Cyanophyceae, and 1 genus of Chlorophyceae (Table 1).

At each month of observation, while on high tide or low tide, Bacillariophyceae was found to have the largest number of genera and abundance than other groups. This is because the class Bacillariophyceae is able to adjust to a diverse environmental condition compared to other classes. According to Arinardi *et al.* (1997), the class Bacillariophyceae adapted well to the environmental conditions, has high tolerance and adaptability, thus, this class is cosmopolitan. This situation is often found in Indonesian waters and territorial water in other regions (Haumahu, 2004; Fathi and Al-Kahtani, 2009; Rokhim *et al.*, 2009; Ismunarti, 2013; Thoha and Aryawati, 2014).

Abundance is a measure of the number of individuals per unit volume of water which generally expressed in cells per unit volume. In general, the abundance of phytoplankton is closely linked to fertility or productivity of a waterbody. When the phytoplankton abundance is high, the water tend to have high productivity (Raymont, 1963).

Banyuasin's abundance of phytoplankton ranged between $6,59 \times 10^5$ to $4,21 \times 10^7$ cell.m^{-3} . In general, the abundance of phytoplankton was found slightly higher during low tide than high tide (Figure 2). The greatest abundance in every month of observation was found in group Bacillariophyceae then Dinophyceae, Cyanophyceae and Chlorophyceae. Genus *Chaetoceros* and *Skeletonema*, part of Bacillariophyceae family, were found to be the most abundant. Arinardi *et al.* (1997) stated that Bacillariophyceae members have




Figure 1. Map of study sites in the coastal waters of South Sumatera

higher reproducibility compared to other phytoplankton.

In August, *Chaetoceros* and *Skeletonema* gave the highest contribution to the phytoplankton abundance, so that the abundance of phytoplankton is commonly highest in this month. Both genera are often found in Banyuasin waters, and allegedly they play an important role there (Aryawati *et al.*, 2005; Isnaini, 2011; Surbakti *et al.*, 2011). In some waters in Indonesia both genera are also often found in large quantities (Soedibjo, 2006; Thoha, 2013; Thoha and Aryawati, 2014). Soedibjo (2007) showed that Jakarta Bay waters also showed a high abundance of *Skeletonema* and *Chaetoceros*, caused by the high level of silicate, supplied from land through the rivers, supported also by temperature and salinity levels that exist. The genera are classified as phytoplankton commonly found especially in seasons with low rainfall (Praseno and Sugestiningih, 2000).

Phytoplankton diversity

The results of the analysis for diversity index (H') uniformity (E) and dominance (D) in the coastal waters of South Sumatera (Table 2 and Figure 3)

showed that, in general, the waters belong to the less stable ones. In the months when certain water conditions are very good in terms of phytoplankton community structure (May high and low tides; November high; February high and low tides). In other months (August high and low tides; November low tide) showed unfavorable water conditions. Based on calculations, it was found that, in general, the month of May has excellent water conditions based on phytoplankton community structure, and the roughest water conditions was found in August. The same research results presented by Isnaini (2011) who conducted the research at Banyuasin river's estuary, which indicated that the phytoplankton community structure was in a better shape at transitional season (May) than the east season (July). These conditions were thought to be caused by environmental factors, existed at those times. In May, water conditions are suspected to be very conducive to the growth and proliferation of various types of phytoplankton, so that, a lot of phytoplankton can grow and proliferate, thus making species richness and abundance to increase. Differently, in August, supposedly the environmental conditions are not conducive to the growth and development of various types of phytoplankton, so that, only certain species can adapt, grow and

Table 1. Abundance of phytoplankton in the coastal waters of South Sumatera

No	Phytoplankton (cell.m ⁻³)	May		August		November		February	
		High tide	Low tide	High tide	Low tide	High tide	Low tide	High tide	Low tide
Cillariophyceae									
1	<i>Amphora</i>	0	0	0	0	0	0	4.247	708
2	<i>Asterionella</i>	0	0	0	46.450	0	0	0	0
3	<i>Bacillaria</i>	0	0	0	0	0	0	37.511	0
4	<i>Bacteriastrium</i>	22.650	4.954	104.388	50.244	7.786	38.556	21.232	15.571
5	<i>Chaetoceros</i>	206.658	141.558	15.582.485	2.628.478	94.140	150.383	60.158	48.845
6	<i>Coscinodiscus</i>	3.539	9.909	144.021	43.171	10.617	1.416	51.666	28.318
7	<i>Dictyosolen</i>	0	0	1.415	0	2.123	9.909	0	0
8	<i>Diploneis</i>	0	0	0	2.123	0	0	0	0
9	<i>Dytilum</i>	9.201	10.618	56.971	37.734	2.123	3.539	8.493	4.247
10	<i>Eucampia</i>	0	0	18.047	16.985	0	24.065	0	0
11	<i>Guinardia</i>	708	0	12.385	9.435	14.156	22.994	1.416	2.124
12	<i>Hemidiscus</i>	0	1.416	1.415	0	0	0	0	0
13	<i>Hemiaulus</i>	0	1.416	147.913	0	12.033	4.955	2.831	0
14	<i>Lauderia</i>	0	0	171.622	212.058	8.494	28.307	2.123	0
15	<i>Leptocylindrus</i>	0	0	4.246	46.685	4.246	0	6.370	0
16	<i>Mellosira</i>	0	8.494	0	0	0	0	0	0
17	<i>Nitzschia</i>	27.602	81.408	164.545	145.530	54.150	38.924	67.236	31.150
18	<i>Odontela</i>	2.123	1.416	67.941	27.125	2.124	1.416	18.402	4.955
19	<i>Pleurosigma</i>	0	0	1.415	0	0	0	708	0
20	<i>Planktoniella</i>	0	0	0	0	0	2.831	0	0
21	<i>Podocystis</i>	0	0	0	0	1.770	28.325	0	0
22	<i>Rhizosolenia</i>	6.370	9.202	115.005	79.245	22.650	9.201	21.233	14.866
23	<i>Skeletonema</i>	175.498	147.225	8.686.175	38.174.782	207.738	1.839.466	627.769	327.809
24	<i>Surirella</i>	708	3.539	0	0	708	2.831	6.370	15.576
25	<i>Synedra</i>	0	0	0	0	0	0	708	0
26	<i>Thalassiosira</i>	67.948	133.807	409.416	251.446	53.091	91.294	323.444	101.222
27	<i>Thalassiothrix</i>	16.280	48.853	1.012.039	169.329	16.634	18.747	352.464	55.921
Dinophyceae									
28	<i>Alexandrium</i>	0	0	2.831	0	0	0	0	0
29	<i>Ceratium</i>	3.538	0	0	0	0	2.477	10.616	4.247
30	<i>Dictyocha</i>	0	0	1.415	0	0	0	0	0
31	<i>Dinophysis</i>	2.831	1.416	2.123	9.908	9.910	9.201	708	0
32	<i>Noctiluca</i>	0	0	0	0	194.289	203.111	0	0
33	<i>Prorocentrum</i>	0	0	0	0	0	708	0	2.833
34	<i>Protoperidinium</i>	18.403	3.539	16.631	12.263	5.663	10.263	2.831	708
Cyanophyceae									
35	<i>Anabaena</i>	0	19.818	74.310	99.750	0	0	0	0
36	<i>Holopedium</i>	0	0	0	0	0	89.174	0	0
37	<i>Microcystis</i>	0	0	0	0	0	0	0	25.484
38	<i>Nostoc</i>	0	0	0	0	1.062	0	0	0
39	<i>Oscillatoria</i>	94.845	82.119	0	0	0	11.324	372.272	156.417
40	<i>Spirulina</i>	0	708	0	21.933	0	0	0	0
Chlorophyceae									
41	<i>Staurastrum</i>	0	0	0	0	0	0	0	708
Abundance (individual)		658.904	711.415	26.798.755	42.084.673	725.507	2.643.417	2.000.806	841.707
Genera		16	19	23	20	21	25	23	19




Figure 2. The abundance of phytoplankton in the coastal waters of South Sumatera (high tide: $6,59 \times 10^5$ cells.m⁻³ on May, $2,7 \times 10^7$ cells.m⁻³ on August, $7,26 \times 10^6$ cells.m⁻³ on November, 2×10^6 cells.m⁻³ on February 3rd; low tide; $7,11 \times 10^5$ cells.m⁻³ on May, $4,21 \times 10^7$ cells.m⁻³ on August, $2,64 \times 10^6$ cell.m⁻³ on November, $8,42 \times 10^5$ cells.m⁻³ on February) **Note.** ■ : high tide, ■ : low tide




Figure 3. The structure of the phytoplankton community in the coastal waters of South Sumatera **Note.** ▨ : Diversity (H'), ■ : Uniformity (E), ▩ : Dominance (D)

Table 2. Phytoplankton community structure in the coastal waters of South Sumatera

Month of Observation	May		August		November		February	
	High tide	Low tide	High tide	Low tide	High tide	Low tide	High tide	Low tide
Diversity (H')	2,72	3,00	1,58	0,64	2,99	1,93	2,81	2,85
Uniformity (E)	0,68	0,71	0,35	0,15	0,68	0,42	0,62	0,67
Dominance (D)	0,15	0,15	0,44	0,83	0,18	0,50	0,19	0,21

develop properly in those conditions. Moreover, in terms of phytoplankton which usually be found, *Chaetoceros* and *Skeletonema* are found to be

dominated. This happens because the member of these groups are able to survive in extreme conditions. According to Basmi (2010), the members

of phytoplankton from division Chrysophyta (class Bacillariophyceae) are cosmopolitan, have properties that easily adapt to the environment, resistant to extreme conditions and high reproduction.

Widiarti (2000) research results indicated that *Skeletonema* was positively correlated with phosphate in waters, while *Chaetoceros* like low salinity, and *Pyrodinium* like salinity and high nitrate content. Lagus *et al.* (2004) reported that diatoms (*Chaetoceros wighamii* and *Skeletonema costatum*) have a very quick response to the addition of nutrients, thus become dominant groups. A research conducted by Hasani *et al.* (2012) showed that there is a strong positive correlation between the concentration of nutrients N and P with the potential emergence of phytoplankton at different aquaculture locations in Lampung Bay. Pednekar *et al.* (2012) suggest that the abundance of phytoplankton species was associated with the increased nutrients that enter coastal waters from the mouth of river. No less important, the factor that triggering algal bloom is the reduced predation by herbivores (Lindahl and Dahl, 1990). Conditions such as these were suspected to affect the number of species and abundance of phytoplankton in the coastal waters of Southern Sumatera, thereby affecting phytoplankton community structure which ultimately affected water conditions in general.

Water quality conditions in the coastal water South Sumatera

Based on the results of measurements of water parameters in the study area, the sea water temperature ranged from 27,6 to 31,9; salinity from 0,09 to 33,05; pH 6,0 to 8,5; dissolved oxygen from 3,49 to 8,22 mg.L⁻¹; phosphate 0,020 to 1,110 mg.L⁻¹; nitrates from 0,64 to 3,98 mg.L⁻¹, silicate from 0,001 to 0,430 mg.L⁻¹. Environmental parameter values in the coastal waters of the South Sumatera, is generally considered to be still feasible for the growth of phytoplankton.

The temperatures of the sea water in this study were generally higher in the dry and rainy season but decreased in transition. The results of measurements in these waters showed that the value of the parameter in sea water temperature is still support for the growth of phytoplankton. The pH value in these waters has similarities with some of the of Indonesia waters (Soedibjo, 2006; Prianto *et al.*, 2013). Generally, pH in ocean and coastal waters ranging from 7,70 to 8,4.

DO levels measured in this study ranged from 3,49 to 8,22 mg.L⁻¹. According to Wardhana (1995), this range is still enough to support life aquatic

organisms normally because it has a value of more than 2 mg.L⁻¹. These results also have similarities with studies conducted in other Indonesia waters (Sembiring *et al.*, 2012). One of the factors that may affect the levels of dissolved oxygen in seawater is the inclusion of waste which needed much oxygen in its decaying process. This type of waste usually comes from people's activities.

Direction and speed of currents in these waters varied from the mouth of the river up to the offshore. Speed of currents at high tide or low tide, has a value of 0,009-1,141 m.s⁻¹. The variations of water flow affects to the variations in the distribution of phytoplankton. Romimohtarto and Juwana (2004) states that the very small size of phytoplankton resulted on its movement much depend on the movement of water.

Nitrate concentrations obtained from the measurement results on the study sites showed a high value. Based on the KepMenLH No. 51-2004, the quality standard of sea water nitrate concentrations eligible to marine life is 0,008 mg.L⁻¹. The high nitrate in these waters is likely to be influenced by the input mass of water from several rivers that carry many nutrients nitrate. This is similar to other research done Risamasu and Prayitno (2011) that found high nitrate concentrations exceeding the threshold value of the exposure limit (standard quality) in Matasiri Islands waters, South Kalimantan. The high nitrate concentrations, indicating the enrichment of nitrogen by the influence of the mass of water enter the Barito River that carries nitrogen compounds from land to sea. High concentrations values of nitrate exceeding the threshold quality standards has also been demonstrated by Isnaini (2014) in the waters around the Maspari Island, Ogan Ilir, South Sumatra (1-2,1mg.L⁻¹) and Handoko *et al.* (2013) in Karimunjawa, Central Java (0,108-1,595 mg.L⁻¹).

Phosphate concentrations obtained in this study had a high value and exceeds the quality standard limits as defined in KepMenLH. KepMenLH No. 51-2004 mentioned that the quality standard of maximum concentration of phosphate eligible to marine life is 0,015 mg.L⁻¹. Compared with some research in the other territorial waters of Indonesia, the concentration of phosphate in this region is higher than the Takalar waters, South Sulawesi ranged from 0,319 to 0,336 mg.L⁻¹ (Pirzan & Pong-Masak, 2008), and Kwanyar waters, Bangkalan from 0,05 to 0,96 mg.L⁻¹ (Rokhim *et al.*, 2009). Otherwise, it has lower value when compared with Karimunjawa Islands, Central Java from 1,769-4,030 mg.L⁻¹ (Handoko *et al.*, 2013).

12

The content of silicate in the 12 coastal waters of South Sumatera during the study ranged from 0,001 to 0,430 mg.L⁻¹. These results are similar to studies conducted by Tambaru (2008) in the coastal waters of Maros, South Sulawesi, but lower than researched Simanjuntak (2007) in Jakarta Bay. Temporally, the concentration of silicate in the coastal waters of South Sumatra was higher at the end of the dry season until the beginning of the rainy season. These results are similar to studies conducted by Lukman *et al.* (2014) in the coastal waters of South Sulawesi. According to the seasons, the concentration of silicate in the dry season is lower than the rainy season. Generally, the environmental conditions in the coastal waters of South Sumatra is still support for the growth and distribution of phytoplankton.

Conclusion

The study found 41 genera of phytoplankton, consisted of family Bacillariophyceae (26 genera), Dinophyceae (7 genera) Cyanophyceae (7 genera) and Chlorophyceae (1 genus). The highest number of genera was recorded while low tides in November (24 genera), and the lowest was on May while high tides (16 genera). The highest abundance of phytoplankton was recorded in August during high tides ($2,68 \times 10^7$ cell.m⁻³), and the lowest was in May during high tides ($6,59 \times 10^5$ cell.m⁻³). The diversity index (H'), the uniformity index (E), and the dominance index (D) ranged between 0,64-3; 0,15-0,71, and 0,15-0,83, respectively.

Acknowledgement

The authors grateful to Directorate General of Higher Education for financially supporting this work. We also wish to thank to Heron Surbakti, M.Si and Isnaini, M.Si for their full support to this research. Additionally, we wish to thank Adriani Sunuddin, M.Si and Suhartono, M.Si for manuscript correction.

References

- American Public Health Association (APHA), American Water Works Association, Water Pollution Control Federation. 1995. Standard methods for the examination of water and wastewater. APHA, AWWA, WPCF. 15th edisi.
- Aquino, J., Flores, B. & Naguit, M. 2010. Harmful Algal Bloom Occurrence in Murcielagos Bay Amidst Climate Change. *E-International Scientific Research J.* 2(4):358-365.
- Arinardi, O.H., Sutomo, A.B., Yusuf, S.A., Trimaningsih, Asnaryanti, E. & Riyono, S.H. 1997. *Kisaran Kelimpahan dan Komposisi Plankton Predominan di Perairan Kawasan Timur Indonesia*. Pusat Penelitian dan Pengembangan Oseanologi Lembaga Ilmu Pengetahuan Indonesia. Jakarta. 140 p.
- Aryawati, R., Surbakti, H. & Ulqodry, T.Z. 2005. Hubungan Kondisi Oseanografi dengan Kelimpahan Fitoplankton di Perairan Banyuasin. Pemanfaatan dan Pengelolaan Perairan Umum secara Terpadu bagi Generasi Sekarang dan Mendatang. 2004 Juli 27-29. BRPPU. Palembang.
- Basmi, J. 2010. *Planktonologi*. Fakultas Perikanan dan Ilmu Kelautan IPB. Bogor. 110 p.
- Edler, L & Elbrachter, M. 2010. The utermohl method for quantitative phytoplankton analysis. In: Karlson, B., C. Cusack and E. Bresnan (Eds.) *Microscopic and molecular methods for quantitative phytoplankton analysis*. Intergovernmental Oceanographic Commission, United Nations Educational, Scientific and Cultural Organization. Spain. pp. 13-15.
- Fathi, A.A. & Al-Kahtani, M.A. 2009. Water Quality and Planktonic Communities in Al-Khadoud Spring, Al-Hassa, Saudi Arabia. *American J. Environ. Sci.* 5(3):434-443.
- Handoko, Yusuf, M. & Wulandari, S.Y. 2013. Sebaran nitrat dan fosfat dalam kaitannya dengan kelimpahan fitoplankton di Kepulauan Karimunjawa. *Bul. Oseanografi Mar.* 2:48-53
- Hasani, Q., Adiwilaga, E.M. & Pratiwi, N.T.M. 2012. The relationship between the harmful algal blooms (habs) phenomenon with nutrients at shrimp farms and fish cage culture sites in pesawaran district Lampung Bay. *Makara J. Science.* 16(3): 183-191.
- Haumahu, S. 2004. Distribusi spasial fitoplankton di Teluk Ambon bagian dalam. *Ichtyos.* 3:91-98.
- Hutabarat, S., Soedarsono, P. & Cahyaningtyas, I. 2013. Studi analisa plankton untuk menentukan tingkat pencemaran di muara Sungai Babon Semarang. *J. Manag. Aquatic Resources.* 2(3):74-84.
- Ismunarti, D.H. 2013. Analisis komponen utama pada hubungan distribusi spasial komunitas fitoplankton dan faktor lingkungan. *Ilmu Kelautan.* 18(1):14-19.

- Isnaini. 2011. Fitoplankton di perairan Muara Sungai Banyuasin Sumatera Selatan. Makalah disampaikan pada Seminar dan Rapat Tahunan (SEMIRATA) BKS Barat Bidang MIPA. Mei 9-10. Unlam. Banjarmasin.
- Isnaini. 2014. Komposisi dan Kelimpahan Fitoplankton di Perairan Sekitar Pulau Maspari, Ogan Komering Ilir. *Maspari J.* 6(1):39-45.
- Isnaini & Surbakti, H. 2015. Tingkat pencemaran perairan Sungai Musi ditinjau dari aspek saprobitas dan kualitas air. Laporan akhir penelitian unggulan kompetitif. Universitas Sriwijaya. Indralaya.
- Keputusan Menteri Negara Lingkungan Hidup Nomor 51 Tahun 2004. Tentang Baku Mutu Air Laut. Jakarta.
- Lagus, A., Suomela, J., Weithoff, G., Heikkila, K., Helminen, H. & Sipura, J. 2004. Species-specific differences in phytoplankton responses to N and P enrichments and the N:P ratio in the Archipelago Sea, northern Baltic Sea. *J. Plankt. Res.* 26:779-798. doi: 10.1093/plankt/fbh 070.
- Lindahl, O., & Dahl, E. 1990. On the development of the *Chrysochromulina polylepis* bloom in the Skagerrak in May-June 1988. In: Graneli, E., Sundstrom, B., Edler, L & Anderson, D. M. (eds). *Toxic Mar. Phytoplankton*. Elsevier, New York. pp. 189-194.
- Lukman, M., Nasir, A., Amri, K., Tambaru, R., Hatta, M., Nurfadilah & Noer, R.J. 2014. Silikat terlarut di perairan pesisir Sulawesi Selatan. *J. ITKT.* 6(2): 461-478.
- Mulyani, Widiarti, R. & Wardhana, W. 2012. Sebaran Spesies Penyebab *Harmful Algal Bloom* (HAB) di Lokasi Budidaya Kerang Hijau (*Perna viridis*) Kamal Muara Jakarta Utara pada Bulan Mei 2011. *J. Akuatika.* III(1):28-39.
- Newell, G.E. & Newell, R.C. 1977. *Marine Plankton; A Practical Guide*. Hutchinson & Co Ltd. London. 207 p.
- Nontji, A. 2005. *Laut Nusantara*. Penerbit Djambatan. Jakarta. 372 p.
- Nybakken, J.W. 1992. *Biologi Laut : Suatu Pendekatan Ekologis*. Terjemahan. Eidman, H. M., Koesoebiono, Bengen, D. G., Hutomo, M dan Subarjo, S. PT. Gramedia Pustaka Utama. Jakarta. 495 p.
- Odum, E.P. 1998. *Dasar-Dasar Ekologi*. Terjemahan. Samingan, T. Edisi Ketiga. Gajah Mada University Press. Yogyakarta. 697 p.
- Pednekar, S.M., Matondkar, S.G.P. & Kerkar, V. 2012. Spatiotemporal Distribution of Harmful Algal Flora in the Tropical Estuarine Complex of Goa, India. *The Scientific World J.* 2012:11 p. doi: 10.1100/2012/596276
- Pirzan, A.M. & Pong-Masak, P.R. 2008. Hubungan keragaman fitoplankton dengan kualitas air di Pulau Bauluang, Kabupaten Takalar, Sulawesi Selatan. *Biodiversitas.* 9(3):217-221
- Prianto, T., Ulqodry, Z. & Aryawati, R.. 2013. Pola sebaran konsentrasi klorofil-a di Selat Bangka dengan menggunakan citra aqua-modis. *Maspari J.* 5(1): 22-33.
- Praseno, D.P & Sugestiningih. 2000. *Retaid di Perairan Indonesia*. Pusat Penelitian dan Pengembangan Oseanologi, LIPI. Jakarta. Hal 2-34.
- Raymont, J.E.G. 1963. *Plankton and Productivity in The Ocean*. A Pergamon Press Book. Mc. Millan, Co. New York. 489 p.
- Richardson, A.J. 2008. In hot water: Zooplankton and Climate Change. *ICES J. Mar Sci.* 65:279-295. doi: 10.1093/icesjms/fsn028.
- Risamasu, F.J.L & Prayitno, H.B. 2011. Kajian Zat Hara Fosfat, Nitrit, Nitrat dan Silikat di Perairan Matasiri, Kalimantan Selatan. *Ilmu Kelautan.* 16(3) : 135-142
- Romimohtarto, K. & Juwana, S.. 2004. *Meroplankton Laut; Larva Hewan Laut yang Menjadi Plankton*. Djambatan. Jakarta.
- Rokhim, K., Arisandi, A. & Abida, I.W. 2009. Analisis kelimpahan fitoplankton dan ketersediaan nutrisi (NO₃ dan PO₄) di perairan Kecamatan Kwanyar Kabupaten Bangkalan. *Jurnal Kelautan.* 2(2):7-16.
- Sembiring, M.R., Melki & Agustriani, F. 2012. Kualitas perairan muara Sungsang ditinjau dari konsentrasi bahan organik pada kondisi pasang surut. *Maspari J.* 2: 238-247.
- Simanjuntak. 2007. Kadar fosfat, nitrat dan silikat di Teluk Jakarta. *J. Fish Sci.* IX(2):274-287.
- Soedibjo, B.S. 2006. Struktur komunitas fitoplankton dan hubungannya dengan beberapa parameter lingkungan di perairan

- Teluk Jakarta. *Oseanologi dan Limnologi di Indonesia*. 40:65-78.
- Soedibjo, B.S. 2007. Fenomena kehadiran *Skeletonema* sp di Teluk Jakarta. *Ilmu Kelautan*. 12(3):119-124.
- Sarinda, F. & Dewiyanti, I. 2013. Keragaman fitoplankton di perairan estuaria Kuala Gigieng Kabupaten Aceh Besar, Provinsi Aceh. *Depik*. 2(1): 20-25.
- Surbakti, H., Aryawati, R. & Isnaini. 2011. Pemodelan Sebaran Nutrien dan Produktivitas Primer untuk Identifikasi Daerah Penangkapan Ikan di Perairan Selat Bangka. Universitas Sriwijaya. Indralaya.
- Tambaru, R. 2008. Dinamika Komunitas Fitoplankton dalam Kaitannya dengan Produktivitas Perairan di Perairan Pesisir Maros Sulawesi Selatan. Disertasi. Pascasarjana IPB, Bogor.
- Thoha, H. 2013. Pengaruh musim terhadap plankton di perairan Riau Kepulauan dan Sekitarnya. *Makara Sains*. 7(2):59-70.
- Thoha, H & R. Aryawati. 2014. Kondisi Fitoplankton di Perairan Teluk Jakarta. Seminar Nasional MIPA: Peran MIPA dalam pengelolaan sumberdaya alam untuk kemakmuran bangsa. Hal 516-526.
- Tomas, C.R. 1997. Identifying Marine Fitoplankton. Academic Press. California. USA
- Wardhana, A.W. 1995. Dampak Pencemaran Lingkungan. Penerbit Andi Offset. Yogyakarta.
- Widiarti, R. 2000. Pola Suksesi Organisme Penyebab Red Tide *Pyrodinium bahamense* Plate di Teluk Hurun Lampung Selatan [Tesis]. IPB. Bogor. 80 hal.
- Yamaji. 1984. Illustration of the Marine Plankton of Japan. Hoikusho, Osaka, Japan. 369 p.

ORIGINALITY REPORT

12%

SIMILARITY INDEX

9%

INTERNET SOURCES

10%

PUBLICATIONS

7%

STUDENT PAPERS

PRIMARY SOURCES

1	www.semanticscholar.org Internet Source	3%
2	ojs.pps.unsri.ac.id Internet Source	2%
3	Submitted to Universitas Diponegoro Student Paper	1%
4	knepublishing.com Internet Source	1%
5	uad.portalgaruda.org Internet Source	1%
6	Submitted to Alexandru Ioan Cuza University of Iasi Student Paper	1%
7	Rima Nabilah Haifa, Zahidah Hasan, Heti Herawati, Isni Nurruhwati, Asep Sahidin. "Spatial Distribution of Phytoplankton in Jatigede Reservoir, West Java, Indonesia", Asian Journal of Fisheries and Aquatic Research, 2020 Publication	1%

8	Submitted to Universitas Indonesia Student Paper	1 %
9	repository.ub.ac.id Internet Source	1 %
10	Mayang Yeanny, Ternala Barus, Herman Mawengkang, Miswar Mulya. "Model of Phytoplankton Diversity in Belawan River, North Sumatera, Indonesia", Pakistan Journal of Biological Sciences, 2023 Publication	1 %
11	ppsub.ub.ac.id Internet Source	1 %
12	Rudy Sugiharto, Agus Tjahjono. "The Relationship between Plankton Abundance and Abiotic Parameters in the Downstream Section of the Musi River, Palembang", Egyptian Journal of Aquatic Biology and Fisheries, 2021 Publication	1 %

Exclude quotes On

Exclude bibliography On

Exclude matches < 1%