

[IJC] Editor Decision

From: Tri Joko Raharjo (trijr_mipa@ugm.ac.id)

To: etihayati74@yahoo.com

Date: Thursday, December 6, 2018 at 11:43 AM GMT+7

Ferlinahayati Ferlinahayati:

We have reached a decision regarding your submission to Indonesian Journal of Chemistry, " α -Glucosidase Inhibitory and A Leptospermone Derivative from *Rhodomyrtus tomentosa*".

Our decision is: Revisions Required

Comments of the reviewers can be seen in the bottom part of this email. The revised paper has to be completed with responds for the reviewer's comments, point by point, in the beginning pages of the paper, and the revised parts should be indicated with different color of letters or author's comments. The revised paper has to be resubmitted in the system within three weeks.

Thank you for your intending to contribute the journal and giving us to read your work.

Best regards,

Tri Joko Raharjo
Laboratory of Organic Chemistry, Department of Chemistry, Universitas Gadjah Mada
trijr_mipa@ugm.ac.id

Reviewer A:

Additional Comment::

1. This compound is not new, it should be written reference of the comparison compounds.
 2. The value of the coupling constant (J) on the H-NMR to be explained
 3. It would be better, when using proof reader
-

Reviewer B:

- The significance and objective of this study have been explained clearly in the introduction. However, there is no originality stated in introduction. It would be better if the authors could confirm the originality of this study on introduction part.
 - and at discussion section, discussion of inhibitory activity must be added more, compare with other papers.
 - Others, please check the manuscript.
-

Reviewer C:

Additional Comment::

Indonesian Journal of Chemistry
<https://jurnal.ugm.ac.id/ijc>
Indexed by SCOPUS since 2012

Re: [IJC] Editor Decision

From: Tri Joko Raharjo (trijr_mipa@ugm.ac.id)

To: etihayati74@yahoo.com

Date: Monday, January 7, 2019, 9:32 AM GMT+7

Please your answer to reviewer's comments as well as submit your revised paper through Indo J Chem OJS system as you did for paper sbmission

On Wed, Dec 26, 2018 at 6:57 AM ferlina hayati <etihayati74@yahoo.com> wrote:

Dear Editor,

Thank you to for reviewing my paper.

I have been resubmitted the revised paper to the system as well as my responds for the reviewer's comment in the separated paper

Here in I also attach the revised paper and my reponds for the reviewer's.

Best regards,

Ferlinahayati

Chemistry department, FMIPA, Sriwijaya University

On Thursday, December 6, 2018, 11:43:08 AM GMT+7, Tri Joko Raharjo <trijr_mipa@ugm.ac.id> wrote:

Ferlinahayati Ferlinahayati:

We have reached a decision regarding your submission to Indonesian Journal of Chemistry, " α -Glucosidase Inhibitory and A Leptospermone Derivative from *Rhodomyrtus tomentosa*".

Our decision is: Revisions Required

Comments of the reviewers can be seen in the bottom part of this email. The revised paper has to be completed with responds for the reviewer's comments, point by point, in the beginning pages of the paper, and the revised parts should be indicated with different color of letters or author's comments. The revised paper has to be resubmitted in the system within three weeks.

Thank you for your intending to contribute the journal and giving us to read your work.

Best regards,

Tri Joko Raharjo

Laboratory of Organic Chemistry, Department of Chemistry, Universitas Gadjah Mada

trijr_mipa@ugm.ac.id

Reviewer A:

Additional Comment::

1. This compound is not new, it should be written reference of the comparison compounds.
2. The value of the coupling constant (J) on the H-NMR to be explained
3. It would be better, when using proof reader

Reviewer B:

- The significance and objective of this study have been explained clearly in the introduction. However, there is no originality stated in introduction. It would be better if the authors could confirm the originality of this study on introduction part.
- and at discussion section, discussion of inhibitory activity must be added more, compare with other papers.
- Others, please check the manuscript.

Reviewer C:

Additional Comment:

Indonesian Journal of Chemistry

<https://jurnal.ugm.ac.id/ijc>

Indexed by SCOPUS since 2012

α -Glucosidase Inhibitory and A Leptospermone Derivative from *Rhodomyrtus tomentosa*

Ferlinahayati^{1*}, Daniel Alfarado¹, Eliza¹, and Budi Untari²

¹Department of Chemistry, Faculty of Mathematics and Natural Sciences, University of Sriwijaya
Jalan Raya Palembang Prabumulih Km 32, Ogan Ilir, South Sumatera, Indonesia 30622

²Department of Pharmacy, Faculty of Mathematics and Natural Sciences, University of Sriwijaya
Jalan Raya Palembang Prabumulih Km 32, Ogan Ilir, South Sumatera, Indonesia 30622

* Corresponding author, tel/: 081394741890, email: etihayati74@yahoo.com

ABSTRACT

One of the treatments for diabetes mellitus disease is to control blood sugar level using an inhibitor of α -glucosidase enzyme. The methanol extracts of the fruit, stem, and leaves of *Rhodomyrtus tomentosa* were found significant in inhibiting α -glucosidase (IC₅₀ 20.57, 20.36 and 43.99 μ g/mL respectively). The ethyl acetate and *n*-butanol fractions from the methanol extract of *R. tomentosa* fruit exhibited the potent inhibition (IC₅₀ 13.49 and 19.29 μ g/mL) compare to acarbose and *n*-hexane fraction (IC₅₀ 383.68 and 1175.16 μ g/mL). A leptospermone derivative, rhodomyrtosone D was isolated from the ethyl acetate fraction of *R. tomentosa* fruit. The structure of rhodomyrtosone D was identified base on spectroscopic analysis, as well as comparing with literature data. The α -glucosidase inhibition of rhodomyrtosone D (IC₅₀ 110.45 μ g/mL) was 3.5 fold more potent than acarbose. Thus, *R. tomentosa* plant could be potential as a natural resource of α -glucosidase inhibitor.

Keywords: α -glucosidase, *Rhodomyrtus tomentosa*, antidiabetic, rhodomyrtosone D, ethyl acetate fraction

ABSTRAK

Salah satu penanganan diabetes mellitus adalah dengan mengontrol kadar gula darah menggunakan penghambat kerja enzim α -glukosidase. Ekstrak metanol buah, batang dan daun *R. tomentosa* menunjukkan penghambatan α -glukosidase yang signifikan (IC₅₀ 20,57; 20,36 dan 43,99 μ g/mL). Fraksi etil asetat dan *n*-butanol yang diperoleh dari ekstrak metanol buah *R. tomentosa* menunjukkan penghambatan yang potensial (IC₅₀ 13,49 dan 19,29 μ g/mL) dibandingkan dengan akarbose dan fraksi *n*-hexana (IC₅₀ 383,68 and 1175,16 μ g/mL). Suatu

turunan leptospermon yaitu rhodomyrtosone D telah diisolasi dari fraksi etil asetat buah *R. tomentosa*. Struktur senyawa rhodomyrtosone D ditetapkan berdasarkan analisis spektroskopi dan membandingkan dengan literatur. Penghambatan α -glukosidase dari rhodomyrtosone D menunjukkan 3,5 kali lebih kuat dibandingkan dengan akar bosa. Dengan demikian, tumbuhan *R. tomentosa* berpotensi sebagai sumber alami penghambat enzim α -glukosidase.

Kata kunci: α -glukosidase, *Rhodomyrtus tomentosa*, antidiabetes, rhodomyrtosone D, fraksi etil asetat.

INTRODUCTION

Diabetes mellitus (DM) is a group of metabolic disorder, in which there are high blood sugar levels (hyperglycemia) over a prolonged period [1]. It will happen if the pancreas does not produce enough insulin that is able to convert sugar into energy, or the body's cells do not respond well to the insulin produced. Some serious complication of hyperglycemia such as cardiovascular disease, damage to the eyes, atherosclerosis, and chronic kidney disease (nephropathy) can also occur [2]. The control of blood sugar level by inhibition of carbohydrate-hydrolyzing enzymes in the digestive organ is believed to be important in hyperglycemia treatment [1]. The α -glucosidase, an enzyme in the small intestine is responsible for the degradation of carbohydrate. The α -glucosidase inhibitor will interfere with the digestion of carbohydrate and thereby reduce the postprandial glucose level and insulin responses in a diabetic patient [2-3]. Acarbose, miglitol, and voglibose have been found as an α -glucosidase inhibitor and currently clinically used to control blood glucose of diabetic patients [4]. However, they have been caused serious gastrointestinal side effects. Nowadays, natural resources have received tremendous attention as a therapeutic agent in the inhibition of α -glucosidase and have shown very promising biological activity.

Karamunting is locally named (Sumatera island) for *Rhodomyrtus tomentosa* and belonging to Myrtaceae family. This plant is an evergreen shrub which is native to Southern Asia and Southeast Asia and is widely distributed in Indonesia. *R. tomentosa* is widely used as traditional medicines to treat a variety of disease caused by bacteria such as diarrhea, dysentery and urinary tract infections [5-6]. In addition, its ripe fruits are used to boost the immune system [7]. Biologically, ethanolic extract of *R. tomentosa* fruits possesses potent antioxidant activities on DPPH radical scavenging activity, reducing power as well as inhibition of lipid peroxidation activity [8]. Furthermore, some extract of this plants were reported as antibacterial and anti-hepatitis properties [9]. Chemically, various secondary metabolites have been reported such as polyketide, flavonoids, anthocyanins, stilbenoids, and triterpenoids [7-11]. Rhodomyrtone, a phloroglucinol polyketide from *R. tomentosa* have displayed significant antibacterial activities against Gram-positive bacteria and suggested as a new candidate as a natural antibacterial drug [10]. Meanwhile, tomentosone A, a hexacyclic phloroglucinol was reported as antimalarial against

chloroquine-resistant and sensitive strains of *Plasmodium falciparum*. Resveratrol and piceatannol, a stilbenoid compound has been characterized from this plant [7]. A stilbenoid compound from *Syagrus romanzoffiana* was reported as a potential hypoglycemic agent. However, there is no literature on the α -glucosidase inhibitory of *R. tomentosa* and its bioactive chemical compound. In a search for potential α -glucosidase inhibitor from natural resources, we have been investigated the ability of *R. tomentosa* plant to inhibit the activity of the α -glucosidase enzyme as well as to isolate the bioactive compound. One active compound, rhodomyrtosone D (1) was isolated and its α -glucosidase inhibition was determined. The following describes the outcomes of these efforts.

EXPERIMENTAL SECTION

Materials

Rhodomyrtus tomentosa (fruits, leaves, and stem) were collected from Inderalaya, Ogan Ilir, South Sumatera. The plant was identified at Herbarium Anda, Department of Biology, University of Andalas. The solvents (methanol, *n*-hexane and ethyl acetate) were a technical quality that is distilled while *n*-butanol and dimethylsulfoxide (DMSO) were pro analysis (p.a) from Merck. The α -glucosidase (from *Saccharomyces cerevisiae*) and *p*-nitro-phenyl- α -D-glucopyranoside were purchased from Sigma-Aldrich. Bovine serum albumin (BSA) was purchased from Merck. Silica gel 60G (Merck) was used for vacuum liquid chromatography and silica gel 60 PF₂₅₄ (Merck) was used for radial chromatography. TLC analysis was performed on Kieselgel 60 GF₂₅₄, 0.25 mm aluminum plate (Merck) and visualized with cerium sulfate.

Instrumentation

Incubator Biosan PST-60HL was used for sample incubation process. The absorbance of *p*-nitrophenol was measured by a Tecan Infinite F50 Microplate reader. UV spectrum was recorded with Shimadzu UV-1240 spectrophotometer. IR spectrum was determined using KBr pellets on a Perkin Elmer FTIR Spectrum One spectrophotometer. ¹H-NMR (500 MHz) and ¹³C-NMR (125 MHz) spectra were recorded with Agilent DD2 spectrometer, using residual and deuterated solvent peaks as reference standards.

Procedure

Extraction of sample for assay

As much as 100 gr of each the dried powdered sample (fruits, leaves, and stem) of *R.tomentosa* were extracted by maceration method using methanol (400 mL) as the solvent at the

room temperature. The maceration process was carried out three times (@ 24 hours). The methanol solvents were evaporated under reduced pressure to give crude extracts of methanol of fruit, leaves, and stem (4.6, 4.2 and 3.9 g respectively). The crude methanol extract of fruit was partitioned successively with *n*-hexane, ethyl acetate, and *n*-butanol and produce of each fraction after the solvent was evaporated.

In-vitro α -glucosidase inhibition assay

The α -glucosidase assay has been performed using the spectrophotometric method as previously described [2, 12, 13] with slight modification. As much as 10 μ L of the sample at various concentrations was added with 55 μ L of 50 mM phosphate buffer (pH 6.8) and 10 μ L of 10 mM *p*-nitrophenyl- α -D-glucopyranoside as the substrate. After preincubated for 5 min at 37 °C, 25 μ L of 0.1 U/mL α -glucosidase (in the phosphate buffer pH 6.8 containing 0.1 mg/mL bovine serum albumin) was added. The mixture was then incubated for 30 min at 37 °C. After that, the stopped solution (100 μ L of 200 mM Na₂CO₃) was added to the mixture. The absorbance of the *p*-nitrophenol released due to hydrolysis of the substrate by the α -glucosidase was measured by microplate reader at 405 nm. The blank solution was prepared by replaced sample solution by DMSO. Acarbose (Glucobay®) is used as a positive control. The percentage inhibition of α -glucosidase was calculated using the following equation: Inhibition % = $[1 - (A_{\text{sample}} / A_{\text{blank}})] \times 100$. The IC₅₀ was calculated by linear regression equation analysis between concentration and percentage inhibition.

Extraction and Isolation of *R. tomentosa* fruits

The dried fruits (2 kg) was extracted with methanol (3 x 7 L, 24 h each) by maceration method. The methanol extract was concentrated under reduced pressure to give 1.1 L syrup which was suspended in distilled water. This suspension was partitioned successively with *n*-hexane, ethyl acetate, and *n*-butanol to afford *n*-hexane, ethyl acetate and *n*-butanol fractions. The ethyl acetate fraction (15 g) was fractionated by vacuum liquid chromatography on silica gel 60 G, eluting with *n*-hexane-ethyl acetate system with increment ethyl acetate gradually (9:1, 8:2, 7:3, 6:4, 4:6, 2:8, 1:9, and 0:10, each 150 mL) to give 8 fractions (A-H). Fraction C (374 mg) was further separated using radial chromatography over silica gel 60 PF₂₅₄ (1 mm), eluted with *n*-hexane-ethyl acetate gradually (85:15, 80:20, 75:25, 70:30, 60:40, 50:50) to yield a leptospermone derivative 1 (8.9 mg)

RESULTS AND DISCUSSION

The α -Glucosidase inhibition of extracts and fractions

The extraction of three parts of *R. tomentosa* (fruit, stem, and leaves) produced methanol extract 4.6, 3.9, and 4.2 g respectively. All of these extracts were tested for α -glucosidase inhibitory using *p*-nitrophenyl- α -D-glucopyranoside as the substrate and acarbose as the reference or positive control. The methanol extract from the stem and fruit have a similar ability to inhibit α -glucosidase activity (IC₅₀ 20.36 and 20.57 μ g/mL). Both of these extracts demonstrated two times more potent than the leaves methanol extract (IC₅₀ 43.99 μ g/mL) (Figure 1). All three methanol extracts possessed high potency in inhibiting α -glucosidase compare to the reference drug, acarbose (IC₅₀ 383.68 μ g/mL) (Table 1). Previously, it has been reported that *R. tomentosa* fruit contains stilbenoid compound, such as resveratrol, and piceatannol [7]. These stilbenoids showed the more potent inhibition of α -glucosidase activity with IC₅₀ 91 and 60 μ g/mL respectively than acarbose (IC₅₀ 247 μ g/mL) [4]. In addition, another phenolic compounds such as flavonoid isolated from *Morus alba* and anthocyanins isolated from noble muscadine grapes have been reported as potential α -glucosidase inhibitory [11,14]. Meanwhile, the triterpenoid saponins from *Gypsophila oldhamiana* and highly oxygenated triterpenoid from *Fagara tessmannii* and *Luculia pinceana* also showed significant α -glucosidase inhibitory comparing to acarbose [1,15].

Base on its inhibition of α -glucosidase, the methanol extract of fruits was partitioned into *n*-hexane, ethyl acetate, and *n*-butanol. Ethyl acetate fraction had the highest α -glucosidase inhibitory (IC₅₀ 13.49 μ g/mL than *n*-butanol fraction (IC₅₀ 19.29 μ g/mL) due to its phenolic content, meanwhile, the *n*-hexane fraction was not as potent as α -glucosidase inhibitory (IC₅₀ 1175.16 μ g/mL) (Table 1 and Figure 1). Compounds typically found in hexane fractions are non-polar triterpenoid and steroid, while triterpenoid saponins and highly oxygenated triterpenoid found in the polar fraction such as ethyl acetate and *n*-butanol [1, 15].

Table 1. Inhibitory effect of the extract, fraction and compound on α -glucosidase activity

Extract/compound	Inhibitor concentration (IC ₅₀ , μ g/mL)
MeOH extract of the leaves	43.99
MeOH extract of the stem	20.36
MeOH extract of the fruit	20.57
<i>n</i> -hexane fraction of the fruit	1175.16
Ethyl acetate fraction of the fruit	13.49
<i>n</i> -butanol fraction of the fruit	19.29
Compound 1	110.45
Acarbose*	383.68

*positive control




Figure 1. Effect of extracts and fractions on the inhibition of α -glucosidase

Isolation and structural elucidation

The sequential partition to the methanol crude extract of *R. tomentosa* fruits (87 g) yielded *n*-hexane, ethyl acetate and *n*-butanol fraction (1.54, 17.81 and 0.44 g respectively). Ethyl acetate fraction with the highest α -glucosidase inhibition was chromatographed over silica gel with some chromatographic technique to afford compound **1**.

Compound **1** was isolated as a white powder with m.p. 120-121 °C. The UV spectrum in methanol showed the maximum absorption at 242 nm which indicated the presence of α,β carbonyl unsaturated. The IR spectrum displayed absorption for the isolated carbonyl group at 1715 cm^{-1} as well as conjugated carbonyl group at 1678 and 1663 cm^{-1} which consisted to UV spectrum. In addition, there is absorption for C-H aliphatic group at 2976 and 2941 cm^{-1} . The ^{13}C -NMR (125 MHz, CDCl_3) was showed the presence of 14 signal. Two of the signal confirmed the existence of the isolated and conjugated carbonyl at δ_{C} 212.2 ppm and δ_{C} 192.2 ppm respectively. In addition, ^{13}C -NMR displayed the presence of five other quaternary carbon signal (δ_{C} 175.5 (oxy-carbon), 128.3, 113.2, 56.6, and 45.3 ppm), two signal for methine carbon (δ_{C} 46.6 and 34.5 ppm), and five signal for methyl carbon (δ_{C} 25.9, 24.5, 24.0, 22.4 and 15.6 ppm). Considering the intensity of quaternary carbon signal at 128.3 ppm with the six other quaternary carbon (included the carbonyl) which has a ratio of 1:2, indicating that the six quaternary carbon is equivalent to twelve carbon. Furthermore, the five methyl carbon signals have an intensity ratio of 2: 1 with a carbon methine signal at δ_{C} 34.5 ppm, consequently each of these methyl signals is identical for 2 methyl carbon (there are a total of 10 methyls). Based on this, compound **1** actually has 25 carbon atoms. The ^1H -NMR (500MHz, CDCl_3) spectrum exhibited the presence of a singlet signal of methine proton at δ_{H} 4.67 ppm. The spectrum also indicated the presence of an isopropyl unit with the appearance of a doublet signal at δ_{H} 1.00 ppm (6H, *d*, J = 6.9 Hz, 2x CH_3) which is adjacent to

the methine proton at δ_{H} 2.35 ppm (1H, *sept*, $J = 6.9$ Hz). These constant coupling value indicates that the both signals are correlated to each other as vicinal aliphatic protons. In addition, there are three singlet signals at δ_{H} 1.41 (12H), 1.32 (6H) and 1.25 ppm (6H) which indicate the presence of 8 methyl groups. The HMBC correlation revealed a correlation of both methyl on a geminal dimethyl group (δ_{H} 1.25 and 1.32 ppm) to the isolated and conjugated carbonyl group (δ_{C} 212.2 and 192.2 ppm) as well as correlation of both methyl on another geminal dimethyl group to the isolated carbonyl (δ_{C} 212.2 ppm) and oxy-carbon (δ_{C} 175.5 ppm). These explained that both of geminal dimethyl are α position in β -triketone unit. Based on the previous NMR data, there is actually two symmetrical unit of β -triketone. Furthermore, the correlation between of proton δ_{H} 4.67 ppm to isopropyl unit (δ_{C} 34.5 ppm) and oxy-carbon (δ_{C} 175.5 and 128.3 ppm) indicating that the isopropyl group was an adjacent bis-furan ring and the bis-furan ring was integrated with the β -triketone unit. According to these spectroscopic evidence and comparing to the those of reported literature [11], the structure of compound **1** was established as rhodomirtosone D. This compound has been previously reported from *R. tomentosa* leaves [11].




Figure 2. Structure of compound **1** (rhodomirtosone D)

Table 2. NMR data of compound **1** in CDCl_3 and rhodomirtosone D

No	Compound 1			Rhodomirtosone D [11]	
	δ_{C}	δ_{H} (ΣH , <i>mult</i> , J_{Hz}) ^b	HMBC (H \rightarrow C)	δ_{C}	δ_{H} (<i>mult</i> , J_{Hz})
1 (1')	192.2	-	-	192.4	-
2(2')	56.6	-	-	56.4	-
3(3')	212.2	-	-	212.1	-
4(4')	45.3	-	-	45.2	-
5(5')	175.5	-	-	175.7	-
6(6')	113.2	-	-	113.0	-
7(7')	25.8	1.25 (6H, s)	C-3(3'), C-1 (1'), C-2(2'), C- 8(8')	25.7	1.27 (s)
8(8')	22.4	1.32 (6H, s)	C-3(3'), C- 1 (1'), C- 2(2'), C- 7(7')	22.3	1.34 (s)
9(9')	24.0	1.41 (6H, s)	C-3(3'), C-5(5'), C-4(4'), C- 9(9'), C-10(10')	23.9	1.44 (s)
10(10')	24.5	1.41 (6H, s)	C-3(3'), C-5(5'), C-4(4'), C- 9(9'), C-10(10')	24.4	1.44 (s)

1''	46.6	4.67 (1H, s)	C-5(5'), C- 2'', C- 6(6'), C- 3''	46.5	4.69 (s)
2''	128.3	-	-	128.2	-
3''	34.5	2.35 (1H, sept, 6.9)	C-2'', C- 1'', C- 4'', C- 5''	34.4	2.37 (sept, 6.9)
4'', 5''	15.6	1.00 (6H, d, 6.9)	C-2'', C- 3'', C- 4', C- 5''	15.5	1.02 (d, 6.9)

The isolated compound **1** (rhodomyrtosone D) was examined for α -glucosidase inhibitory activity with concentration range about 30.77 to 0.24 $\mu\text{g/mL}$. The α -glucosidase inhibitory effect of rhodomyrtosone D (17.7% at 30.77 $\mu\text{g/mL}$) seems higher than the acarbose (8.54 % at 30.77 $\mu\text{g/mL}$). Using the extrapolation method to linear regression, the IC_{50} of rhodomyrtosone D on inhibiting α -glucosidase was 110.45 $\mu\text{g/mL}$.

CONCLUSION

The leaves, the stem, and the fruit of *R. tomentosa* plant were potential as a source of a natural antidiabetic, especially from the ethyl acetate fraction of the fruit. A bioactive compound, rhodomyrtosone D was isolated from the fruit of *Rhodomyrtus tomentosa* and showed higher α -glucosidase inhibition than acarbose.

ACKNOWLEDGMENTS

The author would like thanks to Dr. Nurainas, M.Si from Herbarium Anda, Andalas University for identification of plant specimen. We are grateful to the Ministry of Research, Technology and Higher Education for research grants through PD-UPT scheme (No: 093/SP2H/LT/DPRM/IV/2018).

REFERENCES

- Yin, Z., Zhang, W., Feng, F., Zhang, Y., and Kang, W., 2014, α -Glucosidase Inhibitors Isolated from Medicinal Plants, *Food Science, and Human Wellness*, 3, 136-174.
- Choudhary, M.I., Adhikari, A., Rasheed, S., Marasini, B.P., Hussain, N., Kaleem, W.A., and Rahman, A., 2011, Cyclopeptide Alkaloid of *Ziziphus oxyphylla* Edw as Novel Inhibitors of α -Glucosidase Enzyme and Protein Glycation, *Phytochemistry Letters*, 4, 404-406.
- Lam, S.H., Cheng, J.M., Kang, C.J., Chen, C.H., and Lee, S.S., 2008, α -Glucosidase Inhibitors from the Seed of *Syagrus romanzoffiana*, *Phytochemistry*, 1173-1178.
- Zhang, A.J., Rimando, A.M., Mizuno, C.S., and Mathews, S.T., 2017, α -Glucosidase Inhibitory Effect of Resveratrol and Piceatannol, *The Journal of Nutritional Biochemistry*, 47, 86-93.
- Lavanya, G., Voravuthikunchai, S.P, and Towatana, N.H., 2012, Acetone Extract from *Rhodomyrtus tomentosa*: A Potent Natural Antioxidant, *Evidence-Based Complementary and Alternative Medicine*, 2012, Article ID 535479, 1-8.

6. Limsuwan, S., Kayser, O., and Voravuthikunchai, S.P., 2012, Antibacterial Activity of *Rhodomyrtus tomentosa* (Aiton) Hassk. Leaf Extract against Clinical Isolates of *Streptococcus pyogenes*, *Evidence-Based Complementary and Alternative Medicine*, 2012, Article ID 697183, 1-6.
7. Lai, T.N.H., Herent, M.F., Quetin-Leclercq, J., Nguyen, T.B.T., Larondelle, Y., Andre, C.M., and Rogez, H., 2013, Piceatannol, a Potent Bioactive Stilbene, as Major Phenolic Component in *Rhodomyrtus tomentosa*, *Food Chemistry*, 138: 1421-1430
8. Wu, P., Ma, G., Li, N., Deng, Q., Yin, Y., and Huang, R., 2015, Investigation of In Vitro and in Vivo Antioxidant Activities of Flavonoids Rich Extract from the Berries of *Rhodomyrtus tomentosa* (Ait.) Hassk., *Food Chemistry*, 173, 194-202.
9. Hiranrat, A., Mahabusakaram, W., Carrol, A.R., Duffy, S., and Avery, V.M., 2012, Tomentosones A and B, Hexacyclic Phloroglucinol Derivatives from the Thai Shrub *Rhodomyrtus tomentosa*, *J. Org. Chem.*, 77, 680-683.
10. Limsuwan, S., Trip, E.N., Kouwen, T.R.H.M., Piersma, S., Hiranrat, A., Mahabusakaram, W., Voravuthikunchai, S.P., Dijl, J.M., and Kayse, O., 2009, Rhodomyrtone, A new Candidate as Natural Antibacterial Drug from *Rhodomyrtus tomentosa*, *Phytomedicine*, 16, 645-651.
11. Yang, Z., Wang, Y., Wang, Y., and Zhang, Y., 2012, Bioassay-Guided Screening and Isolation of α -glucosidase and Tyrosinase Inhibitors from Leaves of *Morus alba*, *Food Chemistry*, 617-625.
12. Anisah, L.N., Syafii, W., Pari, G., and Sari, R.K., 2018, Antidiabetic Activities and Identification of Chemical Compound from Samama (*Anthocephalus macrophyllus* (Roxb) Havil), *Indones. J. Chem*, 18(1), 66-74.
13. Hiranrat, A., and Mahabusakaram, W., 2008, New Acylphloroglucinols from the Leaves of *Rhodomyrtus tomentosa*, *Tetrahedron*, 64, 11193-11197.
14. You, Q., Chen, F., Wang, X., Luo, P.G., and jiang, Y., 2011, Inhibitory Effect of Muscadine Anthocyanins on α -Glucosidase and Pancreatic Lipase Activities, *J. Agric. Food. Chem.*, 59, 9506-9511.
15. Luo, J.G., Ma, L., and Kong, L.Y., 2008, New Triterpenoid Saponins with Strong α -Glucosidase Inhibitory Activity from the Roots of *Gypsophila oldhamiana*, *Bioorg. Med. Chem.*, 16, 2912-2920.

Dear reviewer,

Thank you to for reviewing my paper entittled " **α -Glucosidase Inhibitory and A Leptospermone Derivative from *Rhodomyrtus tomentosa***" This is my response to your comments :

Reviewer A

The revised part of **reviewer A** indicated with **red color**

No	Revised/Comment of author
A1	It was deleted
A2	It has been revised to " <i>Rhodomyrtus tomentosa</i> "
A3	It has been revised to "fractions"
A4	It has been added "The structure of rhodomyrtosone D was"
A5	It has been added "rhodomyrtosone D was"
A6	It has been added with "."
A7	It has been added with "rhodomyrtosone D"
A8	It has been added with " menunjukkan"
A9	It Has been revised to " <i>R. tomentosa</i> "
A10	It has been added with "the"
A11	It has been revised to " antimalarial"
A12	It has been added with "The"
A13	It has been revised to " ¹ H-NMR (500 MHz) and ¹³ C-NMR (125 MHz)"
A14	It has been revised to "As much as 100 gr of each"
A15	It has been added with "@ 24 hours"
A16	It has been revised to " extracts"
A17	It has been added with :As much as"
A18	It has been revised
A19	It has been revised
A20	has been revised to "was"
A21	It has been revised
A22	It has been revised and state clearly.
A23	It has been added with "the"
A24	It has been added with " ,"
A25	The sentence has been revised and the reference has been changed to

	numbering
A26	Yes, that is correct
A27	Yes, that is correct
A28 & A29	It has been explained.
A30	The reference has been added
A31	It has been revised
Respond to the additional comments:	
<ul style="list-style-type: none"> • Reference of the comparison compounds have been written. • The value of the coupling constant (J) on the H-NMR have been explained in the text 	

Reviewer B

The revised part of **reviewer B** indicated with **blue color**

No	Revised/Comment of author
W1	The references has been added
W2	It has been added
W3	It has been added the objective of this section and previous
W4	It has been changed
W5	It has been changed to mass units
W6	we didn't have any evidence, only base on the literature [7]. So, the sentence has been deleted
W7	It has been marked
W8	It has been changed to mass units
W9	It has been revised
W10	It has been changed to "spectroscopic evidence"
W11	the literature data has been added
Respond to the additional comments:	
<ul style="list-style-type: none"> • The originality of this study has been added in introduction part. • Discussion on inhibitory activity has been added 	

Reviewer C

The revised part of **reviewer B** indicated with green color

- Butanol has been changed to "*n*-butanol).
- The sign of '→' has been changed to " comma"

Sincerely yours,

Ferlinahayati

Chemistry Department, FMIPA, University of Sriwijaya

Re: IJC article information

From: ferlina hayati (etihayati74@yahoo.com)

To: nuryono_mipa@ugm.ac.id; ijc@ugm.ac.id; ijcugm@yahoo.com

Date: Monday, April 22, 2019, 5:06 PM GMT+7

Yth : Editor IJC

Mohon maaf atas kesalahpahaman dari saya.

Sebelumnya pada tanggal 9 Januari saya telah mengupload kembali revisi artikel sesuai saran reviewer ([40990-113820-4-ED.docx](#)), beserta summary perbaikan dan jawaban atas pertanyaan reviewer pada file terpisah ([40990-113820-3-ED.docx](#)).

Karena isi email pada tanggal 11 Januari tersebut persis sama dengan email yang saya terima sebelumnya, maka saya kira email tersebut terkirim ulang by system saja.

Namun saya akan jawab kembali pertanyaan dari reviewer lebih rinci sebagai berikut :

Response to Reviewer A:

1. The isolated compound is not a new compound. On table 2, we have added the NMR data of the comparison compound.
2. We have added an explanation about the value of coupling constant with this sentence "These constant coupling value indicates that the both signals are correlated to each other as vicinal aliphatic protons".
3. We just checked using Grammarly

Response to Reviewer B:

1. The orinality have been added in the introduction with this sentence "However, there is no literature on the a-glucosidase inhibitory of *R. tomentosa* and its bioactive chemical compound"
2. The discussion of inhibitory activity have been added with this sentence "In addition, another phenolic compounds such as flavonoid isolated from *Morus alba* and anthocyanins isolated from noble muscadine grapes have been reported as potential a-glucosidase inhibitory [11,14]. Meanwhile, the triterpenoid saponins from *Gypsophila oldhamiana* and highly oxigenated triterpenoid from *Fagara tessmannii* and *Luculia pinceana* also showed significant a-glucosidase inhibitory comparing to acarbose [1,15]."

and also with this sentence "

Compounds typically found in hexane fractions are non-polar triterpenoid and steroid, while triterpenoid saponins and highly oxygenated triterpenoid found in the polar fraction such as ethyl acetate and *n*-butanol [1, 15]."

Response to reviewer C:

1. We have revised as reviewer suggestion.

Demikianlah yang dapat saya sampaikan, dan saya sangat berharap artikel tersebut dapat diproses lebih lanjut dan bisa terbit di IJC.

Wassalam

Ferlinahayati
Jurusan Kimia FMIPA UNSRI

On Sunday, April 21, 2019, 9:59:51 PM GMT+7, Nuryono Nuryono <nuryono_mipa@ugm.ac.id> wrote:

Sdr Ferlinahayati

Tanggal 11 Januari editor kami telah mengirim decision sebagai berikut. Namun, samai sekarang tdk ada respon revisi dari author.

Ferlinahayati Ferlinahayati:

We have reached a decision regarding your submission to Indonesian Journal of Chemistry, " α -Glucosidase Inhibitory and A Leptospermone Derivative from *Rhodomyrtus tomentosa*".

Our decision is: Revisions Required

Please answer the

1. This compound is not new, it should be written reference of the comparison compounds.
2. The value of the coupling constant (J) on the H-NMR to be explained
3. It would be better, when using proof reader

1. The significance and objective of this study have been explained clearly in the introduction. However, there is no originality stated in introduction. It would be better if the authors could confirm the originality of this study on introduction part.
2. At discussion section, discussion of inhibitory activity must be added more, compare with other papers.
3. Others, please check the manuscript.

Best regards,

Tri Joko Raharjo

Laboratory of Organic Chemistry, Department of Chemistry, Universitas Gadjah Mada

trijr_mipa@ugm.ac.id

Indonesian Journal of Chemistry

<https://jurnal.ugm.ac.id/ijc>

Indexed by SCOPUS since 2012

Pada tanggal Jum, 19 Apr 2019 pukul 13.13 ferlina hayati <etihayati74@yahoo.com> menulis:

Dear Editor of IJC

Relating to our article ID 40990 with the tittle " α -Glucosidase Inhbitory and A Leptospermone Derivative from *Rhodomyrtus tomentosa*", we need the information about the progress and status of the article.

The revised article has been submitted throught the system on 9th Jan 2019.

We look forward to hear from you

Best Regards,

Ferlinahayati

Department of Chemistry,

FMIPA UNSRI

--

Prof. Dr.rer.nat. Nuryono, MS

Editor in Chief

Indonesian Journal of Chemistry

Accredited by DIKTI; Indexed in Scopus since 2012

Re: [IJC] -Glucosidase Inhibitory and A Leptospermone Derivative from Rhodomyrtus tomentosa

From: ferlina hayati (etihayati74@yahoo.com)

To: trijr_mipa@ugm.ac.id

Date: Saturday, May 4, 2019, 1:41 PM GMT+7

Dear Editor,

Thank you for the information. We really hope to get good news soon.

Best Regards,
Ferlinahayati

On Thursday, May 2, 2019, 9:10:25 PM GMT+7, Tri Joko Raharjo <trijr_mipa@ugm.ac.id> wrote:

Dear Authors

Regarding your submission to IJC #40990, I could inform you that at this moment the manuscript is under review to see if the reviewer's comment from previous round have been addressed properly.

Soon after the reviewer give the feedback I am ready to make decision and I will let you know at the first occasion.

Best regards.

Editor

Indonesian Journal of Chemistry

<https://jurnal.ugm.ac.id/ijc>

Indexed by SCOPUS since 2012

[IJC] Copyediting Completed

From: Aulia Ratri Hapsari (aulia.ratri.h@ugm.ac.id)

To: etihayati74@yahoo.com

Cc: trijr_mipa@ugm.ac.id

Date: Thursday, May 23, 2019, 2:27 PM GMT+7

Dear Ferlinahayati Ferlinahayati,

We have now copyedited your submission "^{EG}_{BT}-Glucosidase Inhibitory and A Leptosperme Derivative from *Rhodomyrtus tomentosa*" for Indonesian Journal of Chemistry. To review the proposed changes and respond to Author Queries, please follow these steps:

1. Log into the journal using URL below with your username and password (use Forgot link if needed).
2. Click on the file at 1. Initial Copyedit File to download and open copyedited version.
3. Review the copyediting, making changes using Track Changes in Word, and answer queries.
4. Save file to desktop and upload it in 2. Author Copyedit.
5. Click the email icon under COMPLETE and send an email to the editor.

This is the last opportunity that you have to make substantial changes. You will be asked at a later stage to proofread the galleys, but at that point, only minor typographical and layout errors can be corrected.

Manuscript URL: <https://jurnal.ugm.ac.id/ijc/author/submissionEditing/40990>

Username: ferlinahayati

If you are unable to undertake this work at this time or have any questions, please contact me. Thank you for your contribution to this journal.

Best regards,
Aulia Ratri Hapsari

Indonesian Journal of Chemistry
<https://jurnal.ugm.ac.id/ijc>
Indexed by SCOPUS since 2012



40990-129453-1-CE.docx
76.3kB

[IJC] Copyediting Review Acknowledgement

From: Dwi Siswanta (dsiswanta@ugm.ac.id)

To: etihayati74@yahoo.com

Date: Wednesday, May 29, 2019, 12:52 PM GMT+7

Dear Ferlinahayati Ferlinahayati:

Thank you for reviewing the copyediting of your manuscript, "[\[REDACTED\]](#)-Glucosidase Inhibitory and A Leptospermone Derivative from *Rhodomyrtus tomentosa*," for Indonesian Journal of Chemistry. We look forward to publishing this work.

Kind Regards,
Dwi Siswanta
Laboratory of Analytical Chemistry,
Department of Chemistry,
Universitas Gadjah Mada
Phone +628157951198
Fax +62545188
dsiswanta@ugm.ac.id

Indonesian Journal of Chemistry

<https://jurnal.ugm.ac.id/ijc>

Indexed by SCOPUS since 2012

Re: [IJC] Proofreading Request (Author)

From: ferlina hayati (etihayati74@yahoo.com)

To: dsiswanta@ugm.ac.id

Date: Friday, June 14, 2019 at 09:10 PM GMT+7

Dear editor,

we have done proofreading and made some improvement as follows:

1. in abstract line 8, we have change *R. tometosa* to *R. tomentosa*
2. In introduction line 14, we have change The ^{En} _{En}-glucosidase inhibitor to the α -glucosidase inhibitor.
3. on table 2 in the 2nd line, we have deleted "b" (**superscribe**)
4. **we have refer fig 2 & table 2 in the text.**

Best regard

Ferlinahayati

On Monday, June 10, 2019, 3:57:12 PM GMT+7, Dwi Siswanta <dsiswanta@ugm.ac.id> wrote:

Dear Ferlinahayati Ferlinahayati,

Your submission " α -Glucosidase Inhibitory and A Leptospermone Derivative from *Rhodomyrtus tomentosa*" to Indonesian Journal of Chemistry now needs to be proofread by following these steps.

1. Click on the Submission URL below.
2. Log into the journal and view PROOFING INSTRUCTIONS
3. Click on VIEW PROOF in Layout and proof the galley in the one or more formats used.
4. Enter corrections (typographical and format) in Proofreading Corrections.
5. Save and email corrections to Layout Editor and Proofreader.
6. Send the COMPLETE email to the editor.

=====
Author is required to complete the proofreading stage in no more than ONE WEEK.
=====

Submission URL: <https://journal.ugm.ac.id/ijc/author/submissionEditing/40990>

Username: ferlinahayati

Best regards,
Dwi Siswanta
Laboratory of Analytical Chemistry,
Department of Chemistry,
Universitas Gadjah Mada
Phone +628157951198
Fax +62545188
dsiswanta@ugm.ac.id

Indonesian Journal of Chemistry
<https://jurnal.ugm.ac.id/ijc>
Indexed by SCOPUS since 2012



40990-131829-1-PB - author.docx

89kB

[IJC] Proofreading Acknowledgement (Author)

From: Dwi Siswanta (dsiswanta@ugm.ac.id)

To: etihayati74@yahoo.com

Date: Saturday, June 15, 2019, 11:48 AM GMT+7

Dear Ferlinahayati Ferlinahayati,

Thank you for proofreading the galleys for your manuscript, " α -Glucosidase Inhibitory and A Leptospermone Derivative from *Rhodomyrtus tomentosa*," in Indonesian Journal of Chemistry. We are looking forward to publishing your work shortly.

If you subscribe to our notification service, you will receive an email of the Table of Contents as soon as it is published. If you have any questions, please contact me.

Best regards,
Dwi Siswanta
Laboratory of Analytical Chemistry,
Department of Chemistry,
Universitas Gadjah Mada
Phone +628157951198
Fax +62545188
dsiswanta@ugm.ac.id

Indonesian Journal of Chemistry
<https://jurnal.ugm.ac.id/ijc>
Indexed by SCOPUS since 2012