



Received: 30-08-2023  
Accepted: 10-10-2023

ISSN: 2583-049X

## Length-Weight Relationship and Condition factor of *Notopterus Notopterus* (Pallas, 1769) from East Pedamaran Floodplain, Ogan Komering Ilir, South Sumatra, Indonesia

Muslim Muslim

Department of Fisheries, Faculty of Agriculture, Aquaculture Study Program, Universitas Sriwijaya, Indralaya, Indonesia

Corresponding Author: Muslim Muslim

### Abstract

The study of length-weight relationships and condition factors is a fundamental component of fisheries management. *Notopterus notopterus* (Pallas, 1769) is a fish species native to Indonesia. They are utilized as a side dish, in the culinary and snack industries, and as ornamental fish. This study aimed to analyze the length-weight relationship and condition factors of *N. notopterus* from the East Pedamaran Floodplain, Ogan Komering Ilir, South Sumatra, Indonesia. A total of 108 *N. notopterus* samples were used in this study. The results showed that the length-weight relationship of *N. notopterus* was negatively allometric, with the regression equation  $W = 0.0948.L^{2.2012}$  ( $R^2 = 0.5332$ ). The condition factor value was 1.02, indicating that the *N. notopterus* population was in good condition. The data from this study are very useful for the sustainable management of *N. notopterus* in the future.

**Keywords:** Fish Resource, Marsh, Knifefish, Native Species, Swamp

### Introduction

The length-weight relationships (LWRs) are important and fundamental components of fisheries management tools (Kumari *et al.*, 2019) <sup>[11]</sup>. It is highly crucial for several biological aspects of the species (Freitas *et al.*, 2017) <sup>[6]</sup>, ecological assessments and monitoring (Orlov & Binohlan, 2019) <sup>[24]</sup>, the life history of fishes (Ferdaushy & Alam, 2015) <sup>[5]</sup>, population stock assessment studies (Augustina *et al.*, 2022) <sup>[2]</sup>, understanding maturity and reproduction (Soni & Ujjania, 2017) <sup>[29]</sup>, growth and body condition (Zuchi *et al.*, 2020) <sup>[32]</sup>, and geographic and seasonal (Lima *et al.*, 2021) <sup>[13]</sup>. The LWRs are an important tool in aquaculture management as they provide information on survival, mortality, growth, and total production (Kaur & Rawal, 2017) <sup>[9]</sup>. The condition factor provides information on the physiological condition of the fish in terms of its well-being (Ighwela *et al.*, 2011) <sup>[8]</sup>, overweight status (Ouahb *et al.*, 2021) <sup>[25]</sup>, gonadal development, and maturity status (Pathak *et al.*, 2022) <sup>[26]</sup>. It is an index reflecting interactions between biotic and abiotic factors in the physiological processes of fish (Lizama & Ambrosio, 2002) <sup>[14]</sup>.

The bronze featherback, or Asian knifefish, *Notopterus notopterus* (Pallas, 1769), is a species of fish native to Southeastern Asia. Its natural habitats are located in Pakistan, India, Bangladesh, Thailand, Myanmar, Malaysia, Laos, the Philippines, Cambodia, Vietnam, and Indonesia (Achakzai *et al.*, 2015; Kyaw *et al.*, 2020; Mohanty & Samanta, 2016; Mustafa *et al.*, 2014; Naeem *et al.*, 2010; Winn *et al.*, 2021) <sup>[1, 12, 15, 22, 23, 31]</sup>. In Indonesia, it has been documented in South Sumatra, Bangka Island, Jambi, Riau, Riau Island, and Kalimantan (Mulyani & Budijono, 2014;

Muslim & Syaifudin, 2022; Rapita *et al.*, 2021; Wibowo *et al.*, 2009) <sup>[17, 19, 28, 30]</sup>. The species is reported to inhabit rivers, floodplains, reservoirs, and lakes (Achakzai *et al.*, 2015; Mulyani & Budijono, 2014; Muslim & Syaifudin, 2022; Naeem *et al.*, 2010; Rapita *et al.*, 2021) <sup>[1, 17, 19, 23, 28]</sup>. It feeds on fish, crustaceans, insects, microalgae, nematodes, annelids, macrophytes, detritus, and some young roots of aquatic plants (Achakzai *et al.*, 2015; Rapita *et al.*, 2021) <sup>[1, 28]</sup>. This fish can easily adjust to the conditions of cultivation. As a result, it may one day become a commodity for aquaculture (Muslim *et al.*, 2020) <sup>[18]</sup>. This species is adaptable in aquariums, box containers, and buckets (Muslim *et al.*, 2023; Muslim & Simanjuntak, 2023) <sup>[21, 20]</sup>. The present study was designed to explore length- frequency distributions, length-weight relationships, and condition factors of *N. notopterus* in the East Pedamaran Floodplain, Ogan Komering Ilir, South Sumatra. Findings from the present study will significantly contribute to the successful management of *N. notopterus* to preserve it from possible depletion in the wild.

### Materials and Methods

This research was conducted in the Floodplain area of Kayu Labu Village, East Pedamaran Sub-district, Ogan Komering Ilir District, South Sumatra, Indonesia (Fig 1). East Pedamaran Sub-district is one of subdistricts in Ogan Komering Ilir Regency which distance about 72 km to capital of regency. It is located in south east capital of regency. East Pedamaran Sub-district has 10 meters from surface of sea level, and total area is 671.11 km<sup>2</sup>.




**Fig 1:** Map of the study site for the length-weight relationship and condition factor of *Notopterus notopterus* (Pallas, 1769) from East Pedamaran Sub-district, Ogan Komering Ilir Regency, South Sumatra, Indonesia

A total of 108 *N. notopterus* were collected from local fishermen at the study site. Traditional fishing gear (gill nets, hand lines, bamboo traps, and fish barriers) was found to be used by fishermen to catch the fish. The fish samples were transported to the UPR Batanghari Sembilan Indralaya, and preserved in 10% formalin until the measurement was taken. In the laboratory, the total length (TL) and body weight (BW) of each specimen were measured to 0.1 cm and 0.1 g accuracy with calipers and an electronic balance, respectively.

The length-weight key of Soni & Ujjania (2017) [29] and the parabolic equation of Froese (2006) [7] were used for the statistical relationship between length and weight.  $W = a.L^b$ , where *W* and *L* are the body weight (g) and total length (cm) of the fish, and *a* and *b* are the regression and allometric constants, respectively. The equation has been transformed into the following logarithmic form:  $\text{Log } W = \log a + b \log L$ , and the values of ‘*a*’ and ‘*b*’ were determined empirically. In the present study, Fulton’s condition factor was determined using the formula:  $K =$

East Pedamaran Floodplain. The total number of *N. notopterus* samples collected was 108. The total length of the fish ranged from 19 to 26 cm ( $22.94 \pm 1.59$ ;  $\text{AVG} \pm \text{SD}$ ) and body weight ranged from 60 to 162 g ( $95.24 \pm 20.43$ ;  $\text{AVG} \pm \text{SD}$ ) (Fig 2). Four total length class intervals were obtained from the results of the length frequency distribution analysis. The most frequent total length class interval was 21-22.99 cm, and the least frequent was 19-20.99 cm. Seven body weight class intervals were also obtained. The most frequent body weight class groups were 90-104.99 g, the least frequent was 150-164.99 g (Table 1).




**Fig 2:** The total length and body weight of *Notopterus notopterus* (Pallas, 1769) from East Pedamaran Floodplain, Ogan Komering Ilir, South Sumatra, Indonesia

**Table 1:** Size composition of *Notopterus notopterus* collected from East Pedamaran Floodplain, Ogan Komering Ilir, South Sumatra, Indonesia

Class interval of body weight (g)	Frequency	%	Class interval of total length (cm)	Frequency	%
60.00-74.99	11	10.19	19.00-20.99	6	5.56
75.00-89.99	23	21.30	21.00-22.99	48	44.44
90.00-104.99	49	45.37	23.00-24.99	37	43.26
105.00-119.99	9	8.33	25.00-26.99	17	15.74
120.00-134.99	11	10.19			
135.00-149.99	3	2.78			
150.00-164.99	2	1.85			
Total	108	100		108	100

The length-weight relationship of *N. notopterus* in this study was formulated as  $W = 0.0948.L^{2.2012}$  (Fig 3). The total length and body weight of *N. notopterus* were positively correlated, with a coefficient of determination ( $R^2$ ) of 0.5332. The t-test results showed that the regression coefficient (b) was significantly different from 3 ( $P > 0.05$ ). The b value of 2.22 means the value of  $b < 3$ , indicating a negative allometric growth pattern of *N. notopterus*.



**Fig 3:** The length-weight relationship of *Notopterus notopterus* (Pallas, 1769) from East Pedamaran Floodplain, Ogan Komering Ilir, South Sumatra, Indonesia

The study of fish length-weight relationships is an essential tool for illustrating some biological aspects of species (Freitas *et al.*, 2017) [6]. It is used to estimate the weight corresponding to a given length, and condition factors are used to compare the "condition", "fatness", or "health" of fish (Basak & Hadiuzzaman, 2019; Froese, 2006; Soni & Ujjania, 2017) [3, 7, 29]. The condition factor indicates the fitness of the species, with the assumption being whether the species retains a reasonable balance between length and weight. The heavier fish of a certain length are in better condition.

The total length of *N. notopterus* sampled in the current study ranged from 19 to 26 cm ( $22.94 \pm 1.59$ ;  $AVG \pm SD$ ) and body weight ranged from 60 to 162 g ( $95.24 \pm 20.43$ ;  $AVG \pm SD$ ). In a previous study, *N. notopterus* samples from Pokoriya River, Morigaon, Assam India sized 15.4-26.1 cm (TL), 20-142 g (BW) (Kaushik *et al.*, 2019) [10], Sukhna Lake, Chandigarh India sized 21-28.3 cm, 60-240 g (Kaur & Rawal, 2017) [9], Brahmaputra River, Dhuburi, Assam India sized 17.2-25.6 cm, 48.7-151.2 g (Paul *et al.*, 2022) [27], Indus River, southern Punjab, Pakistan sized 14.4-29.4 cm (Naeem *et al.*, 2010) [23], Manchar Lake Sindh Pakistan sized 10-30 cm, 16.5-290 g (Achakzai *et al.*, 2015) [1], Kaptai Reservoir Bangladesh sized 10-33.25 cm (Mustafa *et al.*, 2014) [22], Sunye Lake, Mandalay Region, Myanmar sized 17.1-38.7 cm, 34-420 g (Winn *et al.*, 2021) [31], Sei Gesek Reservoir Indonesia sized 16.9-27 cm (Rapita *et al.*, 2021) [28], Sail River Indonesia sized 13.5-27 cm (Mulyani & Budijono, 2014) [17].

The results of this study show that the growth pattern of *N. notopterus* is negatively allometric. These results are similar to research conducted by Naeem *et al.*, (2010) [23] in the Indus River, Pakistan; the growth pattern of *N. notopterus* is negative allometric. However, in contrast to the results of Achakzai *et al.*, (2015) [1], the growth pattern of *N. notopterus* from Manchar Lake, Sindh, Pakistan, is isometric; from Pokoriya River, Morigaon, Assam, and

Sukhna Lake, Chandigarh, it is positive allometric growth (Kaur & Rawal, 2017; Kaushik *et al.*, 2019) [9, 10]. Negative allometric growth implies that the fish becomes leaner with increasing weight, while positive allometric growth implies that the fish becomes relatively fatter or deeper-bodied with increasing length (Fafioye & Ayodele, 2018) [4]. The Fulton condition factor is a method that is generally used to assess the health index of the fish, indicating the nutritional and physiological status of the fish (Muchlisin *et al.*, 2015) [16]. According to Fafioye & Ayodele, (2018) [4] a low condition factor value indicates that the fish habitat is not fully suitable for fish health. The condition factor value of *N. notopterus* in the current study was 1.02, indicating that the fish were in good condition.

## Conclusions

The length-weight relationship of *N. notopterus* has a negative allometric growth pattern ( $b$  value  $2.84 < 3$ ). The condition factor value of 1.02 indicates that the *N. notopterus* population in the East Pedamaran Floodplain, Ogan Komering Ilir was in good condition.

## Conflict of Interest

The authors declare no conflict of interest.

## Acknowledgments

The authors would like to thank Mr. Agus and Mr. Yusuf for their valuable assistance in collecting fish samples. We also wish to acknowledge the fishermen for their cooperation in collecting samples.

## References

1. Achakzai WM, Baloch WA, Soomro AN. Length-weight relationship and condition factor of *Notopterus notopterus* (Pallas, 1769) from Manchar Lake Sindh, Pakistan. *Sindh University Research Journal (Science Series)*. 2015; 47(3):515-518.
2. Augustina AXT, Sreeram MP, Sreekumar KM, Aju KR. Length-weight relation and relative condition factor of six deep-sea fishes from the south-eastern Arabian Sea. *Journal of the Marine Biological Association of India*. 2022; 64(2):118-122. Doi: <https://doi.org/10.6024/jmbai.2022.64.2.2336-19>
3. Basak SS, Hadiuzzaman M. Length-weight relationship (LWR), condition factor (K) and relative condition factor (KN) of Kalibaus fish *Labeo calbasu* (Hamilton, 1822) of Kaptai Lake, Rangamati, Bangladesh. *International Journal of Fisheries and Aquatic Studies*. 2019; 7(5):231-235.
4. Fafioye O, Ayodele O. Length-weight relationship and condition factor of four commercial fish species of Oyan Lake, Nigeria. *Examines in Marine Biology and Oceanography*. 2018; 2(2):227-230. Doi: <https://doi.org/10.31031/EIMBO.2018.02.000543>
5. Ferdaushy MH, Alam MM. Length-length and length-weight relationships and condition factor of nine freshwater fish species of Nageshwari, Bangladesh. *International Journal of Aquatic Biology*. 2015; 3(3):149-154.
6. Freitas TMD, Souza JBD, Prudente BD, Montag LFD. Length-weight relationship in ten fish species from the Nhamundá River, the Amazon Basin, Brazil. *Acta Amazonica*. 2017; 47(1):75-78.
7. Froese BR. Cube law, condition factor and weight-

- length relationships: History, meta-analysis and recommendations. *Journal of Applied Ichthyology*. 2006; 22:241-253. Doi: <https://doi.org/10.1111/j.1439-0426.2006.00805.x>
8. Ighwela KA, Ahmed A, Bin, Abol-Munafi A. Condition factor as an indicator of growth and feeding intensity of Nile tilapia fingerlings (*Oreochromis niloticus*) fed on different levels of maltose. *American-Eurasian Journal of Agricultural & Environmental Sciences*. 2011; 11(4):559-563.
  9. Kaur V, Rawal YK. Length-weight relationship (LWR) in *Notopterus notopterus* (Pallas) from Sukhna Lake, Chandigarh. *IOSR Journal of Pharmacy and Biological Sciences*. 2017; 12(4):63-65. Doi: <https://doi.org/10.9790/3008-1204046365>
  10. Kaushik KK, Sahu P, Nath R. Length-weight relationship of *Notopterus notopterus* (Pallas, 1769), *Anabas testudineus* (Bloch, 1792) and *Clarias batrachus* (Linnaeus, 1758) from Pokoriya River, Morigaon, Assam, India. *International Journal of Scientific and Technology Research*. 2019; 8(10):3087-3088. Doi: <https://doi.org/10.2307/1540>
  11. Kumari S, Sandhya KM, Karnatak G, Lianthuamluaia, Sarkar UK, Panda D, *et al.* Length-weight relationship and condition factor of *Gudusia chapra* (Hamilton, 1822) from Panchet Reservoir, Jharkhand, India. *Indian Journal of Fisheries*. 2019; 66(3):136-139. Doi: <https://doi.org/10.21077/ijf.2019.66.3.81017-18>
  12. Kyaw MM, Kyaw MM, Lum MR, Naing SSS. Relationship of ecological factors and commercial bronze featherback fish, *Notopterus notopterus* (Pallas, 1769); Irrawaddy River, Myitkyina Segment, Kachin State, Myanmar. *University of Mandalay Research Journal*. 2020; 11:98-111.
  13. Lima JS, Costa I, Zalmon I. Length-weight relationship of fish species captured around an artificial offshore reef (northern Rio de Janeiro, Brazil). *Journal of Applied Ichthyology*. 2021; 37(2):337-341. Doi: <https://doi.org/10.1111/jai.14109>
  14. Lizama M, de los AP, Ambrosio AM. Condition factor in nine species of fish of the characidae family in the upper Paraná River Floodplain, Brazil. *Brazilian Journal of Biology*. 2002; 62(1):113-124.
  15. Mohanty D, Samanta L. Chemosphere Multivariate analysis of potential biomarkers of oxidative stress in *Notopterus notopterus* tissues from Mahanadi River as a function of concentration of heavy metals. *Chemosphere*. 2016; 155:28-38. Doi: <https://doi.org/10.1016/j.chemosphere.2016.04.035>
  16. Muchlisin ZA, Batubara AS, Siti-azizah MN, Adlim M, Hendri A, Fadli N, *et al.* Feeding habit and length weight relationship of keureling fish, *Tor tambra Valenciennes*, 1842 (Cyprinidae) from the western region of Aceh Province, Indonesia. *Biodiversitas*. 2015; 16(1):89-94. Doi: <https://doi.org/10.13057/biodiv/d160112>
  17. Mulyani I, Budijono B. Morphometric and meristic analysis of Asian knifefish (*Notopterus notopterus*) in Sail River, Pekanbaru Riau Province. *Jurnal Ilmiah Biologi Eksperimen dan Keanekaragaman Hayati*. 2014; 7(2):59-64.
  18. Muslim M, Heltonika B, Sahusilawane H, Wardani WW, Rifai R. Ikan lokal perairan tawar Indonesia yang prospektif dibudidayakan. Purwokerto: Pena Persada, 2020.
  19. Muslim M, Syaifudin M. Biodiversity of freshwater fish in Kelekar Floodplain Ogan Ilir Regency in Indonesia. *Journal of Tropical Biodiversity and Biotechnology*. 2022; 7(1):p67494. Doi: <https://doi.org/10.22146/jtbb.67494>.
  20. Muslim M, Simanjuntak WJ. Growth and survival of bronze featherback (*Notopterus notopterus*, Pallas 1769) reared on bucket. *Magna Scientia Advanced Research and Reviews*. 2023; 9(1):101-105. Doi: <https://doi.org/10.30574/msarr.2023.9.1.0133>.
  21. Muslim M, Pitriani E, Agustina H. Growth and survival of bronze featherback (*Notopterus notopterus*) adapted on box container and aquarium. *GSC Advanced Research and Review*. 2023; 16(3):133-137. Doi: <https://doi.org/10.30574/gscarr.2023.16.3.0368>
  22. Mustafa MG, Singha S, Islam M, Mallick N. Population dynamics of *Notopterus notopterus* (Pallas, 1769) from the Kaptai Reservoir of Bangladesh. *SAARC Journal of Agriculture*. 2014; 12(2):112-122. Doi: <https://doi.org/10.3329/sja.v12i2.21920>
  23. Naeem BM, Salam A, Gillani Q, Ishtiaq A. Length-weight relationships of *Notopterus notopterus* and introduced *Oreochromis niloticus* from the Indus River, southern Punjab, Pakistan. *Journal of Applied Ichthyology*. 2010; 26:p620. Doi: <https://doi.org/10.1111/j.1439-0426.2010.01480.x>
  24. Orlov A, Binohlan C. Length-weight relationships of deep-sea fishes from the western Bering Sea. *Journal of Applied Ichthyology*. 2019; 25:223-227. Doi: <https://doi.org/10.1111/j.1439-0426.2009.01215.x>
  25. Ouahb S, Bousseba M, Ferraj L, Moujtahid AEL, Hasnaoui M. Weight-length relationship and relative condition factor of *Micropterus salmoides* (Lacépède, 1802), *Cyprinus carpio* (Linnaeus, 1758) and *Oreochromis niloticus* (Linnaeus, 1758) caught in the Al-Massira Dam Lake. *E3S Web of Conferences*. 2021; 314:p01005.
  26. Pathak AK, Kantharajan G, Dayal R, Kumar R, Saini VP, Mohindra V, *et al.* Length-weight relationships and relative condition factor of fish species dwelling in the inland saline river Luni, India, with a note on the associated risks and encounters obstructing the fish diversity and ecosystem services. *Indian Journal of Fisheries*. 2022; 69(4):59-64. Doi: <https://doi.org/10.21077/ijf.2022.69.4.127612-07>
  27. Paul MG, Nath P, Dutta A. Length-weight relationship, relative condition factor and fecundity of *Notopterus notopterus* (Pallas, 1769) from river Brahmaputra in Dhubri, Assam, India. *Journal of Fisheries*. 2022; 10(3):p103201. Doi: <https://doi.org/10.17017/j.fish.398>
  28. Rapita R, Susiana S, Kurniawan D, Lestari F, Sabriaty D, Rianti U. Food habits of belida fish (*Notopterus notopterus*, Pallas 1769) in Sei Gesek Reservoir, Bintan Regency, Riau Island, Indonesia. *IOP Conference Series: Earth and Environmental Science*. 2021; 919:p012003. Doi: <https://doi.org/10.1088/1755-1315/919/1/012003>
  29. Soni N, Ujjania NC. Length-weight relationship and condition factor of Indian major carps of Vallabhsagar reservoir, Gujarat, India. *Indian Journal of Fisheries*. 2017; 64(76263):186-189. Doi:

<https://doi.org/10.21077/ijf.2017.64.special-issue.76263-27>

30. Wibowo A, Sunarno MT, Subagja S, Hidayah T. Population characteristic of the putak (*Notopterus notopterus*) using fenotipic diversity analysis and 16sRNA region mitochondria DNA. *Jurnal Penelitian Perikanan Indonesia*. 2009; 15(1):1-12.
31. Winn NA, Sandi P, Khaing T, Nyunt KT, Kyaw HT, Sabai M, *et al.* Length weight relationship of twelve freshwater fish species from Sunye Lake, Mandalay Region, Myanmar. *Greener Journal of Biological Sciences*. 2021; 11(2):74-80.
32. Zuchi N, Röpke C, Shibuya A, Farago T, Carmona M, Zuanon J, *et al.* Length-weight relationship of fish species from Central Amazon floodplain. *Journal of Applied Ichthyology*. 2020; 36(6):837-841. Doi: <https://doi.org/10.1111/jai.14082>