



Rozirwan unsri &lt;rozirwan@unsri.ac.id&gt;

---

## [STI] Submission Acknowledgement

1 pesan

---

**Prof. Aldes Lesbani, Ph.D** <scitechindones@gmail.com>  
Kepada: "Dr.Rozirwan, S.Pi., M.Sc." <rozirwan@unsri.ac.id>

21 Februari 2024 pukul 08.54

Dear Dr.Rozirwan, S.Pi., M.Sc.:

Thank you for submitting your manuscript entitled "Anti-inflammatory activity and phytochemical profile from the leaves of the mangrove *Sonneratia caseolaris* (L.) Engl. for future drug discovery" to Science and Technology Indonesia. Now, your manuscript will be considered by the editor and section editor before further peer-review process. With the online journal management system that we are using, you will be able to track its progress through the editorial process by logging in to the journal website:

Submission URL: <https://scitechindonesia.com/index.php/jsti/authorDashboard/submission/1163>

Username: rozirwan

If you have any questions, please contact me. Thank you for considering this journal as a venue for your work.

### Science and Technology Indonesia

*A Peer-Reviewed Research Journal of Science and Technology*

p-ISSN: 2580-4405 | e-ISSN: 2580-4391

**E-mail:** [admin@scitechindonesia.com](mailto:admin@scitechindonesia.com) | [scitechindonesia@gmail.com](mailto:scitechindonesia@gmail.com)

**Homepage:** <http://scitechindonesia.com/index.php/jsti>



Rozirwan unsri &lt;rozirwan@unsri.ac.id&gt;

## [STI] Editor Decision

1 pesan

**Prof. Aldes Lesbani** <scitechindones@gmail.com>

4 Maret 2024 pukul 13.40

Kepada: "Dr.Rozirwan, S.Pi., M.Sc." <rozirwan@unsri.ac.id>, Ade Siswanto <adeeafhd@gmail.com>, Nadila Nur Khotimah <nadilakhotimah1142@gmail.com>, Redho Yoga Nugroho <redhoyn.29@gmail.com>, Wike Ayu Eka Putri <wike\_ayu\_ep@unsri.ac.id>, Fauziyah <siti\_fauziyah@yahoo.com>, Rezi Apri <rezi\_apri@unsri.ac.id>, Hartoni <hartoniunsri@gmail.com>

Dear Dr.Rozirwan, S.Pi., M.Sc., Ade Siswanto, Nadila Nur Khotimah, Redho Yoga Nugroho, Wike Ayu Eka Putri, Fauziyah, Rezi Apri, Hartoni:

Reviewers have now commented on your paper (attached below this email). You will see that they are advising that you must revise your manuscript. If you are prepared to undertake the work required, I would be pleased to reconsider my decision.

For your guidance, reviewers' comments can be read in your Author online interface. If you decide to revise the work, please submit a list of changes or a rebuttal against each point which is being raised when you submit the revised manuscript.

\*\*\*\*>>>Please be noted that you have up to **2 (two) weeks from now** to revise your manuscript, unless your manuscript will be considered as a new submitted manuscript. If you need additional time to complete your revision, please let us know by replying to this email and informing us of the date you expect to submit it. <<<<\*\*\*\*

To submit a revision, please upload your revised manuscript documents to SCIENCE AND TECHNOLOGY INDONESIA online submission interface at (<https://sciencetechindonesia.com>) after you log in as Author. Important: Please indicate the revision as RED-highlighting the revision sentences or words within your revised manuscript.

-----  
The revised document files (three files) MUST include:

- One (1) file of Revision Note file in a table form with respect to Reviewers comments including the location of the revision on the revised manuscript.
- One (1) file of Revised Manuscript file according to Template-based format (MS Word file) (please **RED**-color highlighted texts in the revised sentences).
- One (1) file of Graphical Abstract file according to Guideline of Graphical Abstract

-----  
GUIDELINE TO UPLOAD REVISION: To upload your revised manuscript, please login to Science and Technology Indonesia journal submission interface (<https://sciencetechindonesia.com>) login using your user and password as usual. **Therefore, click on "Review" and upload to "revision box"**.

For second and continuing order of revision of manuscript, please just add your latest version file(s), do not delete the previous round of file(s) of manuscript.

IMPORTANT NOTE: A GRAPHICAL ABSTRACT of each article is mandatory for this journal. This graphic should capture the reader's attention and, in conjunction with the manuscript title, should give the reader a quick visual impression of the essence of the manuscript without providing specific results.

Choosing/Creating a Graphical Abstract: The graphic should be simple, but informative; The use of color is encouraged; The graphic should uphold the standards of a scholarly, professional publication; The graphic must be entirely original, unpublished artwork created by one of the coauthors; The graphic should not include a photograph, drawing, or caricature of any person, living or deceased; Do not include postage stamps or currency from any country, or trademarked items (company logos, images, and products); Avoid choosing a graphic that already appears within the text of the manuscript. Image size: please provide an image with a size of 500 × 800 pixels (height × width). Preferred file types: TIFF, JPG, PNG, PDF, or MS Office files.

The Graphical Abstracts should be submitted as a separate file during the revised manuscript submission system or otherwise, it can be submitted by email to: [sciencetechindonesia@gmail.com](mailto:sciencetechindonesia@gmail.com)

Thank you for submitting to this journal.

## Science and Technology Indonesia

*A Peer-Reviewed Research Journal of Science and Technology*

p-ISSN: 2580-4405 | e-ISSN: 2580-4391

E-mail: [admin@sciencetechindonesia.com](mailto:admin@sciencetechindonesia.com) | [sciencetechindonesia@gmail.com](mailto:sciencetechindonesia@gmail.com)

Homepage: <http://sciencetechindonesia.com/index.php/jsti>

---

### 3 lampiran



**A-review-sti-1163.docx**

16K



**B-STI-1163 Rvw.pdf**

56K



**C-Review 1163 STI.docx**

14K

### Table of Responses

No.	Questions	Answers
<b>A-review-sti-1163</b>		
<b>1.</b>	Please add data on the presentation of inflammation test results that are more suitable for the interpretation of inflammation results.	Data presenting inflammatory test results have been presented in Figure 3 and explained in more detail in Table 2. (Page 26-28)
<b>2.</b>	Statistical analysis must be included in the graphic	The results of the statistical analysis have been added to the graph (Figure 3).
<b>3.</b>	Please discuss in detail about the GC-MS results on the anti-inflammatory test.... In your manuscript there are many components from the GC-MS results... What is the effect of these components for anti-inflammatory?	<p>GC-MS results show that there are various compounds in the ethanol extract of <i>S. caseorolis</i> leaves that have anti-inflammatory potential, supported by references from several previous studies, including aldehydes, glucosinolates, coumarins, esters, terpenoids, alcohols, lipids, tocopherols, and steroids. Most of these compounds have been associated with anti-inflammatory activity in previous studies.</p> <p>For example, some aldehyde compounds, such as nonanal, have been shown to have potent anti-inflammatory activity. In addition, glucosinolates such as desulphosininigrin and 4-methylsulfinylbutyl glucosinolate also have anti-inflammatory effects as well as antimicrobial and antioxidant activities. Coumarin compounds, such as 5,8-Epoxy-3H-2-benzopyran, have anti-inflammatory potential associated with inhibition of prostaglandin production and anticoagulant effects. Esters, such as hexadecanoic acid, methyl esters, and terpenoid compounds such as phytol, have been studied for their anti-inflammatory activity. Likewise, alcohol compounds</p>



		<p>such as 1-hexadecanol have also been found to have anti-inflammatory properties.</p> <p>Tocopherols, like <math>\alpha</math>-tocopheryl acetate, are known for their antioxidant properties, which can help fight inflammation by neutralizing free radicals. Additionally, steroid compounds such as <math>\zeta</math>-sitosterol have also been found to have significant anti-inflammatory effects in certain tissues.</p> <p>Thus, the combination of various compounds found in <i>S. caseoralis</i> leaf extract has the potential to produce powerful anti-inflammatory effects. Their effect on inflammation may vary depending on the mechanism of action of each compound and the interactions between them. Therefore, the use of <i>S. caseoralis</i> leaves as a potential source for traditional medicine in the future can provide an effective alternative in the treatment of inflammatory diseases and contribute to the maintenance of public health (Page 11-13).</p>
4.	<p>What components/compounds have a good effect related to the anti-inflammatory test from the components of the GC-MS test results</p>	<p>Of the various compounds found based on supporting references and the various mechanisms of action involved, coumarin is a promising compound for reducing inflammation and has the potential to be used in the treatment of inflammatory diseases (Page 12).</p>
5.	<p>Please pay attention to comments no. 3-4 with abstracts. abstracts are too general,,,,,should be specific about the results.</p>	<p>The results presented in the abstract are specific because adding compound information components or compounds has a good effect on the anti-inflammatory test (Page 1).</p>
6.	<p>What is 150 mg/kgBB? isn't that Indonesian? please check it carefully.</p>	<p>Indeed, "mg/kgBB" is a unit generally used in Indonesia. We have changed it to international units, namely "mg/kg body weight" or "mg/kg BW"</p>

7.	please separate between leaf collection with process and materials/instrumentation.	Leaf collection by process and materials/instrumentation has been separated into a new subchapter (page 3-4).
8.	Please check...GC-MS is not spectrum but Chromatogram (for GC) and Spectrum (for MS)> be careful to show the results.	Based on the GC-MS results, we have displayed the chromatogram results (Page 11).
9.	Did you obtain pure compound of your extract? please explain why you did not use pure compound from your extract	<p>In our study on the anti-inflammatory activity and phytochemical profile of <i>Sonneratia caseolaris</i> (L.) Engl. mangrove leaves, we indeed utilized extracts as the primary test material. This decision was based on several considerations. Firstly, leaf extracts contain a diverse range of chemical compounds that can provide holistic information regarding the potential anti-inflammatory activity. Secondly, leaf extracts naturally reflect the complexity of interactions among compounds within the plant matrix, which may not be fully represented by single compounds.</p> <p>While we acknowledge that isolating pure compounds could offer additional clarity on the specific compounds responsible for the observed activity, in the context of this research, our focus lies in comprehensively understanding the phytochemical contribution of <i>Sonneratia caseolaris</i> leaf extracts to anti-inflammatory potential. Furthermore, this choice aligns with our research goal of providing a broad foundation of information for the development of future medicines.</p>
<b>B-STI-1163 Rvw</b>		
1.	Abstract first sentence should state a background problem that the study tries to solve.	The first sentence of the abstract has been added to the research background on page 1.
2.	Chemicals and Instrumentation should be created in a new section in the Experimental	Chemicals and Instrumentation have been created in a new section in the

	Section. Indicate the source and purity of the chemicals. Indicate the country of origins and brand names of the instruments or any materials.	Experimental Section along with supporting information on page 3-4.
3.	In the Preparation of a Suspension of Na. CMC 0.5%, how many grams of Na. CMC used in this procedure. Please correct.	A total of 1.5 g Na. CMC used in making Na. CMC 0.5% on page 4.
4.	Please change ml to mL in your manuscript.	Every ml has been changed to mL.
5.	The x and y axes in Figure 3 still use commas in decimal numbers. Please pay attention again, I am telling you to use the origin application in processing graphs.	Figure 3 has been updated with a display without using commas in decimal numbers (Page 26-27).
6.	The quality of your Figure in Table 5 is very poor, please improve it.	The quality of the images in Table 5 has been improved (Page 30-32).
7.	I suggest the authors to mention the novelty of this work clearly in the introduction section.	The explanation regarding the novelty in this research has been corrected on page 3.



Rozirwan unsri &lt;rozirwan@unsri.ac.id&gt;

---

## Request Revision\_Round 2 (STI-1163)

1 pesan

---

**Aldes Lesbani** <sciencetechindonesia@gmail.com>  
Kepada: Rozirwan unsri <rozirwan@unsri.ac.id>

25 Maret 2024 pukul 09.12

Dear Author,

We hope you are doing well. We have checked the revision of your article, but there are still some parts that you need to improve. Please revise your article according to the comments below:

- i) This paper has a high similarity index when checked with Turnitin (attached file). Please improve and address this similarity index issue by lowering it to below 20%.
- ii) All Figures, Tables, and Equations should be mentioned in the text of the manuscript.
- iii) Figures 3 and 4 do not match the journal format, please refer to the journal template.
- iv) The quality of Figure in Table 5 should be improved

Please revise your article and submit it before **March 30, 2023**. If you have any questions, please do not hesitate to contact us. Thank you for your cooperation

Sincerely Yours,

Editor-in-Chief

**Prof. Aldes Lesbani, Ph.D.**

Science &amp; Technology Indonesia

<http://sciencetechindonesia.com>

---

 **STI 1163.pdf**  
6758K



Rozirwan unsri &lt;rozirwan@unsri.ac.id&gt;

## Request Revision\_Round 3 (STI-1163)

1 pesan

**Aldes Lesbani** <sciencetechindonesia@gmail.com>  
Kepada: Rozirwan unsri <rozirwan@unsri.ac.id>

30 Maret 2024 pukul 13.11

Dear Author,

We hope you are doing well. We have checked the revision of your article, but there are still some parts that you need to improve. Please revise your article according to the comments below:

- i) Please correct Figure 3 in your manuscript. Please use lower case in the captions a) and b). Pay attention to the description of the x-axis, because it still uses uppercase and lowercase letters in hours.
- ii) Please pay attention to self-citation in your manuscript. According to our journal rules, you can only self-cite a maximum of 10% of the total number of references.
- iii) The quality of Figure in Table 5 should be improved

Please revise your article and submit it before **2 April 2023**, if you wish to be **included in the April issue**. If you have any questions, please do not hesitate to contact us. Thank you for your cooperation

Sincerely Yours,


Editor-in-Chief

**Prof. Aldes Lesbani, Ph.D.**

Science &amp; Technology Indonesia

<http://sciencetechindonesia.com>

### 2 lampiran



**Figure 3a STI 1163.jpg**  
71K



**revision2-article-dr.rozirwan-sti.docx**  
1479K



Rozirwan unsri &lt;rozirwan@unsri.ac.id&gt;

---

**[STI] Editor Decision**

1 pesan

**Prof. Aldes Lesbani** <scitechindones@gmail.com>

1 April 2024 pukul 11.08

Kepada: Rozirwan <rozirwan@unsri.ac.id>, Ade Siswanto <adeeafhd@gmail.com>, Nadila Nur Khotimah <nadilakhotimah1142@gmail.com>, Redho Yoga Nugroho <redhoyn.29@gmail.com>, Wike Ayu Eka Putri <wike\_ayu\_ep@unsri.ac.id>, Fauziah <siti\_fauziah@yahoo.com>, Rezi Apri <rezi\_apri@unsri.ac.id>, Hartoni <hartoniunsri@gmail.com>

Dear Rozirwan, Ade Siswanto, Nadila Nur Khotimah, Redho Yoga Nugroho, Wike Ayu Eka Putri, Fauziah, Rezi Apri, Hartoni:

We have reached a decision regarding your submission to Science and Technology Indonesia, "Anti-inflammatory activity and phytochemical profile from the leaves of the mangrove *Sonneratia caseolaris* (L.) Engl. for future drug discovery".

Our decision is to accept your submitted manuscript for publication in

Thank you for publishing with us and please do not hesitate to contact us if you have any inquiry.

**Science and Technology Indonesia**

*A Peer-Reviewed Research Journal of Science and Technology*

p-ISSN: 2580-4405 | e-ISSN: 2580-4391

**E-mail:** [admin@scitechindonesia.com](mailto:admin@scitechindonesia.com) | [scitechindonesia@gmail.com](mailto:scitechindonesia@gmail.com)

**Homepage:** <http://scitechindonesia.com/index.php/jsti>

**D-Revision3\_Article\_Dr.Rozirwan\_STI.docx**

1767K



Rozirwan unsri &lt;rozirwan@unsri.ac.id&gt;

---

## Proofread and Invoice Article (STI-1163)

1 pesan

---

**Aldes Lesbani** <sciencetechindonesia@gmail.com>  
Kepada: Rozirwan unsri <rozirwan@unsri.ac.id>

2 April 2024 pukul 07.25

Dear Author,

I am sending the proof of the manuscript for your approval and final check before publishing it in Science and Technology Indonesia. If anything needs to be changed, please inform us as soon as possible.

*References marked in red are not in the reference list of your manuscript, please complete them*

To cover processing costs and provide open access for articles that have been accepted, the Journal now charges a publication fee of 3,700,000 IDR. This publication fee should be transferred to the bank account shown below, and details of the transfer either e-mailed to <admin@sciencetechindonesia.com> and <sciencetechindonesia@gmail.com>.

The payment must be made to Science and Technology Indonesia to the below bank account.

### Bank Transfer/Credit card To

**Bank: Bank Central Asia (BCA)**  
**Account Name: Aldes Lesbani**  
**Account No: 0214167855**  
**Bank SWIFT Code: CENAIJJA**

Your accepted manuscript will be transferred to the production section after your payment has been confirmed by our administration department.

Sincerely Yours,

Editor-in-Chief

**Prof. Aldes Lesbani, Ph.D.**

Science &amp; Technology Indonesia

<http://sciencetechindonesia.com>

---

### 2 lampiran

 **STI\_template.pdf**  
9391K **STI-INVOICE ARTICLE (1163).pdf**  
19K



## ARTICLE INVOICE

Invoice Date: 02/04/2024

Due Date: 05/04/2024

### Science and Technology Indonesia

Research Centre of Inorganic Materials and Coordination Complexes  
Magister Program of Materials Science, Graduate School  
Universitas Sriwijaya  
Jl. Padang Selasa, Bukit Besar, Palembang, 30139  
South Sumatra-Indonesia

### Bill To:

**Rozirwan**

rozirwan@unsri.ac.id

Reference Number: STI-1163-Manuscript

Item	Description	Quantity	Unit Price	Amount
1	Anti-Inflammatory Activity and Phytochemical Profile from the Leaves of the Mangrove <i>Sonneratia caseolaris</i> (L.) Engl. for Future Drug Discovery	1	Rp. 3,700,000	Rp. 3,700,000
			Sub Total	Rp. 3,700,000
			Total	Rp. 3,700,000

### Payment Address:

Bank: Bank Central Asia (BCA)  
Account Name: Aldes Lesbani  
Account No: 0214167855  
Bank SWIFT Code: CENAIJJA

We take your privacy seriously. We do not share your personal information with any third parties.

---



## Transfer Berhasil

**Rp3.700.000**

Nomor Referensi 20240402101710154655  
Tanggal Transaksi 2-April-2024  
Waktu Transaksi 10:17:10 WIB

Layanan Transfer Realtime Online  
Bank Tujuan BCA  
Nomor Rekening 0214167855  
Nama Penerima ALDES LESBANI, DR

Nama Pengirim ROZIRWAN  
Rekening Debet 06\*\*\*\*\*4

Berita Publication fee RF ST1163  
Manuscript Antiinflammasia

Nominal Rp3.700.000  
Biaya Admin Rp6.500

**Total Rp3.706.500**



## ARTICLE INVOICE

Invoice Date: 02/04/2024

Due Date: 05/04/2024

### Science and Technology Indonesia

Research Centre of Inorganic Materials and Coordination Complexes  
Magister Program of Materials Science, Graduate School  
Universitas Sriwijaya  
Jl. Padang Selasa, Bukit Besar, Palembang, 30139  
South Sumatra-Indonesia

### Bill To:

**Rozirwan**

rozirwan@unsri.ac.id

Reference Number: STI-1163-Manuscript

Item	Description	Quantity	Unit Price	Amount
1	Anti-Inflammatory Activity and Phytochemical Profile from the Leaves of the Mangrove <i>Sonneratia caseolaris</i> (L.) Engl. for Future Drug Discovery	1	Rp. 3,700,000	Rp. 3,700,000
			Sub Total	Rp. 3,700,000
			Total	Rp. 3,700,000

### Payment Address:

Bank: Bank Central Asia (BCA)  
Account Name: Aldes Lesbani  
Account No: 0214167855  
Bank SWIFT Code: CENAIJJA

We take your privacy seriously. We do not share your personal information with any third parties.



Rozirwan unsri &lt;rozirwan@unsri.ac.id&gt;

---

**Re: Proofread and Invoice Article (STI-1163)**

1 pesan

---

**Aldes Lesbani** <sciencetechindonesia@gmail.com>  
Kepada: Rozirwan unsri <rozirwan@unsri.ac.id>

2 April 2024 pukul 09.48

Dear Author,

We have received your publication payment and included your article in the Upcoming Edition (April 2024) section which will be published soon, we attach your final article below. Thank you for your cooperation.

Sincerely Yours,

Editor-in-Chief

**Prof. Aldes Lesbani, Ph.D.**

Science &amp; Technology Indonesia

<http://sciencetechindonesia.com>

Pada Sel, 2 Apr 2024 pukul 10.29 Rozirwan unsri &lt;rozirwan@unsri.ac.id&gt; menulis:

Dear Editor,

Thank you for sending the proof of our manuscript for final approval before publishing it in Science and Technology Indonesia. We have reviewed the proof and made the necessary revisions to ensure its accuracy and completeness.

We have also completed the payment of the publication fee of 3,700,000 IDR.

Additionally, we have emailed the proof of payment and the revised manuscript as instructed.

Please let me know if there are any further steps required from our end. We appreciate your assistance and look forward to the publication of our manuscript in Science and Technology Indonesia.

Thank you  
Best regards,

Pada Sel, 2 Apr 2024 pukul 09.25 Aldes Lesbani &lt;sciencetechindonesia@gmail.com&gt; menulis:

Dear Author,

I am sending the proof of the manuscript for your approval and final check before publishing it in Science and Technology Indonesia. If anything needs to be changed, please inform us as soon as possible.

*References marked in red are not in the reference list of your manuscript, please complete them*

To cover processing costs and provide open access for articles that have been accepted, the Journal now charges a publication fee of 3,700,000 IDR. This publication fee should be transferred to the bank account shown below, and details of the transfer either e-mailed to <admin@sciencetechindonesia.com> and <sciencetechindonesia@gmail.com>.

The payment must be made to Science and Technology Indonesia to the below bank account.

**Bank Transfer/Credit card To****Bank: Bank Central Asia (BCA)****Account Name: Aldes Lesbani****Account No: 0214167855****Bank SWIFT Code: CENAIDJA**

Your accepted manuscript will be transferred to the production section after your payment has been confirmed by our administration department.

Sincerely Yours,

Editor-in-Chief

**Prof. Aldes Lesbani, Ph.D.**

Science & Technology Indonesia

<http://sciencetechindonesia.com>

--

**Dr. Rozirwan**

Head of Marine Bioecology Laboratory

Department of Marine Science

Faculty of Mathematics and Natural Sciences

Sriwijaya University

Jalan Raya Palembang-Prabumulih KM 32, Indralaya

Ogan Ilir, Sumatera Selatan, Indonesia, Pos Code: 30862

Email: [rozirwan@unsri.ac.id](mailto:rozirwan@unsri.ac.id), [rozirwan@gmail.com](mailto:rozirwan@gmail.com)

---

## 2 lampiran



**STI-INVOICE ARTICLE (1163) (1).pdf**

39K



**STI\_template.pdf**

9392K



Rozirwan unsri &lt;rozirwan@unsri.ac.id&gt;

---

**[STI] Editor Decision**

1 pesan

**Prof. Aldes Lesbani** <scitechindones@gmail.com>

2 April 2024 pukul 09.43

Kepada: Rozirwan <rozirwan@unsri.ac.id>, Ade Siswanto <adeeafhd@gmail.com>, Nadila Nur Khotimah <nadilakhotimah1142@gmail.com>, Redho Yoga Nugroho <redhoyn.29@gmail.com>, Wike Ayu Eka Putri <wike\_ayu\_ep@unsri.ac.id>, Fauziah <siti\_fauziah@yahoo.com>, Rezi Apri <rezi\_apri@unsri.ac.id>, Hartoni <hartoniunsri@gmail.com>

Dear Rozirwan, Ade Siswanto, Nadila Nur Khotimah, Redho Yoga Nugroho, Wike Ayu Eka Putri, Fauziah, Rezi Apri, Hartoni:

Your article "Anti-Inflammatory Activity and Phytochemical Profile from the Leaves of the Mangrove Sonneratia caseolaris (L.) Engl. for Future Drug Discovery," is has been pre-published in upcoming issue.

URL of the issue in progress is here: <https://scitechindonesia.com/index.php/jsti/>

Thank you for your great contribution.

Best regards

**Science and Technology Indonesia**

*A Peer-Reviewed Research Journal of Science and Technology*

p-ISSN: 2580-4405 | e-ISSN: 2580-4391

**E-mail:** [admin@scitechindonesia.com](mailto:admin@scitechindonesia.com) | [scitechindonesia@gmail.com](mailto:scitechindonesia@gmail.com)

**Homepage:** <http://scitechindonesia.com/index.php/jsti>

## Anti-Inflammatory Activity and Phytochemical Profile from the Leaves of the Mangrove *Sonneratia caseolaris* (L.) Engl. for Future Drug Discovery

Rozirwan<sup>1\*</sup>, Ade Siswanto<sup>1</sup>, Nadila Nur Khotimah<sup>2</sup>, Redho Yoga Nugroho<sup>1</sup>, Wike Ayu Eka Putri<sup>1</sup>, Fauziyah<sup>1</sup>, Rezi Apri<sup>1</sup>, Hartoni<sup>1</sup>

<sup>1</sup>Department of Marine Science, Faculty of Mathematics and Natural Sciences, Universitas Sriwijaya, Indralaya, South Sumatra, 30862, Indonesia

<sup>2</sup>Environmental Management Study Program, Graduate Program, Universitas Sriwijaya, Palembang, South Sumatra, 30139, Indonesia

\*Corresponding author: rozirwan@unsri.ac.id

### Abstract

The increasing demand for effective and natural anti-inflammatory agents prompts an investigation into the properties of *Sonneratia caseolaris* (L.) Engl., a plant traditionally used in medicine. This study aimed to explore the ability of *S. caseolaris* leaves extract to inhibit inflammation and accelerate wound healing. *S. caseolaris* leaves were collected from Tanjung Api-Api area, Banyuasin, South Sumatra. The method involved carrageenan induction in rat paws as an inflammatory model. The results showed that the most effective dose was found in the group with a dose of 150 mg/kg BW measured using a digital caliper and plethysmometer. The qualitative phytochemical test contain alkaloids, steroids, flavonoids, and phenols. The results GC-MS analysis were thought to contain aldehydes, glucosinolates, coumarins, esters, terpenoids, alcohols, lipids, tocopherols, and steroids. Due to their diverse range of mechanisms of action, coumarins show significant promise in mitigating inflammation and hold potential for treating inflammatory conditions. This study provides new insights related to the potential of *S. caseolaris* as a source of natural anti-inflammatory agents, supporting public comprehension regarding the utilization of traditional herbal remedies.

### Keywords

Anti-Inflammatory, *S. caseolaris*, Leaves, Phytochemical Profile, Drug Discovery

Received: 20 February 2024, Accepted: 1 April 2024

<https://doi.org/10.26554/sti.2024.9.2.502-516>

## 1. INTRODUCTION

Inflammation is a natural reaction to harm and irritation resulting from infection, allergic reactions, or exposure to chemicals (MacLeod and Mansbridge, 2016). In inflammatory situations, common signs include vasodilation, which causes a rise in temperature and results in red discoloration. Increased vascular permeability leads to swelling, while nociceptors are physically and chemically stimulated, causing pain. Disruption of tissue structure results in the loss of normal tissue function (de Araújo et al., 2016; Bai et al., 2021). Frequently used clinical treatments for inflammatory conditions involve the use of anti-inflammatory compounds, both nonsteroidal and steroidal in nature (Shaukat et al., 2023). The use of steroid drugs as anti-inflammatory agents is currently controversial due to their many side effects (Itoh et al., 2023; Tsujimae et al., 2024). Therefore, it is necessary to be expanded new agents with stronger anti-inflammatory activity but with fewer side effects.

Nature is the ultimate source of all living things, and its various medicinal plants have been a resource for obtaining cures for various diseases since the dawn of human civilization

(Kundu et al., 2023). Despite the continuous improvement of modern techniques through methods like combinatorial chemistry or computational drug design, over fifty percent of modern pharmaceuticals originate from medicinal plants (Wainwright et al., 2022). Medicinal plants play a crucial role as a natural reservoir for combating enduring illnesses, offering a diverse array of treatments for various diseases (Ríos and Andújar, 2020; Maldonado Miranda, 2021). Native medicinal plants are a new source of anti-inflammatory agents with more tolerable side effects (Nascimento et al., 2021).

Mangroves are arboreal species found in tropical and subtropical coastal areas influenced by tidal fluctuations, primarily situated within the belt spanning from the northern to southern tropics (Biswas and Biswas, 2021; Anu et al., 2024). Several bioactive compounds with diverse effects, including insecticidal, antioxidant, and antimicrobial properties, have been extracted from mangrove plants (Rozirwan et al., 2020, 2023b,a). Various biologically active compounds, including limonoids, alkaloids, glycosides, terpenoids, steroids, flavonoids, esters, phenols, lactones, aliphatic alcohols, amides, acids, aliphatic ketones, quinones, and benzodioxols, have been extracted from



different parts of mangrove trees, such as bark, stems, leaves, fruits, and seeds (Parthiban et al., 2023). The *Sonneratia* genus thrives continuously in coastal areas where land meets the sea, encountering difficulties like oxygen-deprived soil, increased salinity, and strong ultraviolet radiation throughout the year (Nasrin et al., 2021). *Sonneratia* adapted to its environment and metabolism in unique ways to deal with these problems. For example, it developed high osmotic pressure, adaptive root growth, and salt secretion mechanisms. These adaptation systems allow *Sonneratia* to survive harsh environmental conditions and produce unusual secondary metabolites (Liu et al., 2023). In addition, *Sonneratia caseolaris* is also one of the mangrove plant species known to have a variety of pharmacological activities.

Previous studies have shown that mangrove leaves contain various bioactive compounds. One of the compounds found was a triterpenoid compound, which has been demonstrated to possess properties that reduce inflammation (Kundu et al., 2022). In addition, flavonoid compounds are also found in mangrove leaves, which are also known to have strong antioxidant activity properties (Rozirwan et al., 2023a). A previous study indicated that the ethanol extract derived from *S. caseolaris* leaves contains a diverse range of phytochemicals, including phenolics, alkaloids, flavonoids, saponins, and tannins (Rahim and Abu Bakar, 2018). According to (Komakech et al., 2019), these compounds hold promise for inhibiting the function of proinflammatory enzymes, reduce the production of inflammatory mediators, and affect inflammatory signaling pathways.

However, the novelty of our work lies in a comprehensive exploration of the chemical compounds found in the leaves of *S. caseolaris*, aiming to identify previously undiscovered active components. This study merges ethnobotanical knowledge with modern scientific techniques, presenting a unique approach that is poised to contribute significantly to the advancement of novel, effective, and sustainable anti-inflammatory medicines. The exploration of novel compounds in *S. caseolaris* adds a distinctive dimension to the ongoing efforts in the field, promising innovative solutions to the challenges posed by inflammatory diseases.

## 2. EXPERIMENTAL SECTION

### 2.1 Materials

The materials and specifications used in this research are White Rats (*Rattus norvegicus*), Aquades, Seaweed Carrageenan, Carboxymethyl Cellulose Sodium (Shandong Yulong, China), Sodium Diclofenac (Phapros, Indonesia), Ethanol, HCL, and NaOH from Merck KGaA, Darmstadt, Germany. Mangrove leaf extract was obtained through the extraction process using a Biobase RE-301 Rotary Evaporator (China). Edema volume measurements were taken using the Orchid PLM01 Plethysmometer (Orchid Scientific, India) and a digital caliper with 6 carbon (Nankai, Japan Technology).

Identification, preparation, maceration, deconstruction, and antioxidant activity tests on samples were carried out at the Ma-

rine Bioecology Laboratory; phytochemical tests were carried out at the Agricultural Products Chemistry and Microbiology Laboratory; and screening of secondary metabolite compounds was carried out at the Integrated Testing Laboratory, Sriwijaya University. Mangrove identification refers to Giesen et al. (2007).

## 2.2 Method

### 2.2.1 Leaf Collection

This study was conducted in October 2023. The leaf samples used were *Sonneratia caseolaris* (L.) Engl. species obtained from the innermost zonation close to the sea with muddy areas lined up along the Tanjung Api-Api port in Banyuasin Regency, South Sumatra (Figure 1). Mangrove forests in the area are brackish forest ecosystems that grow in the area between land and sea waters (Rozirwan et al., 2023c). Mangroves also provide a unique and productive habitat for various types of flora and fauna, including fish, shrimp, crabs, birds, and other animals (Rozirwan et al., 2022).

### 2.2.2 Plant Maseration and Extraction

*S. caseolaris* mangrove leaves as much as 200 gr (dry weight) that have been mashed and macerated with ethanol solvent as much as 1 L (1: 5 b/v) for 2 x 24 hours. According to Rozirwan et al. (2023c), ethanol solvents that are polar are more effective for extracting secondary metabolites. The result of maceration in the form of a solution is then filtered with a filter. The macerated mixture was subjected to evaporation using a water bath set at 40°C until the solvent evaporated entirely, resulting in the formation of a paste-like substance (crude extract) from mangrove leaves. This crude extract was subsequently kept at ambient temperature.

### 2.2.3 Preparation of 1% Carrageenan Solution

The carrageenan solution was formulated by weighing 0.1 g of carrageenan, then dissolved with 10 mL of physiological saline (NaCl 0.9%), so that a 1% carrageenan solution was obtained (Fehrenbacher et al., 2012; Elfiruk et al., 2021).

### 2.2.4 Preparation of a Suspension of Na. CMC 0.5%

The preparation of 0.5% Na.CMC suspension refers to Liang et al. (2023), a total of 1.5 g Na. CMC is sprinkled evenly into a mortar that already contains 100 mL of heated distilled water, stirred until a clear gel is achieved, diluted with a little water, put into a 100-mL volumetric flask until a transparent mass is formed, and then added with distilled water until the mark line is reached.

### 2.2.5 Diclofenac Sodium Suspension Preparation

A total of 10 diclofenac sodium tablets (each tablet contains 50 mg diclofenac sodium) were weighed and then crushed. Diclofenac sodium tablet powder was weighed according to the dose calculation, then put into a 5 mL volumetric flask and suspended in 0.5% Na-CMC, and enough volume was added to make it 5 mL (Das et al., 2023).




Figure 1. Map of Mangrove Leaves Collection




Figure 2. Leaf of *Sonneratia caseolaris* (L.) Engl.




Figure 3. Reducing the Volume of Edema on the Soles of Rats' Feet. a) Plethysmometer, b) Digital Vernier Caliper

**2.2.6 Preparation of *S. caseolaris* Leaves Extract Suspension Doses of 100 mg/kg BW, 150 mg/kg BW, and 200 mg/kg BW**

Ethanol leaf extracts were prepared in suspension form with three different doses: 100 mg/kg BW, 150 mg/kg BW, and

200 mg/kg BW. Portions of 100 mg, 150 mg, and 200 mg of ethanol extract from *S. caseolaris* leaves were placed into a mortar. Then, 0.5% Na-CMC suspension was gradually added






Figure 4. Analysis Chromatogram of Mangrove Leaves Extract *S. caseolaris*

while crushing until a uniform mixture was obtained. Finally, the mixture was transferred into a 10 mL volumetric flask and diluted to the mark line with 0.5% Na-CMC suspension.

### 2.2.7 Anti-Inflammatory Activity Assay

In this study, the carrageenan induction method on the soles of rat feet was used as an experimental method which was modified, referring to (Farida et al., 2022). Test animals included as many as 15 rats, which were divided into 5 experimental groups. The experiment was repeated three times, using three rats in each repetition. During the testing period, rats were weighed and randomly assigned to different groups, including a negative control group, a positive control group, and a test group. Each test group consisted of three rats.

Before treatment, each rat was given a boundary mark on the hind paw joint so that when measuring using a plethysmometer, the position was always the same. As a first step before treatment, the volume of the rat's paw was measured to determine the initial volume ( $V_0$ ). Measurements were made using two measurement tools, namely a plethysmometer and a vernier caliper, with the aim of obtaining more accurate and comprehensive results. Combination: The use of these two tools allows for more complete measurements, especially if the object being measured involves changes in volume and length dimensions simultaneously.

Test animals were induced with 1% carrageenan in as much as 0.1 mL. The mice were then treated orally using a sonde at the third hour after the induction. The edema volume of the mice's feet was measured after treatment every 30 minutes for 5 hours. The volume of edema was determined based on the increase in the plethysmometer and Vernier caliper. Mice in each group were administered treatment according to the following procedure:

- The positive control group received diclofenac sodium at a dosage of 0.13 mg/kg BW.
- The group (negative control) was given Na. CMC 0.5% 0.5 mL (500 mg)/100 g BW.
- Test group 1 was given 100 mg/kg BW of *S. caseolaris* leaves ethanol extract.
- Test group 2 was given *S. caseolaris* leaves ethanol extract

150 mg/kg BW.

- Test group 3 was given 200 mg/kg BW of the ethanol extract of *S. caseolaris* leaves.

### 2.2.8 Qualitative Phytochemical Analysis

Qualitative phytochemical tests of *S. caseolaris* leaves extracts were carried out by qualitative methods as described in Harborne (1998), Bouabida and Dris (2022), and Olasunkanni et al. (2022) in the form of alkaloid, flavonoid, terpenoid/steroid, saponin, and tannin tests.

### 2.2.9 Gas Chromatography-Mass Spectroscopy (GC-MS) Analysis

An analysis was performed to identify the bioactive compound constituents present in the *S. caseolaris* leaf extract. The spectral graph from the analysis findings was cross-referenced with the data repository in Wiley Library 7 (Hossain et al., 2017; Rahim and Abu Bakar, 2018).

### 2.3 Data Analysis

The observed variable is the change in the volume of inflammation in the sole of the test animal's foot per unit of time that has been determined. The use of the formula (1) for percent inflammation refers to Su et al. (2011).

$$\text{Percent Inflammation} = \left( \frac{V_t - V_o}{V_o} \right) \times 100\% \quad (1)$$

Where  $V_o$  is the initial volume of the mouse paw and  $V_t$  represents the inflammation of the mouse paw after treatment.

### 2.4 Statistical Analysis

Data analysis with the one-way ANOVA test is used to see differences in the mean of data groups with more than two treatments (Rozirwan et al., 2023d; García-Suárez et al., 2021). If the results obtained are significantly different or show that the  $p$ -value  $< 0.05$  is considered significant, subsequently, the Tukey Honestly Significant Difference (HSD) test was employed to ascertain any disparities in the mean of each treatment using the IBM SPSS v.26 program.

### 3. RESULTS AND DISCUSSIONS

#### 3.1 Description of Mangrove Leaf

Leaves are the characteristic part of a mangrove species. In the process of identifying each mangrove species, morphological observation of leaf shape is very necessary to be able to determine the characteristics of *Sonneratia caseolaris* leaf shape (Figure 2). *S. caseolaris* has skinned leaves that are round in shape; the base of the yellow peduncle has no glands; and the leaf tip is rounded. The base of the leaf is narrowed, with a leaf width of 3–4 cm and a leaf length of 7–9 cm.

Various mangrove species are found along the Tanjung Api-api Port area, South Sumatra. The Tanjung Api-api area is an area affected by anthropogenic activities like agriculture, residential areas, and port operations can lead to alterations in the marine ecosystem (Rozirwan et al., 2024; Fitria et al., 2023). Mangroves can act as natural filters that help clean water of polluting substances such as heavy metals and excess nutrients (Rozirwan et al., 2020; Ivorra et al., 2021). It can protect the marine environment from pollution that may come from plantation and settlement activities. The *Sonneratia* genus is one of the mangroves that are in the leading and dominant zone around the area (Rozirwan et al., 2023d). *S. caseolaris* is a characteristic mangrove species that reproduces via non-viviparous means and serves as an essential element of mangrove ecosystems in the Indo-West Pacific area (Chen et al., 2008). According to Ragavan et al. (2014), the morphology of *S. caseolaris* leaves ranges from lanceolate to broadly elliptical, featuring either a pointed or rounded apex; its branches do not hang downwards; it blooms with 1–3 flowers; mature buds are elliptical and slender in the middle; and its fruits are relatively larger in size.

#### 3.2 Characteristics of Mangrove Leaves Extract

Percentage weight shrinkage of wet and dry samples of *Sonneratia caseolaris* leaves are summarized in Table 1. Shrinkage percentage of water content of *S. caseolaris* leaves have a shrinkage percentage of 69% and a sample weight of 30.9%. The moisture content serves as a measure of the water content within the leaves. Excessive moisture levels can diminish leaf quality as they promote microbial growth (Zheng et al., 2011; Ramses et al., 2020). According to Brendel (2021), The water content decreases as the plant ages. The solvent used in maceration is 70% ethanol, with the aim of attracting both polar and non-polar compounds (Batubara et al., 2020). Ethanol is an amphipathic solvent that can dissolve compounds that are both polar and nonpolar (Hikmawanti et al., 2021). Mangroves often contain various types of compounds with polar and nonpolar properties, so ethanol can effectively extract a large number of diverse bioactive compounds (Altemimi et al., 2017; Acquaviva et al., 2023).

#### 3.3 Anti-Inflammatory Activity

The graph of the lower edema volume shows that the 150 mg/kg BW dose of mangrove leaves extract works faster than the 100 mg/kg BW dose, the 200 mg/kg BW dose, and the

positive control (Figure 3). This is indicated by the decrease in the volume of edema of mangrove leaf extract dose 150 mg/kg BW continuously from 1 to 5 hours compared to other controls using both digital vernier caliper and plethysmometer. Based on Figure 3, the greatest anti-inflammatory effectiveness was obtained at a dose of 150 mg/kg BW leaf extract group using a digital vernier caliper and plethysmometer, which is effective for binding to receptors. The anti-inflammatory effect is evidenced by the lowest percentage of inflammation observed over the course of 5 hours.

*Sonneratia caseolaris* mangrove leaf extract has anti-inflammatory activity against carrageenan-induced test animals. According to Maehle (2009), The strength of drug effects corresponds to the receptors they interact with or bind to. Maximum effect occurs when all receptors are engaged by the drug, so increasing the dosage does not enhance the anti-inflammatory effect. The carrageenan test is commonly employed as a model for acute inflammation in experimental animals and exhibits high sensitivity to nonsteroidal anti-inflammatory drugs (Bouyahya et al., 2022; Chitsaz et al., 2023). There is a complicated process by which carrageenan causes an inflammatory response. It does this by releasing several acute inflammatory mediators and making blood vessels more permeable (Annamalai and Thangam, 2017). Previous studies have documented the anti-inflammatory properties of various mangrove species. The genus *Rhizophora* exhibits a range of pharmacological effects, such as antifungal, antibacterial, antiseptic, anti-inflammatory, and wound-healing activities (Sormin et al., 2021; Dat et al., 2022).

#### 3.4 Analysis of Variance (ANOVA) and Honestly Significant Difference (HSD)

The results of the ANOVA test of the treatment group of anti-inflammatory activity on the plethysmometer with the values of F count and F table, namely  $5.09 > 2.71$  and on the term thrust  $7.21 > 2.71$ , declare the presence of a notable distinction and proceed with the Tukey Honestly Significant Difference (HSD) examination, as outlined in Table 2. The results of the Tukey HSD test state that each treatment group has a significant difference, which is indicated by different numbers in each treatment at each hour.

Based on Table 2, in the plethysmometer measurement method for the Na CMC treatment group, the letter notation (kl) is shown for 1 hour, (ij) 2 hours, (ef) 3 hours, (d) 4 hours, and (c) 5 hours, which means that each treatment experienced a significantly different decrease in edema volume. Furthermore, Na Diclofenac is shown with the letter notation (jk) for 1 hour, (gh) 2 hours, (fg) 3 hours, (c) 4 hours, and (b) 5 hours, which is interpreted as each treatment experiencing a significantly different decrease in edema volume. Then the test group received a 100 mg/kg BW dose of *Sonneratia caseolaris* leaf extract is indicated by the notation (kl) for 1 hour, (ij) 2 hours, (fg) 3 hours, (d) 4 hours, and (b) 5 hours, then the 150 mg/kg BW dose test group is indicated by the notation (m), which is the notation with the highest letter, which is interpreted as

**Table 1.** Percentage Shrinkage of Leaf Extract

Leaf Sample	Sample Weight (gr)		Percentage Depreciation (%)	Percentage Sample Weight (%)
	Wet	Dry		
<i>S. caseolaris</i>	1780	551	69	30,9

**Table 2.** Tukey HSD Further Tested the Results of *S. caseolaris* Leaves Anti-Inflammation

Methods Measurements	Group Treatment	Time (Hour)				
		1	2	3	4	5
Plethysmometer	Na CMC	0.073±0.006kl	0.067±0.006ij	0.053±0.006ef	0.047±0.006d	0.043±0.006c
	Na Diklofenak	0.070±0.10jk	0.060±0.010gh	0.057±0.012fg	0.043±0.006c	0.037±0.006b
	EDM100	0.073±0.006kl	0.067±0.006ij	0.057±0.006fg	0.047±0.006d	0.037±0.006b
	EDM150	0.090±0.010m	0.077±0.012l	0.060±0.000gh	0.043±0.006c	0.027±0.006a
	EDM200	0.073±0.006kl	0.067±0.006ij	0.063±0.006hi	0.050±0.000d	0.033±0.006b
Vernier caliper	Na CMC	3.57±0.25bc	3.12±0.09bc	3.00±0.13bc	2.86±0.09bc	2.80±0.10b
	Na Diklofenak	4.09±0.32c	3.69±0.31c	3.42±0.40bc	3.16±0.29bc	2.96±0.25ab
	EDM100	3.50±0.37c	3.00±0.16b	2.72±0.30ab	2.61±0.23ab	2.46±0.15ab
	EDM150	3.71±0.04bc	3.13±0.03ab	3.04±0.05ab	2.67±0.13ab	2.48±0.12a
	EDM200	3.58±0.24bc	3.25±0.48bc	3.01±0.47ab	2.61±0.21ab	2.47±0.03ab

**Table 3.** Qualitative Phytochemicals

Leaf Sample	Compound				
	Alkaloids	Steroids	Flavonoids	Saponins	Phenols
<i>S. caseolaris</i>	+	+	+	-	+
	+	+	+	-	+

the greatest inflammation of the edema at 1 hour, (l) 2 hours, (gh) 3 hours, (c) 4 hours, and (a) 5 hours, which is interpreted as each treatment experiencing a very significantly different edema volume decrease. Then the dose of 200 mg/kgBB is indicated by the letter notation (kl) for 1 hour, (ij) for 2 hours, (hi) for 3 hours, (d) for 4 hours, and (b) for 5 hours, which means that each treatment has a significantly different edema volume.

Furthermore, in the method of measuring the term thrust for the Na CMC treatment group, the same letter notation (bc) is shown from 1 hour to 4 hours, which means that the treatment is not significantly different, and at the 5 hour, there is a decrease in edema volume with the notation (b), which is interpreted as significantly different. Next, with the Na Diclofenac treatment group, the letter notation (c) is shown for 1 hour to 2 hours, indicating that the treatment is not significantly different, then there is a decrease in edema volume with a change in notation to (bc) at 3 hours to 4 hours, and again a decrease at 5 hours with the notation (ab). In the *S. caseolaris* leaf extract test group for a dose of 100 mg/kgBB indicated by notation (c) at 1 hour and changed at 2 hours with notation (b), which showed a real difference. Furthermore, the decrease in edema volume from 3 hours to 5 hours with notation (c) means that the treatment is not significantly different. Then, at a dose of

150 mg/kgBB indicated by the notation (bc) at 1 hour, there is a real difference with the notation (c), which means that the treatment is not significantly different. The decrease in edema volume from 2 hours with notation (ab) until 3 hours means that the treatment is not significantly different, and only at 5 hours is there a decrease in edema volume with notation (a), which is interpreted as a significantly different treatment. Furthermore, the dose of 200 mg/kg BW is indicated by notation (bc) at 1 hour to 2 hours, which is interpreted as not significantly different, and there is a decrease in edema volume at 3 hours to 5 hours with notation (ab).

The outcomes from the Tukey examination revealed that across the board category of reduction in the volume of edema on the soles of the feet of rats, there were differences in each treatment group. The results of the decrease in edema volume with different letter notation signs in the Tukey HSD test showed that in the 150 mg/kg BW dose of mangrove leaves extract in the digital vernier caliper and plethysmometer method. The use of a digital plethysmometer will make the results or value of the decrease in edema more thorough. According to [Bucan et al. \(2022\)](#), measurements with digital calipers and plethysmometers are reliable and can be applied interchangeably. A plethysmometer is a precise and easy measurement method for decreasing edema volume ([Zhang et al., 2020](#)).

**Table 4.** Retention Time, Peak Area, Name of Compounds, Formula, and Compound Group

Ret. Time	Peak Area %	Name of Compounds	Formula	Compound Group
5.26	3.71	Nonanal	C <sub>9</sub> H <sub>18</sub> O	Aldehydes (Hao, 2019)
12.13	9.16	Desulphosinigrin	C <sub>10</sub> H <sub>17</sub> NO <sub>6</sub> S	Glucosinolates (Youssef et al., 2023)
14.71	2.84	5,8-Epoxy-3H-2-benzopyran, 4,4a,5,8-tetrahydro-5,8-dimethyl-, (4α,5α,8α)-	C <sub>9</sub> H <sub>6</sub> O <sub>2</sub>	Coumarin (Bansal et al., 2013)
17.57	3.40	Hexadecanoic acid, methyl ester	C <sub>17</sub> H <sub>34</sub> O <sub>2</sub>	Ester (Habtemariam, 2019)
20.48	4.61	1-Hexadecanol, 2-methyl-6,9,12,15-	C <sub>17</sub> H <sub>36</sub> O	Alcohol (Prasansang et al., 2023)
20.80	3.88	Docosatetraenoic acid, methyl ester	C <sub>23</sub> H <sub>38</sub> O <sub>2</sub>	Ester (Yu et al., 2020)
21.02	17.26	Phytol	C <sub>20</sub> H <sub>40</sub> O	Terpenoids (Hao et al., 2015)
21.97	3.70	1-Heptatriacotanol	C <sub>37</sub> H <sub>76</sub> O	Alcohol (Zaidan et al., 2019)
22.18	3.07	7-Methyl-Z-tetradecen-1-ol acetate	C <sub>17</sub> H <sub>32</sub> O <sub>2</sub>	Ester (Oluwasina et al., 2023)
27.96	2.75	Hexadecanoic acid, 2-hydroxy-1-(hydroxymethyl)ethyl ester	C <sub>19</sub> H <sub>38</sub> O <sub>4</sub>	Ester (Syed et al., 2022)
31.34	9.64	1-Monolinoleoylglycerol trimethylsilyl ether	C <sub>18</sub> H <sub>32</sub> O <sub>2</sub>	Lipids (Taha et al., 2023)
33.96	6.72	α-Tocopheryl acetate	C <sub>31</sub> H <sub>52</sub> O <sub>3</sub>	Tocopherol (Zahid et al., 2019)
36.24	17.45	β-Sitosterol	C <sub>29</sub> H <sub>50</sub> O	Steroids (Petrova et al., 2023)

### 3.5 Qualitative Phytochemical Analysis of Leaves Extract

Qualitative phytochemical analysis of leaves extract were positive for alkaloids, steroids, flavonoids, and phenols (Table 3). Bioactive compounds contained in *Sonneratia caseolaris* leaves include alkaloids, steroids, and flavonoids. The qualitative phytochemical tests for *S. caseolaris* indicate the presence of bioactive compounds like flavonoids and phenols, which are believed to exhibit potent biological activity, such as anti-inflammatory effects. While saponins had negative results with no shown or detected foam at the time of testing.

Qualitative phytochemical tests on *S. caseolaris* leaves extracts were positive for bioactive compounds that lead to anti-inflammatory activity. In line with the results of the study Kalasuba et al. (2023), the mangrove species *Rhizophora stylosa* is rich in alkaloids, flavonoids, phenolic acids, tannins, terpenoids, saponins, and steroids and is widely used in traditional medicine for its anti-inflammatory, antibacterial, antioxidant, and antipyretic effects. Flavonoids and saponins are recognized for their anti-inflammatory properties (Meshram et al., 2016; Chang et al., 2023). According to Syahidah and Subekti (2019), various metabolite compounds like alkaloids, flavonoids, phenolics, tannins, and terpenoid saponins are present in mangrove plants. Consequently, this plant is extensively utilized in traditional medicine across diverse regions of Indonesia, notably for













its anti-inflammatory and antipyretic properties (Arbiastutie et al., 2021; Al Kazman et al., 2022).

### 3.6 GC-MS Analysis of *S. caseolaris* Leaves Extract

The graph of GC analysis results obtained 13 peak points (Figure 4). Compounds identified based on chromatogram peak heights were matched identically to the library data base: WILEY 7 (Table 4). The chromatogram analysis of *Sonneratia caseolaris* leaf extract fractions clearly showed the presence of 13 secondary metabolite compounds, which were found in the library database (WILEY 7). The main groups of compounds that were found were aldehydes, glucosinolates, coumarins, esters, terpenoids, alcohols, lipids, tocopherols, and steroids. Information related to compound names, chemical structures, and mass spectra is summarized in Table 5.

According to the findings from Table 4, the majority of ethanol extracts from *S. caseolaris* leaves consist of aldehyde compound, glucosinolates, coumarins, esters, terpenoids, alcohols, lipids, tocopherols, and steroids. Nonanal, which is classified as an aldehyde compound, is a compound that is often found in various essential oils and can give a distinctive aroma to some plants. In addition, based on reports Tseng et al. (2023), aldehyde compounds are potent anti-inflammatory active compounds in *Cinnamomum cassia*. Desulphosinigrin, a type of glucosinolate, is also documented for its anti-cancer and antiimi-

**Table 5.** Compound name, Chemical Structure, and Mass Spectral Information

Compound Name, Chemical Structure, and Mass Spectra	
  <p>Nonanal</p>  	  <p>Desulphosinigrin</p>  
<p>5,8-Epoxy-3H-2-benzopyran, 4,4a,5,8-tetrahydro-5,8-dimethyl-, (4<math>\alpha</math>,5<math>\alpha</math>,8<math>\alpha</math>)-</p>   <p>1-Hexadecanol, 2-methyl-</p>	<p>Hexadecanoic acid, methyl ester</p>   <p>6,9,12,15-Docosatetraenoic acid, methyl ester</p>















icrobial properties (Youssef et al., 2023). 4-methylsulfinylbutyl glucosinolate is a compound originating from the amino acid methionine, known for its antifungal, antioxidant, and antimicrobial properties (Elaiyaraja and Chandramohan, 2018).

Hexadecanoic acid, methyl ester, 6,9,12,15-Docosatetraenoic acid, methyl ester, 7-Methyl-Z-tetradecen-1-ol acetate, and Hexadecanoic acid, 2-hydroxy-1-(hydroxymethyl)ethyl ester, are classified as ester class compounds. Research results of Sokeng et al. (2020), it is evident that compounds within the ester group exhibit both acute and chronic anti-inflammatory

effects as well as central analgesic properties. 1-Hexadecanol, 2-methyl- and 1-Heptatriacotanol belong to alcohol compounds. So far, some alcohol compounds, especially those found in plant essential oils, can show anti-inflammatory activity that can help reduce inflammatory reactions in the body (Ali et al., 2015; Agarwal et al., 2022). Phytol is classified among terpenoid compounds, which represent one of the most diverse categories of natural substances, garnering significant interest because of their wide range of biological activities (Masyita et al., 2022). Terpenoids exhibit potent anti-inflammatory effects, making



Table 5. Cont.

Compound Name, Chemical Structure, and Mass Spectra	
  <b>Phytol</b>	  <b>1-Heptatriacotanol</b>
  <b>7-Methyl-Z-tetradecen-1-ol acetate</b>	  <b>Hexadecanoic acid, 2-hydroxy-1-(hydroxymethyl)ethyl ester</b>
  <b>1-Monolinoleoylglycerol trimethylsilyl ether</b>	  <b>α-Tocopheryl acetate</b>
  <b>β-Sitosterol</b>	

Source : (Kim et al., 2022).

them viable candidates for subsequent structural modifications and optimization (Ge et al., 2022; Ghallab et al., 2024). Another discovered component, 1-Monolinoleoylglycerol trimethylsilyl ether, has been reported as an antioxidant, antimicro-

bial, and anti-inflammatory agent (Mohan et al., 2015; Sezer and Uysal, 2021). α-Tocopheryl acetate is classified as a tocopherol compound. Based on the results of research Cassano (2012), each form of vitamin E (α-tocopherol) has biologi-

cal activity, with its natural configuration being 2R,4'R,8'R whose main function of vitamin E is as an antioxidant. The last compound is  $\beta$ -Sitosterol classified as a steroid compound. According to the results of research Chang et al. (2023),  $\beta$ -Sitosterol (BS) is present in a variety of plant seeds, vegetable oils, and nuts demonstrating diverse anti-inflammatory effects on peritoneal macrophages and peripheral tissues.

Furthermore, 5,8-Epoxy-3H-2-benzopyran, 4,4a,5,8-tetrahydro-5,8-dimethyl-, (4 $\alpha$ ,5 $\alpha$ ,8 $\alpha$ )- belongs to the class of coumarin compounds. Several naturally occurring compounds containing coumarinic groups have been documented to possess diverse biological activities (Menezes and Diederich, 2019; Sun et al., 2021). According to Annunziata et al. (2020), coumarin in the pharmaceutical field is known for its potential as an anticoagulant, which means it can affect blood clotting. Similar to isomeric flavonoids, coumarin is anticipated to impact the generation and elimination of reactive oxygen species (ROS) and affect processes associated with free radical-induced damage. Coumarin has the potential to alleviate tissue swelling and inflammation. Also, coumarins and their 7-hydroxy derivatives stop the production of prostaglandins, which encompasses intermediates of fatty acid hydroperoxides (Fylaktakidou et al., 2004). Several studies show that coumarin compounds have the ability to inhibit the enzymes cyclooxygenase (COX) and lipoxygenase (LOX), which are the main enzymes in the prostaglandin and leukotriene synthesis pathways (Lončarić et al., 2020; Chahal et al., 2023; Rudrapal et al., 2023). Through inhibiting the function of this enzyme, coumarin compounds can decrease the generation of inflammatory mediators, thus mitigating the inflammatory response (Al-Duhaidahawi et al., 2022; Chen et al., 2024). Coumarin compounds can also influence the activity of immune cells, which play a role in the inflammatory process (Sandhiutami et al., 2017; Rostom et al., 2022). This can be accomplished by diminishing the release of pro-inflammatory cytokines and hindering the migration of inflammatory cells to the inflammation site, or increasing the activity of immune cells responsible for the recovery and resolution of inflammation (Wu et al., 2022; Zhang et al., 2023; Chen et al., 2024).

Therefore, the ethanol extract derived from *Sonneratia caseolaris* leaves contains a diverse array of compounds that have been documented to possess anti-inflammatory, anti-cancer, antioxidant, and antimicrobial properties. Based on the multiple mechanisms of action involved, coumarins are promising compounds for reducing inflammation and may have potential for use in the treatment of inflammatory diseases. These findings open up great opportunities for the utilization of *S. caseolaris* leaves as a potential source for future traditional medicines, which can help people maintain their health.

#### 4. CONCLUSIONS

*S. caseolaris* leaves extract has anti-inflammatory activity against carrageenan-induced test animals. The anti-inflammatory effect is denoted by the lowest percentage of inflammation observed during the 5-hour monitoring period, which is calcu-

lated as the average decrease. The largest dose of *S. caseolaris* leaves extract is found in the 150 mg/kg BW mangrove leaf extract group dose using a digital caliper and plethysmometer. *S. caseolaris* mangrove leaves positively contain secondary metabolites, namely flavonoids, steroids, alkaloids, and phenols. The findings from the GC-MS examination of *S. caseolaris* leaves extracts are thought to contain groups of aldehyde compounds, glucosinolates, coumarins, esters, terpenoids, alcohols, lipids, tocopherols, and steroids.

#### 5. ACKNOWLEDGMENT

The research of this article was funded by DIPA of Public Service Agency of Universitas Sriwijaya 2023. Nomor SP DIPA-023.17.2.677515/2023 on the 30th of November 2022. In accordance with the Rector's Decree Number: 0188/UN9.3.1/SK/2023 on the 18th of April 2023.

#### REFERENCES

- Acquaviva, A., S. C. Di Simone, Nilofar, A. Bouyahya, G. Zengin, L. Recinella, S. Leone, L. Brunetti, A. I. Uba, O. Guler, M. Balos, U. Cakilcioglu, L. Menghini, C. Ferrante, G. Orlando, M. L. Libero, and A. Chiavaroli (2023). Screening for Chemical Characterization and Pharmacological Properties of Different Extracts from *Nepeta italica*. *Plants*, **12**(15)
- Agarwal, P., Z. Sebghatollahi, M. Kamal, A. Dhyani, A. Shrivastava, K. K. Singh, M. Sinha, N. Mahato, A. K. Mishra, and K. H. Baek (2022). Citrus Essential Oils in Aromatherapy: Therapeutic Effects and Mechanisms. *Antioxidants*, **11**(12)
- Al-Duhaidahawi, D., H. F. S. AL-Zubaidy, K. Al-Khafaji, and A. AL-Ameiry (2022). Synthesis, Anti-Inflammatory Effects, Molecular Docking and Molecular Dynamics Studies of 4-Hydroxy Coumarin Derivatives As Inhibitors of COX-II Enzyme. *Journal of Molecular Structure*, **1247**; 131377
- Al Kazman, B. S. M., J. E. Harnett, and J. R. Hanrahan (2022). Traditional Uses, Phytochemistry and Pharmacological Activities of *Annonaceae*. *Molecules*, **27**(11)
- Ali, B., N. A. Al-Wabel, S. Shams, A. Ahamad, S. A. Khan, and F. Anwar (2015). Essential Oils Used in Aromatherapy: A Systemic Review. *Asian Pacific Journal of Tropical Biomedicine*, **5**(8); 601–611
- Altemimi, A., N. Lakhssassi, A. Baharlouei, D. G. Watson, and D. A. Lightfoot (2017). Phytochemicals: Extraction, Isolation, and Identification of Bioactive Compounds from Plant Extracts. *Plants*, **6**(4)
- Annamalai, P. and E. B. Thangam (2017). Local and Systemic Profiles of Inflammatory Cytokines in Carrageenan-induced Paw Inflammation in Rats. *Immunological Investigations*, **46**(3); 274–283
- Annunziata, F., C. Pinna, S. Dallavalle, L. Tamborini, and A. Pinto (2020). An Overview of Coumarin as a Versatile and Readily Accessible Scaffold with Broad-Ranging Biological Activities. *International Journal of Molecular Sciences*, **21**(13); 1–83
- Anu, K., H. P. K, V. K. Sneha, P. Busheera, J. Muhammed,

- and A. Augustine (2024). Results in Engineering Mangroves in Environmental Engineering: Harnessing the Multifunctional Potential of Nature's Coastal Architects for Sustainable Ecosystem Management. *Results in Engineering*, **21**(January); 101765
- Arbiastutie, Y., F. Diba, and Masriani (2021). Short Communication: Ethnobotanical and Ecological Studies of Medicinal Plants in a Mangrove Forest in Mempawah District, West Kalimantan, Indonesia. *Biodiversitas Journal of Biological Diversity*, **22**(6); 3164–3170
- Bai, R., C. Yao, Z. Zhong, J. Ge, Z. Bai, X. Ye, T. Xie, and Y. Xie (2021). Discovery of Natural Anti-inflammatory Alkaloids: Potential Leads for the Drug Discovery for the Treatment of Inflammation. *European Journal of Medicinal Chemistry*, **213**; 113165
- Bansal, Y., P. Sethi, and G. Bansal (2013). Coumarin: A Potential Nucleus for Anti-inflammatory Molecules. *Medicinal Chemistry Research*, **22**(7); 3049–3060
- Batubara, R., Surjanto, T. Ismanelly Hanum, A. Handika, and O. Affandi (2020). The Screening of Phytochemical and Antioxidant Activity of Agarwood Leaves (*Aquilaria malaccensis*) from Two Sites in North Sumatra, Indonesia. *Biodiversitas*, **21**(4); 1588–1596
- Biswas, P. L. and S. R. Biswas (2021). Mangrove Forests: Ecology, Management, and Threats. In *Encyclopedia of the UN Sustainable Development Goals*. Springer, Cham, pages 627–640
- Bouabida, H. and D. Dris (2022). Phytochemical Constituents and Larvicidal Activity of *Ruta graveolens*, *Ruta montana* and *Artemisia absinthium* Hydro-methanolic Extract against Mosquito Vectors of Avian Plasmodium (*Culiseta longiareolata*). *South African Journal of Botany*, **151**; 504–511
- Bouyahya, A., F. E. Guaouguaou, N. El Omari, N. El Menyiy, A. Balahbib, M. El-Shazly, and Y. Bakri (2022). Anti-inflammatory and Analgesic Properties of Moroccan Medicinal Plants: Phytochemistry, In Vitro and In Vivo Investigations, Mechanism Insights, Clinical Evidences and Perspectives. *Journal of Pharmaceutical Analysis*, **12**(1); 35–57
- Brendel, O. (2021). The Relationship between Plant Growth and Water Consumption: A History from the Classical Four Elements to Modern Stable Isotopes. *Annals of Forest Science*, **78**(2); 1–16
- Bucan, A., A. Wiinholt, F. Dalaei, O. Gerke, C. R. Hansen, and J. A. Sørensen (2022). Microcomputed Tomography versus Plethysmometer and Electronic Caliper in the Measurements of Lymphedema in the Hindlimb of Mice. *Scientific Reports*, **12**(1); 12267
- Cassano, R. (2012). Vitamin E Chemistry, Biological Activity and Benefits on the Skin. In *Handbook of Diet, Nutrition and the Skin*. Wageningen Academic, pages 144–163
- Chahal, S., P. Rani, Kiran, J. Sindhu, G. Joshi, A. Ganesan, S. Kalyanamoorthy, Mayank, P. Kumar, R. Singh, and A. Negi (2023). Design and Development of COX-II Inhibitors: Current Scenario and Future Perspective. *ACS Omega*, **8**(20); 17446–17498
- Chang, Y., S. Wang, J. Xu, T. Zhu, S. Ma, A. Zhou, Y. Song, M. Liu, and C. Tian (2023). Optimization of Extraction Process of *Dioscorea nipponica* Makino Saponins and their UPLC-QTOF-MS Profiling, Antioxidant, Antibacterial and Anti-inflammatory Activities. *Arabian Journal of Chemistry*, **16**(4); 104630
- Chen, I. C., W. L. Chen, K. H. Chang, J. W. Lee, T. H. Lin, W. Lin, C. M. Chen, and G. J. Lee-Chen (2024). Investigating the Therapeutic Effects of Novel Compounds Targeting Inflammatory IL-1 $\beta$  and IL-6 Signaling Pathways in Spinocerebellar Ataxia Type 3. *European Journal of Pharmacology*, **967**; 176370
- Chen, T., R. Zhou, X. J. Ge, and S. Shi (2008). Development and Characterization of Microsatellite Markers for a Mangrove Tree Species *Sonneratia caseolaris* (L.) Engler (Lythraceae sensu lato). *Conservation Genetics*, **9**(4); 957–959
- Chitsaz, R., A. Zarezadeh, J. Asgarpanah, P. Najafzadeh, and Z. Mousavi (2023). Rubiadin Exerts an Acute and Chronic Anti-Inflammatory Effect in Rodents. *Brazilian Journal of Biology*, **83**; 1–8
- Das, S., A. Chakravorty, S. Luktuke, A. Raj, A. A. Mini, K. Ramesh, A. N. Grace, S. K. Pandey, and V. Raghavan (2023). Graphene/Gadolinium Oxide Composite Modified Screen-Printed Electrochemical Sensor for Detection of Diclofenac Sodium. *Results in Chemistry*, **6**; 101189
- Dat, T. T. H., L. C. V. Cuong, D. V. Ha, P. T. T. Oanh, N. P. K. Nhi, H. L. T. Anh, P. T. Quy, T. Q. Bui, N. T. Triet, and N. T. A. Nhung (2022). The Study on Biological Activity and Molecular Docking of Secondary Metabolites from *Bacillus* sp. Isolated from the Mangrove Plant *Rhizophora apiculata* Blume. *Regional Studies in Marine Science*, **55**; 102583
- de Araújo, I. W. F., J. A. G. Rodrigues, A. L. G. Quinderé, J. d. F. T. Silva, G. d. F. Maciel, N. A. Ribeiro, E. de Sousa Oliveira Vanderlei, K. A. Ribeiro, H. V. Chaves, K. M. A. Pereira, M. M. Bezerra, and N. M. B. Benevides (2016). Analgesic and Anti-Inflammatory Actions on Bradykinin Route of a Polysulfated Fraction from Alga *Ulva lactuca*. *International Journal of Biological Macromolecules*, **92**; 820–830
- Elaiyaraja, A. and G. Chandramohan (2018). Comparative Phytochemical Profile of *Crinum defixum* Ker-Gawler Leaves Using GC-MS. *Journal of Drug Delivery and Therapeutics*, **8**(4); 365–380
- Elfaruk, M. S., C. Wen, C. Chi, X. Li, and S. Janaswamy (2021). Effect of Salt Addition on Iota-Carrageenan Solution Properties. *Food Hydrocoll*, **113**; 106491
- Farida, S., D. K. Pratami, M. Sahlan, D. R. Laksmiawati, E. Rohmatin, and H. Situmorang (2022). In-Vitro Antioxidant, In-Vivo Anti-Inflammatory, and Acute Toxicity Study of Indonesian Propolis Capsule from *Tetragonula sapiens*. *Saudi Journal of Biological Sciences*, **29**(4); 2489–2500
- Fehrenbacher, J. C., M. R. Vasko, and D. B. Duarte (2012). Models of Inflammation: Carrageenan- or Complete Freund's Adjuvant-Induced Edema and Hypersensitivity in the Rat. *Current Protocols in Pharmacology*, **0** 5(SUPPL.56); Unit5.4



- Fitria, Y., M. Fitriani, R. Y. Nugroho, and W. A. E. Putri (2023). Gastropods As Bioindicators of Heavy Metal Pollution in the Banyuasin Estuary Shrimp Pond Area, South Sumatra, Indonesia. *Acta Ecologica Sinica*, **43**(6); 1129–1137
- Fylaktakidou, K., D. Hadjipavlou-Litina, K. Litinas, and D. Nicolaidis (2004). Natural and Synthetic Coumarin Derivatives with Anti-Inflammatory/Antioxidant Activities. *Current Pharmaceutical Design*, **10**(30); 3813–3833
- García-Suárez, P. C., I. Rentería, E. P. Plaisance, J. Moncada-Jiménez, and A. Jiménez-Maldonado (2021). The Effects of Interval Training on Peripheral Brain Derived Neurotrophic Factor (BDNF) in Young Adults: A Systematic Review and Meta-Analysis. *Scientific Reports*, **11**(1); 8937
- Ge, J., Z. Liu, Z. Zhong, L. Wang, X. Zhuo, J. Li, X. Jiang, X. Y. Ye, T. Xie, and R. Bai (2022). Natural Terpenoids with Anti-Inflammatory Activities: Potential Leads for Anti-Inflammatory Drug Discovery. *Bioorganic Chemistry*, **124**; 105817
- Ghallas, D. S., R. S. Ibrahim, M. M. Mohyeldin, and E. Shawky (2024). Marine Algae: A Treasure Trove of Bioactive Anti-Inflammatory Compounds. *Marine Pollution Bulletin*, **199**; 116023
- Giesen, W., S. Wulffraat, M. Zieren, and L. Scholten (2007). *Mangrove Guidebook for Southeast Asia*. FAO Regional Office for Asia and the Pacific
- Habtemariam, S. (2019). The Chemical and Pharmacological Basis of Fenugreek (*Trigonella foenum-graecum* L.) as Potential Therapy for Type 2 Diabetes and Associated Diseases. *Medicinal Foods as Potential Therapies for Type-2 Diabetes and Associated Diseases*, **1**; 579–637
- Hao, D. (2019). Mining Chemodiversity From Biodiversity: Pharmacophylogeny of Ranunculaceae Medicinal Plants. In D.-C. Hao, editor, *Ranunculales Medicinal Plants*. Academic Press, pages 35–71
- Hao, D. C., X. Gu, and P. G. Xiao (2015). Phytochemical and Biological Research of Cannabis Pharmaceutical Resources. In D. C. Hao, X.-J. Gu, and P. G. Xiao, editors, *Medicinal Plants*. Woodhead Publishing, pages 431–464
- Harborne, J. B. (1998). *Phytochemical Methods: A Guide to Modern Techniques of Plant Analysis*. Chapman and Hall
- Hikmawanti, N. P. E., S. Fatmawati, and A. W. Asri (2021). The Effect of Ethanol Concentrations as the Extraction Solvent on Antioxidant Activity of Katuk (*Sauropus androgynus* (L.) Merr.) Leaves Extracts. *IOP Conference Series: Earth and Environmental Science*, **755**(1)
- Hossain, S. J., M. R. Islam, T. Pervin, M. Iftekharuzzaman, O. A. Hamdi, S. Mubassara, M. Saifuzzaman, and J. A. Shilpi (2017). Antibacterial, Anti-Diarrhoeal, Analgesic, Cytotoxic Activities, and GC-MS Profiling of *Sonneratia apetala* (buch.-Ham.) Seed. *Preventive Nutrition and Food Science*, **22**(3); 157
- Itoh, H., N. Yamashita, S. Kamijo, K. Masuda, H. Kato, and S. Yamaori (2023). Effects of Acidic Non-Steroidal Anti-Inflammatory Drugs on Human Cytochrome P450 4A11 Activity: Roles of Carboxylic Acid and a Sulfur Atom in Potent Inhibition by Sulindac Sulfide. *Chemical Biology Interactions*, **382**; 110644
- Ivorra, L., P. G. Cardoso, S. K. Chan, C. Cruzeiro, and K. A. Tagulao (2021). Can Mangroves Work as an Effective Phytoremediation Tool for Pesticide Contamination? An Interlinked Analysis Between Surface Water, Sediments and Biota. *Journal of Cleaner Production*, **295**; 126334
- Kalasuba, K., M. Miranti, S. R. Rahayuningsih, W. Safriansyah, R. R. P. Syamsuri, K. Farabi, D. Oktavia, A. N. Alhasnawi, and F. Doni (2023). Red Mangrove (*Rhizophora stylosa* Griff.)—A Review of Its Botany, Phytochemistry, Pharmacological Activities, and Prospects. *Plants*, **12**(11)
- Kim, S., T. Cheng, S. He, P. A. Thiessen, Q. Li, A. Gindulyte, and E. E. Bolton (2022). PubChem Protein, Gene, Pathway, and Taxonomy Data Collections: Bridging Biology and Chemistry Through Target-Centric Views of PubChem Data. *Journal of Molecular Biology*, **434**(11); 167514
- Komakech, R., Y. Kim, G. M. Matsabisa, and Y. Kang (2019). Anti-Inflammatory and Analgesic Potential of *Tamarindus indica* Linn. (Fabaceae): A Narrative Review. *Integrative Medicine Research*, **8**(3); 181–186
- Kundu, P., S. L. Debnath, M. F. Ahad, H. S. Devnath, L. Saha, U. K. Karmakar, and S. K. Sadhu (2023). Exploration of In Vivo and In Vitro Biological Effects of *Sonneratia caseolaris* (L.) Fruits Supported by Molecular Docking and ADMET Study. *BioMed Research International*, **2023**; 1–21
- Kundu, P., S. L. Debnath, H. S. Devnath, L. Saha, and S. K. Sadhu (2022). Analgesic, Anti-Inflammatory, Antipyretic, and In Silico Measurements of *Sonneratia caseolaris* (L.) Fruits from Sundarbans, Bangladesh. *BioMed Research International*, **2022**; 1–16
- Liang, C., S. Meng, Y. Wang, X. Xie, Z. Zhang, and D. Cheng (2023). Preparation and Activity of Sodium Carboxymethyl Cellulose (CMC-Na) and *Metarhizium rileyi* ZHKUMR1 Composite Membrane. *International Journal of Biological Macromolecules*, **253**; 126858
- Liu, B., X. Wang, Y. Wang, X. Chen, X. Jin, and X. Luo (2023). Review of Compounds and Activities from Mangrove *Sonneratia* Genus and Their Endophytes. *Journal of Holistic Integrative Pharmacy*, **4**(3); 218–227
- Lončarić, M., I. Strelec, V. Pavić, D. Šubarić, V. Rastija, and M. Molnar (2020). Lipoxigenase inhibition activity of coumarin derivatives—qsar and molecular docking study. *Pharmaceuticals*, **13**(7); 154
- MacLeod, A. S. and J. N. Mansbridge (2016). The Innate Immune System in Acute and Chronic Wounds. *Advances in Wound Care*, **5**(2); 65
- Maehle, A. H. (2009). A Binding Question: The Evolution of the Receptor Concept. *Endeavour*, **33**(4); 135
- Maldonado Miranda, J. J. (2021). Medicinal plants and their traditional uses in different locations. *Phytomedicine A Treasure Pharmacol. Act. Prod. from Plants*; 207–223
- Masyita, A., R. Mustika Sari, A. Dwi Astuti, B. Yasir, N. Rahma Rumata, T. B. Emran, F. Nainu, and J. Simal-Gandara (2022). Terpenes and Terpenoids As Main Bioac-

- tive Compounds of Essential Oils, Their Roles in Human Health and Potential Application As Natural Food Preservatives. *Food Chemistry*; X, **13**; 100217
- Menezes, J. C. J. M. D. S. and M. F. Diederich (2019). Natural Dimers of Coumarin, Chalcones, and Resveratrol and the Link between Structure and Pharmacology. *European Journal of Medicinal Chemistry*, **182**; 111637
- Meshram, G. G., A. Kumar, W. Rizvi, C. D. Tripathi, and R. A. Khan (2016). Evaluation of the Anti-Inflammatory Activity of the Aqueous and Ethanolic Extracts of the Leaves of *Albizia lebeck* in Rats. *Journal of Traditional and Complementary Medicine*, **6**(2); 172–175
- Mohan, V., P. B. Pg, and S. Mgt (2015). GC-MS Analysis of Phytocomponents in *Pleiospermium alatum* (Wall. ex Wight & Arn.) Swingle, (Rutaceae). *Journal of Pharmacognosy and Phytochemistry*, **4**(1); 216–222
- Nascimento, R. d. P. d., A. M. T. M. Moya, A. P. d. F. Machado, M. V. Geraldi, P. Diez-Echave, T. Vezza, J. Galvez, C. B. B. Cazarin, and M. R. Maróstica Junior (2021). Review on the Potential Application of Non-Phenolic Compounds from Native Latin American Food Byproducts in Inflammatory Bowel Diseases. *Food Research International*, **139**; 109796
- Nasrin, S., H. Mahmood, and M. M. Rahman (2021). Parental Influences on Salt Adaptive Divergence of *Sonneratia apetala* (Buch.-Ham.) Seedlings along the Salinity Gradient in the Sundarbans of Bangladesh. *Environmental Challenges*, **4**; 100121
- Olasunkanmi, A. A., O. S. Fadahunsi, and P. I. Adegbola (2022). Gas Chromatography-Mass Spectroscopic, High Performance Liquid Chromatographic and In-Silico Characterization of Antimicrobial and Antioxidant Constituents of *Rhus longipes* (engl). *Arabian Journal of Chemistry*, **15**(2); 103601
- Oluwasina, O. O., S. O. Idris, C. O. Ogidi, and F. O. Igbe (2023). Production of Herbal Toothpaste: Physical, Organoleptic, Phyto-Compound, and Antimicrobial Properties. *Heliyon*, **9**(3); e13892
- Parthiban, A., R. Sivasankar, V. Sachithanandam, S. A. Khan, A. Jayshree, K. Murugan, and R. Sridhar (2023). An Integrative Review on Bioactive Compounds from Indian Mangroves for Future Drug Discovery. *South African Journal of Botany*, **149**(September); 899–915
- Petrova, A. V., E. F. Khusnutdinova, A. N. Lobov, L. M. Zakirova, N. T. T. Ha, and D. A. Babkov (2023). Selective synthesis of A-ring *E*-arylidene derivatives from  $\beta$ -sitosterol and their activity. *Natural Product Research*, **38**(1); 52–59
- Prasaneang, W., K. Choojun, Y. Poo-arporn, A. L. Huang, Y. C. Lin, E. Chen, H. H. Lee, P. W. Chung, and T. Sooknoi (2023). Tuning Cu<sup>+</sup> species/Brønsted acids of copper phyllosilicate by K<sup>+</sup> doping for selective hydrogenation of methyl palmitate to hexadecanol. *Journal of Catalysis*, **428**; 115115
- Ragavan, P., K. Ravichandran, P. M. Mohan, A. Sxaena, R. S. Prasanth, R. S. C. Jayaraj, and S. Saravanan (2014). Short Communication: New Distributional Records of *Sonneratia* Spp. From Andaman and Nicobar Islands, India. *Biodiversitas Journal of Biological Diversity*, **15**(2); 251–260
- Rahim, A. C. and M. F. Abu Bakar (2018). Pidada—*Sonneratia caseolaris*. In S. Rodrigues, E. de Oliveira Silva, and E. S. de Brito, editors, *Exotic Fruits*. Academic Press, pages 327–332
- Ramses, Ismarti, F. Amelia, Rozirwan, and Suheryanto (2020). Diversity and abundance of polychaetes in the west coast waters of batam island, Kepulauan Riau province-indonesia. *AACL Bioflux*, **13**(1); 381–391
- Ríos, J. L. and I. Andújar (2020). *Apoptotic activities of Mediterranean plants*. Academic Press, second edition edition
- Rostom, B., R. Karaky, I. Kassab, and M. Sylla-Iyarreta Veitía (2022). Coumarins derivatives and inflammation: Review of their effects on the inflammatory signaling pathways. *European Journal of Pharmacology*, **922**; 174867
- Rozirwan, F. Fauziyah, R. Y. Nugroho, M. Melki, T. Z. Ulqodry, F. Agustriani, E. N. Ningsih, W. A. E. Putri, A. Absori, and M. Iqbal (2022). An Ecological Assessment of Crab's Diversity Among Habitats of Migratory Birds at Berbak-Sembilang National Park Indonesia. *International Journal of Conservation Science*, **13**(3); 961–972
- Rozirwan, H. Hananda, R. Y. Nugroho, R. Apri, N. N. Khotimah, F. Fauziyah, W. A. E. Putri, and R. Aryawati (2023a). Antioxidant Activity, Total Phenolic, Phytochemical Content, and HPLC Profile of Selected Mangrove Species from Tanjung Api-Api Port Area, South Sumatra, Indonesia. *Tropical Journal of Natural Product Research*, **7**(7); 3482–3489
- Rozirwan, N. N. Khotimah, W. A. E. Putri, Fauziyah, R. Aryawati, N. Damiri, Isnaini, and Nigr (2024). Ecological Risk Assessment of Heavy Metal (Pb, Cu) Contamination in Water, Sediment, and Polychaeta (*Neoleanira Tetragona*) from Coastal Areas Affected by Aquaculture, Urban Rivers, and Ports in South Sumatra. *Journal of Ecological Engineering*, **25**(1); 303–319
- Rozirwan, H. I. Muda, and T. Z. Ulqodry (2020). Short Communication: Antibacterial Potential of Actinomycetes Isolated from Mangrove Sediment in Tanjung Api-Api, South Sumatra, Indonesia. *Biodiversitas Journal of Biological Diversity*, **21**(12); 5723–5728
- Rozirwan, Nanda, R. Y. Nugroho, G. Diansyah, Muhtadi, Fauziyah, W. A. E. Putri, and A. Agusalm (2023b). Phytochemical Composition, Total Phenolic Content and Antioxidant Activity of *Anadara granosa* (Linnaeus, 1758) Collected from the East Coast of South Sumatra, Indonesia Abstract. *Baghdad Science Journal*; 1–8
- Rozirwan, S. Ramadani, W. Ayu, E. Putri, N. Nur, and R. Y. Nugroho (2023c). Evaluation of Calcium and Phosphorus Content in Scallop Shells (*Placuna placenta*) and Blood Cockle Shells (*Anadara granosa*) from Banyuasin Waters, South Sumatra. *Egyptian Journal of Aquatic Biology and Fisheries*, **27**(3); 1053–1068
- Rozirwan, A. P. Saputri, R. Y. Nugroho, N. N. Khotimah, W. A. E. Putri, Fauziyah, and A. I. S. Purwiyanto (2023d). An Assessment of Pb and Cu in Waters, Sediments, and Mud Crabs (*Scylla serrata*) from Mangrove Ecosystem Near Tanjung Api-Api Port Area, South Sumatra, Indonesia. *Science*

- and *Technology Indonesia*, **8**(4); 675–683
- Rudrapal, M., W. A. Eltayeb, G. Rakshit, A. A. El-Arabey, J. Khan, S. M. Aldosari, B. Alshehri, and M. Abdalla (2023). Dual Synergistic Inhibition of COX and LOX by Potential Chemicals from Indian Daily Spices Investigated Through Detailed Computational Studies. *Scientific Reports*, **13**(1); 1–27
- Sandhiutami, N. M. D., M. Moordiani, D. R. Laksmiawati, N. Fauziah, M. Maesaroh, and W. Widowati (2017). In Vitro Assessment of Anti-inflammatory Activities of Coumarin and Indonesian Cassia Extract in RAW264.7 Murine Macrophage Cell Line. *Iranian Journal of Basic Medical Sciences*, **20**(1); 99
- Sezer, E. N. S. and T. Uysal (2021). Phytochemical Analysis, Antioxidant and Anticancer Potential of *Sideritis niveotomentosa*: Endemic Wild Species of Turkey. *Molecules*, **26**(9); 4420
- Shaukat, T. M., M. O. Omer, A. Javeed, H. u. Rehman, and T. M. Shaukat (2023). Isolation of Alkaloidal and Glycosidal Fractions from Leaves of *Trigonella foenum-graecum* L. cv. Desi Indigenous to Pakistan for Antiprostaglandin Evaluation as Substitute of Nonsteroidal Anti-inflammatory Drugs. *Journal of Ethnopharmacology*, **317**; 116730
- Sokeng, S. D., E. Talla, P. Sakava, M. A. Fokam Tagne, C. Henoumont, L. Sophie, J. T. Mbafor, and F. N. Tchuenguem Fohouo (2020). Anti-Inflammatory and Analgesic Effect of Arachic Acid Ethyl Ester Isolated from Propolis. *Biomedical Research International*
- Sormin, R. B. D., D. M. Nendissa, M. N. Mailoa, F. Rieuwpassa, and M. R. Wenno (2021). Antibacterial Activity of *Rhizophora apiculata* Extract Originated from Inner Ambon Bay against Selected Pathogen Bacteria. *IOP Conference Series: Earth and Environmental Science*, **797**(1); 012017
- Su, J. Y., Q. C. Li, and L. Zhu (2011). Evaluation of the in Vivo Anti-inflammatory Activity of a Flavone Glycoside from *Cancerina discoidea* (Ledeb.) Poljak. *EXCLI Journal*, **10**; 110
- Sun, M., M. Sun, and J. Zhang (2021). Osthole: An Overview of Its Sources, Biological Activities, and Modification Development. *Medicinal Chemistry Research*, **30**(10); 1767–1794
- Syahidah and N. Subekti (2019). Phytochemical Analysis of Mangrove Leaves (*Rhizophora sp.*). *IOP Conference Series: Materials Science and Engineering*, **593**(1); 012007
- Syed, R. U., S. S. Moni, R. H. Alfaisal, R. H. Alrashidi, N. F. Alrashidi, K. M. Wadeed, F. N. Alshammary, A. M. Habib, F. M. Alharbi, Z. ur Rehman, M. Shamsheer Alam, V. K. Basode, and A. A. Abdulhaq (2022). Spectral Characterization of the Bioactive Principles and Antibacterial Properties of Cold Methanolic Extract of *Olea europaea* from the Hail Region of Saudi Arabia. *Arabian Journal of Chemistry*, **15**(8); 104006
- Taha, M., S. T. Elazab, O. Abdelbagi, A. A. Saati, O. Babateen, T. A. S. Baokbah, N. F. Qusty, M. E. Mahmoud, M. M. Ibrahim, and A. M. Badawy (2023). Phytochemical Analysis of *Origanum majorana* L. Extract and Investigation of Its Antioxidant, Anti-inflammatory and Immunomodulatory Effects against Experimentally Induced Colitis Downregulating Th17 Cells. *Journal of Ethnopharmacology*, **317**; 116826
- Tseng, S. H., C. J. Lee, S. H. Chen, C. H. Chen, P. W. Tsai, M. S. Hsieh, J. S. Chu, and C. C. Wang (2023). Cinnamic Aldehyde, an Anti-inflammatory Component in Du-Huo-Ji-Sheng-Tang, Ameliorates Arthritis in II Collagenase and Monosodium Iodoacetate Induced Osteoarthritis Rat Models. *Journal of Traditional and Complementary Medicine*, **13**(1); 51–61
- Tsujimae, M., A. Masuda, M. Takagi, T. Kato, R. Nakano, K. Fujita, Y. Hirata, S. Kakuyama, K. Furumatsu, T. Nakagawa, K. Ogisu, S. Fujigaki, T. Iemoto, T. Ezaki, Y. Yagi, T. Ikegawa, K. Yamanaka, Y. Sato, N. Juri, and Y. Kodama (2024). Relapse and Side Effects of Steroid Therapy Beyond 3 Years in Autoimmune Pancreatitis: A Multicenter Retrospective Study. *Pancreatology*
- Wainwright, C. L., M. M. Teixeira, D. L. Adelson, E. J. Buenz, B. David, K. B. Glaser, Y. Harata-Lee, M. J. R. Howes, A. A. Izzo, P. Maffia, A. M. Mayer, C. Mazars, D. J. Newman, E. Nic Lughadha, A. M. Pimenta, J. A. Parra, Z. Qu, H. Shen, M. Spedding, and J. L. Wolfender (2022). Future Directions for the Discovery of Natural Product-Derived Immunomodulating Drugs: An IUPHAR Positional Review. *Pharmacological Research*, **177**; 106076
- Wu, M., L. J. Hill, L. E. Downie, and H. R. Chinnery (2022). Neuroimmune Crosstalk in the Cornea: The Role of Immune Cells in Corneal Nerve Maintenance During Homeostasis and Inflammation. *Progress in Retinal and Eye Research*, **91**; 101105
- Youssef, A. M. M., D. A. M. Maaty, and Y. M. Al-Saraireh (2023). Phytochemistry and Anticancer Effects of Mangrove (*Rhizophora mucronata Lam.*) Leaves and Stems Extract against Different Cancer Cell Lines. *Pharmaceuticals*, **16**(1); 4
- Yu, J., H. M. Yang, Y. Y. Lai, X. L. Wan, and Z. Y. Wang (2020). The Body Fat Distribution and Fatty Acid Composition of Muscles and Adipose Tissues in Geese. *Poultry Science*, **99**(9); 4634–4641
- Zahid, S., H. Khalid, F. Ikram, H. Iqbal, M. Samie, L. Shahzadi, A. T. Shah, M. Yar, A. A. Chaudhry, S. J. Awan, A. F. Khan, and I. u. Rehman (2019). Bi-layered  $\alpha$ -tocopherol Acetate Loaded Membranes for Potential Wound Healing and Skin Regeneration. *Materials Science and Engineering: C*, **101**; 438–447
- Zaidan, U. H., N. I. Mohamad Zen, N. A. Amran, S. Shamsi, and S. S. A. Gani (2019). Biochemical Evaluation of Phenolic Compounds and Steviol Glycoside from *Stevia rebaudiana* Extracts Associated with In vitro Antidiabetic Potential. *Bio-catalysis and Agricultural Biotechnology*, **18**; 101049
- Zhang, X., X. Han, Y. Jiang, Y. Wang, X. He, D. Liu, J. Huang, H. Liu, T. Ye, and S. Li (2023). Impact of Inflammation and Anti-Inflammatory Modalities on Diabetic Cardiomyopathy Healing: From Fundamental Research to Therapy. *International Immunopharmacology*, **123**; 110747

- Zhang, Z., A. Chinnathambi, S. Ali Alharbi, and L. Bai (2020). Copper Oxide Nanoparticles from *Rabdosia rubescens* Attenuates the Complete Freund's Adjuvant (CFA) Induced Rheumatoid Arthritis in Rats via Suppressing the Inflammatory Proteins COX-2/PGE2. *Arabian Journal of Chemistry*, **13**(6); 5639–5650
- Zheng, Y., M. Yates, H. Aung, Y. S. Cheng, C. Yu, H. Guo, R. Zhang, J. Vandergheynst, and B. M. Jenkins (2011). Influence of Moisture Content on Microbial Activity and Silage Quality During Ensilage of Food Processing Residues. *Bioprocess and Biosystems Engineering*, **34**(8); 987–995