

1. Submission (13 Maret 2024)
2. Review Round 1 (27 Maret 2024)
3. Review Round 2 (28 Maret 2024)
4. Editor decision (28 Mei 2024)
5. Accepted ( 29 Mei 2024)
6. Copy editing (2 Juni 2024)
7. Published (3 Juni 2024)

## Submissions

Workflow

Publication

Submission



Review

Copyediting

Production

### Submission Files

 Search

	3046-1	<a href="#">muslimmuslim, template-jsai.docx</a>	March 13, 2024	Article Text
	3074-1	<a href="#">muhdailami, Review 420-Article Text-3046-1-2-20240313.docx</a>	March 26, 2024	Article Text

[Download All Files](#)

### Pre-Review Discussions

[Add discussion](#)

Platform &  
workflow by  
OJS / PKP

## Growth and survival of bronze featherback (*Notopterus notopterus*, Pallas 1769) reared on net cages

Pertumbuhan dan Kelangsungan Hidup Ikan Putak (*Notopterus notopterus*, Pallas 1769) yang Dipelihara dalam Waring

Disubmit: xx November 2023, Direvisi: xx Februari 2024, Diterima: xx Februari 2024

### ABSTRAK

Ikan putak (*Notopterus notopterus*, Pallas 1769) dipelihara pada tiga kepadatan yang berbeda di waring untuk mengevaluasi efek kepadatan tebar terhadap pertumbuhan, tingkat kelangsungan hidup, dan efisiensi pakan. *N. notopterus* ditebar pada tiga kepadatan yang berbeda yaitu 1, 3, dan 5 ekor.m<sup>-2</sup> dengan tiga kali ulangan. Bobot awal ikan putak (rata-rata±std) masing-masing perlakuan yakni 55,40±3,40; 55,41±2,90; dan 56,07±1,06 g. Uji coba pertumbuhan berlangsung selama 60 hari, dari bulan Mei hingga Juli 2023. Bobot individu ikan di setiap waring dicatat setiap bulan. Bobot akhir (rata-rata ± std) ikan yang ditebar pada kepadatan 1, 3, dan 5 ekor.m<sup>-2</sup> mencapai 66,87±2,17, 67,63±3,06, dan 64,93±0,72 g, masing-masing. Nilai laju pertumbuhan spesifik adalah 0,45, 0,48, dan 0,34% hari<sup>-1</sup> pada suhu antara 25,1 dan 30,3°C. Nilai efisiensi pakan adalah 8,36, 9,04, dan 6,30%, dan tingkat kelangsungan hidup dihitung masing-masing 100, 100, dan 93,3%. Hasil penelitian menunjukkan bahwa kepadatan tebar memiliki pengaruh yang signifikan terhadap pertumbuhan dan efisiensi pakan *N. notopterus*.

**Kata kunci:** Asli Indonesia, budidaya ikan, domestikasi, ikan-pisau, komoditi ikan

### ABSTRACT

The bronze featherback (*Notopterus notopterus*, Pallas 1769) were reared at three different densities in the net cages to evaluate the effects of stocking density on growth, survival rate, and feed efficiency. The *N. notopterus* were stocked in three different densities of 1, 3, and 5 fish.m<sup>-2</sup> with three replicates and an initial body weight (mean±std) of 55.40±3.40, 55.41±2.90, and 56.07±1.06 g, respectively. The growth trial lasted for 60 days, from May to July 2023. The individual weights of fish in each net cage were recorded monthly. The final body weights (mean ±std) of the fish stocked at densities of 1, 3, and 5 fish.m<sup>-2</sup> reached 66.87±2.17, 67.63±3.06, and 64.93±0.72 g, respectively. The specific growth rate values were 0.45, 0.48, and 0.34% day<sup>-1</sup> at temperatures ranging between 25.1 and 30.3°C. The feed efficiency values were 8.36, 9.04, and 6.30%, and the survival rates were calculated at 100, 100, and 93.3%, respectively. The results revealed that stocking density had a significant effect on the growth and feed efficiency of *N. notopterus*.

**Keywords:** domestication, native Indonesia, fish farming, fishery commodities, knifefish.

### INTRODUCTION

Density is an important aquaculture parameter. It has a significant impact on the variable growth performance of fish (Liu *et al.*, 2017; Li

*et al.*, 2021). In fisheries production, stocking density refers to the number of fish initially cultivated per unit area (Battisti *et al.*, 2020). The stocking density of the fish is one of the most important factors to reflect and evaluate

**Comment [WU1]:** Add the location to the research title

**Comment [WU2]:** 1. It is necessary to add the core of the research, data analysis method.  
2. From the three different densities treatments, which treatment is the best?  
3. What your recommendation from this result of research

**Comment [WU3]:** Key word must be corrected and related to the research written

**Comment [WU4]:** Same as the Abstrak above.

the level of physiological stress (Barton, 2002). According to Riche *et al.*, (2013), stocking density has a significant impact on growth, water quality, and fish welfare in aquaculture systems, all of which are related to production efficiency (Yang *et al.*, 2020). The welfare and growth of fish can be affected positively or negatively by extreme low or high densities, depending on the species (Millán-Cubillo *et al.*, 2016; Liu *et al.*, 2019; Yang *et al.*, 2020). Fish wellbeing is especially affected by stocking density, which is a significant factor affecting fish health in commercial production (Li *et al.*, 2021). In aquaculture, stocking density is a crucial factor and a topic of frequent debate. It can cause chronic stress, leading to physiological changes such as stress responses, reduced growth, and health impairment. It is important to carefully consider stocking density to ensure the well-being of aquatic animals (Carbonara *et al.*, 2020). Determining optimal stocking densities is difficult due to the complex relationship between fish welfare and stocking density (Kozłowski & Piotrowska, 2023).

The bronze featherback (*Notopterus notopterus*, Pallas 1769) is one of the freshwater fish species. Its distribution in Asia includes Indonesia, Malaysia, Philippines, Vietnam, Thailand, Cambodia, Laos, Myanmar, Pakistan, India, and Bangladesh (Naem *et al.*, 2010; Gupta *et al.*, 2013; Mustafa *et al.*, 2014; Achakzai *et al.*, 2015; Mohanty & Samanta, 2016; Conallin *et al.*, 2023; Winn *et al.*, 2021). In Indonesia, *N. notopterus* is found in South Sumatra, Lampung, Riau, Riau Island, Jambi, Bangka Islands, and Kalimantan (Ammar *et al.*, 2014; Mulyani & Budijono, 2014; Rapita *et al.*, 2021; Muslim & Syaifudin, 2022; Muslim, 2023). *N. notopterus* belongs to the omnivorous fish group; its diet consists of insects, small fish, insect larvae, crustaceans, nematodes, shrimps, annelids, and detritus (Srivastava *et al.*, 2012; Achakzai *et al.*, 2015). *N. notopterus* is a nocturnal fish species. Its unique body shape has the potential to be

used as an ornamental fish commodity (Muslim *et al.*, 2023). Several studies on *N. notopterus* in Indonesia include aspects of reproductive biology (Gustomi *et al.*, 2016), feeding habits (Rapita *et al.*, 2021), rearing with different stocking densities and different feeds (Sukendi *et al.*, 2020), rearing in box container and aquarium (Muslim *et al.*, 2023), rearing in buckets (Muslim & Simanjuntak, 2023), and length-weight relationships and condition factors (Muslim, 2023).

In South Sumatra, Indonesia, *N. notopterus*, is highly valued. The price of fresh fish ranges from IDR 50.000 to 100.000 kg<sup>-1</sup>, depending on the size of the fish. As an ornamental fish, *N. notopterus* with a total length of 10 cm, is sold for IDR 20.000 to 30.000 fish<sup>-1</sup>. In addition, the price of pulverized *N. notopterus* meat ranges from IDR 100.000 to IDR 150.000 kg<sup>-1</sup>. This fish meat paste is used to make pempek and crackers, a culinary specialty of South Sumatra (Muslim *et al.*, 2023). The increasing demand for this fish has led to increased exploitation of this commodity. This increase in exploitation has caused the population of *N. notopterus* in the wild to decline dramatically. The production of *N. notopterus* is wild-caught, while aquaculture is not yet available. Studies on the rearing of *N. notopterus* into net cages are very important as a basic piece of information. This is the first step towards providing *N. notopterus* that has adapted to the cultural environment. This study focused on the growth of *N. notopterus* reared in net cages in the Kelekar River, South Sumatra, Indonesia. The objective of this study was to evaluate the effects of different stocking densities on the growth performance, survival rate, and feed efficiency of *N. notopterus*.

## MATERIAL AND METHODS

### 1. Study site and time

This study was conducted in Kelekar River, Tanjung Pering Village, North Indralaya District, Ogan Ilir

Comment [WU5]: Must be italic

Comment [WU6]: Consistency is needed in writing references according to author guidelines

Comment [WU9]: Why is the stocking density 1,3 fish/m<sup>2</sup> especially *N. notopterus*? Maybe from another species of fish that is related

Comment [WU7]: Need to check again in reference

Comment [WU8]: 2020?

Comment [WU10]: The objective must be in sync with the research title or vice versa

Regency, South Sumatra, at ordinate point 3°14'48.0012 "S 104°38'43.674 "E (Figure 1) in May-July 2023.




Figure 1. Map of the research location of the bronze featherback (*Notopterus notopterus*, Pallas 1769) in Kelekar River, Ogan Ilir Regency, South Sumatra

## 2. Experimental Design

This study used nine net cages measuring 1 m in length, 1 m in width, and 1 m in height. A net cage is made of high-density polyethylene by weaving it using a flat loom. Bamboos were anchored to the riverbank to serve as pillars for the net cage. Each corner of the net cage was strapped to the bamboo using a strap, and the bottom of the net cage was tied to a stone so that the net cage would not float. The water level in the net cage ranged from 70–80 cm from the water surface to the bottom. The experimental design used in the study was a complete randomized design. The treatments had different stocking densities: 1, 3, and 5 fish.m<sup>-2</sup>. Each treatment was repeated three times.

## 3. Fish Stocking

The *N. notopterus* used were caught by local fishermen. The body weight of the fish used ranged from 55 to 58 g, and the total length ranged from 19 to 20 cm. The number of fish was stocked in net cages according to the treatment (1, 3, and 5 fish.m<sup>-2</sup>). The body weight and total length of the fish were measured individually before being stocked into the net cages. The fish were

stocked into the net cage in the afternoon at 4:00 p.m.

## 4. Fish Rearing and Feeding

The *N. notopterus* were reared for 60 days (May–July 2023). During the rearing period, the fish were fed with seluang fish (*Rasbora* sp.) at a feeding rate of 3%. The frequency of feeding was twice a day, at 8:00 a.m. and 4:00 p.m. The seluang fish used as feed are live fish, not dead ones. The protein content of seluang fish is 15.25%, fat is 4.46%, ash is 15.2%, and fiber is 0.8%.

Comment [WU12]: The proximate composition of seluang fish must be complete.

## 5. Measured Parameters

The growth in body weight was calculated as follows:  $W = W_t - W_0$  (where  $W$  is growth in body weight (g),  $W_t$  is the final body weight of fish (g), and  $W_0$  is the initial body weight of fish (g)). The growth in total length was calculated as follows:  $L = L_t - L_0$  (where  $L$  is growth in total length (cm),  $L_t$  is the final total length of fish (cm), and  $L_0$  is the initial total length of fish (cm)). The specific growth rate (SGR) and feed efficiency (FE) were calculated as follows:

$$SGR = \frac{\ln W_t - \ln W_0}{t} \times 100$$

$$FE = \frac{(W_t + D) - W_0}{F} \times 100\%$$

where  $t$  is the day of feeding,  $W_0$  is the initial body weight of fish (g),  $W_t$  is the final body weight of fish (g),  $D$  is the body weight of the dead fish (g), and  $F$  is the amount of feed given (g).

The survival rate (SR) was calculated as follows:  $SR = [N_t/N_0] \times 100$  (where  $N_0$  is the number of fish at the beginning of the experiment (fish), and  $N_t$  is the number of fish at the end of the experiment (fish)). The water quality parameters measured included temperature using a thermometer, pH using a pH-meter, dissolved oxygen using a DO-meter, and ammonia using a spectrophotometer. Water temperature and pH measurements were taken daily at 7:00 a.m. and 5:00 p.m., while dissolved oxygen and ammonia were measured weekly.

Comment [WU11]: Referred to by whom?

## 7. Statistical Analysis

Data were represented as arithmetic means of individual weight (mean±std). Data on body weight growth, total length growth, specific growth rate, survival, and feed efficiency were analyzed with one-way analyses of variance (ANOVA) and the Duncan's multiple range test using SPSS version 20.0. Water quality data were analyzed descriptively.

## RESULTS AND DISCUSSION

This study successfully reared *N. notopterus* in net cages in the Kelekar River for 60 days. The growth performance, feed efficiency, and survival of *N. notopterus* from this study are presented in Table 1, Figure 2, and Figure 3. The growth of body weight, total length, and specific growth rate was best in the 3 fish.m<sup>-2</sup> treatment. There was no significant difference ( $P > 0.05$ ) between the 1 fish.m<sup>-2</sup> treatment and the 3 fish.m<sup>-2</sup> treatment. Feed efficiency in the three treatments was significantly different ( $P < 0.05$ ). The survival rate of fish was not significantly different for all treatments ( $P > 0.05$ ). Water quality parameters during rearing were within the tolerance range for the growth and survival of *N. notopterus*. The water temperature ranged from 25.1 to 30.3 °C, the pH value ranged from 5.63 to 7.15, and the dissolved oxygen ranged from 3.6 to 5.5 mg. L<sup>-1</sup>, the ammonia ranged from 0.09 to 0.017 mg. L<sup>-1</sup>, and the transparency of water ranged from 50 to 57 cm.





Figure 2. The final body weight and total length of the bronze featherback (*Notopterus notopterus*, Pallas 1769)




Figure 3. The feed efficiency of the bronze featherback (*Notopterus notopterus*, Pallas 1769)

The results of this study showed that different stocking density treatments had a significant effect on the growth of body weight and total length of *N. notopterus*. The best treatment for growth in body weight and total length of *N. notopterus* was a stocking density of 3 fish.m<sup>-2</sup>. A significant determining factor in fish productivity in cages is stocking density, which is based on the volume of water or surface area per fish. The higher stocking density causes more stress, which increases energy requirements and slows growth and food consumption (Aksungur et al., 2007; Jia et al., 2022). In teleost fish, growth processes follow steps that are characteristic for each species and are more or less directly under the control of environmental factors (Xu et al., 2022; Abdel-Latif et al., 2023). The fish depends on internal and external factors, which control or synchronize many activities or functions,

**Comment [WU13]:** Where did the data come from? Missing in water quality measurement methods?

**Comment [WU14]:** When compared with the 1 fish/m<sup>2</sup> treatment, why is the 3 fish/m<sup>2</sup> treatment better even though the density is higher and can cause stress?



including growth performance (Bœuf & Payan, 2001). Internal factors include health, stress, and reproductive status, while external factors include food quality and quantity, water temperature, and some water quality parameters. In the 3 fish.m<sup>-2</sup> treatment, feed utilization by fish is more efficient than other treatments. Competition for food and niches is low, so food is fully utilized for the growth process. Fish reared in low densities will be more aggressive (Andersson *et al.*, 2022). The body weight growth of *N. notopterus* in this study was higher than the results of Sukendi *et al.*, (2020).

The specific growth rate of *N. notopterus* in this study ranged from 0.34% to 0.48%. Specific growth rate is closely related to body weight growth. The growth of body weight is in line with the specific growth rate; the higher the growth of body weight, the higher the specific growth rate. In this study, the highest specific growth rate was observed in the stocking density treatment of 3 fish.m<sup>-2</sup>. According to Li *et al.*, (2021), the specific growth rate of fish increases significantly at low densities. A low stocking density provides more space for movement than a high stocking density, resulting in less competition for niches and food (Ullah *et al.*, 2018). The results of this study showed that different stocking density treatments significantly affected the feed efficiency of *N. notopterus* reared. The highest feed efficiency was 9.04% in the stocking density treatment of 3 fish.m<sup>-2</sup>. Feed efficiency value is closely related to the growth value of fish. The feed efficiency

is an essential indicator, as high-quality feed and effective aquaculture management can reduce the amount of feed used and result in better fish growth with less feed consumed (Zlaugotne *et al.*, 2022).

The survival of *N. notopterus* in the treatment of 1 and 3 fish.m<sup>-2</sup> was 100.00%, and in the treatment of 5 fish.m<sup>-2</sup>, it was 93.33%. These results show that stocking density has no significant effect on survival. According to Budiardi *et al.*, (2007), the optimal stocking density is when a high number of fish are stocked, but competition for food and space is still tolerated by the fish, resulting in high survival rates, fish growth rates, and low size variations. Water quality plays a critical role as it has a direct impact on the overall health status of the reared fish (Abdel-Aziz *et al.*, 2021). The results of this study indicate that water quality was favorable for the growth and survival of *N. notopterus*. The water quality of the *N. notopterus* habitat in the Kelekar Floodplain was as follows: water temperature ranged from 24 to 31°C, pH ranged from 4 to 7, and dissolved oxygen ranged from 4.32 to 7.61 mg. L<sup>-1</sup> (Muslim & Syaifudin, 2022). The water quality of *N. notopterus* rearing media in the box container was as follows: water temperature ranged from 27 to 29.7°C, pH ranged from 6 to 7.92, dissolved oxygen ranged from 4.5 to 7.8 mg. L<sup>-1</sup>, and ammonia ranged from 0.041 to 0.055 mg. L<sup>-1</sup> (Muslim *et al.*, 2023).

**Comment [WU15]:** Must be update references

**Comment [WU17]:** Does the quality of the water obtained comply with SNI for the *N. notopterus*?

**Comment [WU16]:** This statement contra with the results of your research

Tabel 1. The growth performance, feed efficiency, and survival of the bronze featherback (*Notopterus notopterus*, Pallas 1769)

Parameters	Treatments		
	1 fish.m <sup>-2</sup>	3 fish.m <sup>-2</sup>	5 fish.m <sup>-2</sup>
Initial body weight (g)	55.40±3.40 <sup>a</sup>	55.41±2.90 <sup>a</sup>	56.07±1.06 <sup>a</sup>
Final body weight (g)	66.87±2.17 <sup>b</sup>	67.63±3.06 <sup>b</sup>	64.93±0.72 <sup>a</sup>

**Comment [WU18]:** This data has been found in Figure 2. It's best not to repeat it on the table again



Growth in body weight (g)	11.47±1.29 <sup>b</sup>	12.22±1.09 <sup>b</sup>	8.86±0.57 <sup>a</sup>
Initial total length (cm)	20.03±0.42 <sup>a</sup>	20.16±0.40 <sup>a</sup>	19.87±0.09 <sup>a</sup>
Final total length (cm)	22.30±0.70 <sup>b</sup>	22.52±0.33 <sup>b</sup>	21.33±0.06 <sup>a</sup>
Growth in total length (cm)	2.27±0.32 <sup>b</sup>	2.37±0.12 <sup>b</sup>	1.46±0.10 <sup>a</sup>
Specific growth rate (% day <sup>-1</sup> )	0.45±0.07 <sup>b</sup>	0.48±0.05 <sup>b</sup>	0.34±0.03 <sup>a</sup>
Feed efficiency (%)	8.36±1.03 <sup>b</sup>	9.04±0.81 <sup>c</sup>	6.30±0.05 <sup>a</sup>
Survival rate (%)	100.00±0.00 <sup>a</sup>	100.00±0.00 <sup>a</sup>	93.33±11.55 <sup>a</sup>

Note: mean±std in the same column with different superscript (a, b, c) are significantly different (P < 0.05)

## CONCLUSION

In conclusion, the results of this study showed that stocking density affected the growth performance and feed efficiency of *N. notopterus* but not its survival. This study indicated that *N. notopterus* can be adapted to a net cage. This study provides a valuable reference for the domestication of *N. notopterus* in the future.

## ACKNOWLEDGEMENT

We are grateful to Mr. Nurrohman, a fisherman from the village of Tanjung Pring, who has been a great help in the technical work in the field.

## REFERENCES

- Abdel-Aziz, M.F.A., Hassan, H.U., Yones, A.M., Abdel-Tawwab, Y.A., & Metwalli, A.A.A.T. (2021). Assessing the effect of different feeding frequencies combined with stocking density, initial weight, and dietary protein ratio on the growth performance of tilapia, catfish and carp. *Scientific African*, 12, e00806. <https://doi.org/10.1016/j.sciaf.2021.e00806>
- Abdel-Latif, H.M.R., Ahmed, H.A., Shukry, M., Khafaga, A.F., Elkhayat, B.K., Abdel-Tawwab, M., & Abd-elaziz, R.A. (2023). Growth performance, physiological responses, and histoarchitectural changes in juvenile *Pangasianodon hypophthalmus* under different environmental salinities. *Fishes*, 8(6), 282. <https://doi.org/10.3390/fishes8060282>
- Achakzai, W. M., Saddozai, S., Baloch, W. A., Soomro, A. N., & Memon, N. (2015). Length-weight relationship and condition factor of *Notopterus notopterus* (Pallas, 1769) from Manchar Lake Sindh, Pakistan. *Sindh University Research Journal-SURJ (Science Series)*, 47(3), 515-518.
- Aksungur, N., Aksungur, M., Akbulut, B., & Kutlu, İ. (2007). Effects of stocking density on growth performance, survival and food conversion ratio of Turbot (*Psetta maxima*) in the net cages on the southeastern coast of the Black Sea. *Turkish Journal of Fisheries and Aquatic Sciences*, 7(2), 147-152.
- Ammar, J. A., & Kamal, M. M. (2014). Keragaman ikan di Danau Cala, Kabupaten Musi Banyuasin Sumatera Selatan. *Depik*, 3(3), 216-220. <https://doi.org/10.13170/depik.3.3.2147>
- Andersson, M., Roques, J. A., Aliti, G. M., Ademar, K., Sundh, H., Sundell, K., Ericson, M. & Kettunen, P. (2022). Low holding densities increase stress response and aggression in zebrafish. *Biology*, 11(5), 725.

**Comment [WU19]:** This data has been found in Figure 2. It's best not to repeat it on the table again

**Comment [WU20]:** This data has been found in Figure 3. It's best not to repeat it on the table again

**Comment [WU21]:** It needs to be added, the best treatment so that it becomes a reference for domestication

- <https://doi.org/10.3390/biology11050725>
- Barton, B. A. (2002). Stress in fishes: a diversity of responses with particular reference to changes in circulating corticosteroids. *Integrative and comparative biology*, 42(3), 517-525.  
<https://doi.org/10.1093/icb/42.3.517>
- Battisti, E. K., Rabaioli, A., Uczay, J., Sutili, F. J., & Lazzari, R. (2020). Effect of stocking density on growth, hematological and biochemical parameters and antioxidant status of silver catfish (*Rhamdia quelen*) cultured in a biofloc system. *Aquaculture*, 524, 735213.  
<https://doi.org/10.1016/j.aquaculture.2020.735213>
- Bœuf, G., & Payan, P. (2001). How should salinity influence fish growth?. *Comparative Biochemistry and Physiology Part C: Toxicology and Pharmacology*, 130(4), 411-423.  
[https://doi.org/10.1016/s1532-0456\(01\)00268-x](https://doi.org/10.1016/s1532-0456(01)00268-x)
- Budiardi, T., Gemawaty, N., & Wahjuningrum, D. (2007). Production of *Paracheirodon innesi* on different densities in recirculating system. *Jurnal Akuakultur Indonesia*, 6(2), 211-215.  
<https://journal.ipb.ac.id/index.php/jai/article/view/4030>
- Carbonara, P., Alfonso, S., Gai, F., Gasco, L., Palmegiano, G., Spedicato, M. T., Zupa, W., & Lembo, G. (2020). Moderate stocking density does not influence the behavioural and physiological responses of rainbow trout (*Oncorhynchus mykiss*) in organic aquaculture. *Aquaculture Research*, 51(7), 3007-3016.  
<https://doi.org/10.1111/are.14640>
- Conallin, J., Tun, N. N., Swe, A. M., Baumgartner, L. J., Lunn, Z., Mallen-Cooper, M., ... & Thew, P. (2023). Using fish swimming ability to refine criteria for fishway construction in Myanmar. *Fisheries Research*, 262, 106680.  
<https://doi.org/10.1016/j.fishres.2023.106680>
- Gupta, A., Lal, K. K., Punia, P., Singh, R. K., Mohindra, V., Sah, R. S., Kumar, J., Luhariya, R.K., Dwivedi, A.K., Masih, P., Mishra, R.M., & Jena, J. K. (2013). Characterization of polymorphic microsatellite markers and genetic diversity in wild bronze featherback, *Notopterus notopterus* (Pallas, 1769). *Molecular Biology Reports*, 40, 6625-6631.  
<https://doi.org/10.1007/s11033-013-2776-z>
- Gustomi, A., Sulistiono, S., & Yonvitner, Y. (2016). Reproductive biology feather back (*Notopterus notopterus*, Pallas, 1769) in Simpung Reservoir, Bangka Island. *Indonesian Journal of Agricultural Sciences*, 21(1), 56-62.  
<https://doi.org/10.18343/jipi.21.1.56>
- Jia, R., Wang, L., Hou, Y., Feng, W., Li, B., & Zhu, J. (2022). Effects of stocking density on the growth performance, physiological parameters, redox status and lipid metabolism of *Micropterus salmoides* in integrated rice–fish farming systems. *Antioxidants*, 11(7), 1215.  
<https://doi.org/10.3390/antiox11071215>
- Kozłowski, M., & Piotrowska, I. (2023). Effect of stocking density on

- growth, survival and cannibalism of juvenile pikeperch, *Sander lucioperca* (L.), in a recirculating aquaculture system. *Aquaculture International*, 1-9. <https://doi.org/10.1007/s10499-023-01339-6>
- Li, L., Shen, Y., Yang, W., Xu, X., & Li, J. (2021). Effect of different stocking densities on fish growth performance: A meta-analysis. *Aquaculture*, 544, 737152. <https://doi.org/10.1016/j.aquaculture.2021.737152>
- Liu, B., Jia, R., Zhao, K., Wang, G., Lei, J., & Huang, B. (2017). Stocking density effects on growth and stress response of juvenile turbot (*Scophthalmus maximus*) reared in land-based recirculating aquaculture system. *Acta Oceanologica Sinica*, 36, 31-38. <https://doi.org/10.1007/s13131-017-0976-4>
- Liu, Y., Liu, H., Wu, W., Yin, J., Mou, Z., & Hao, F. (2019). Effects of stocking density on growth performance and metabolism of juvenile Lenok (*Brachymystax lenok*). *Aquaculture*, 504, 107-113. <https://doi.org/10.1016/j.aquaculture.2019.01.058>
- Millán-Cubillo, A. F., Martos-Sitcha, J. A., Ruiz-Jarabo, I., Cárdenas, S., & Mancera, J. M. (2016). Low stocking density negatively affects growth, metabolism and stress pathways in juvenile specimens of meagre (*Argyrosomus regius*, Asso 1801). *Aquaculture*, 451, 87-92. <https://doi.org/10.1016/j.aquaculture.2015.08.034>
- Mohanty, D., & Samanta, L. (2016). Multivariate analysis of potential biomarkers of oxidative stress in *Notopterus notopterus* tissues from Mahanadi River as a function of concentration of heavy metals. *Chemosphere*, 155, 28-38. <https://doi.org/10.1016/j.chemosphere.2016.04.035>
- Mulyani, I., & Budijono, B. (2020). Morphometric and meristic analysis of Asian knifefish (*Notopterus notopterus*) in Sail River, Pekanbaru Riau Province. *Jurnal Ilmiah Biologi Eksperimen dan Keanekaragaman Hayati*, 7(2), 59-64.
- Muslim, M. (2023). Length-weight relationship and condition factor of *Notopterus notopterus* (Pallas, 1769) from East Pedamaran Floodplain, Ogan Komering Ilir, South Sumatra, Indonesia. *International Journal of Advanced Multidisciplinary Research and Studies*, 3(5), 1227-1231.
- Muslim, M., & Simanjuntak, W. J. (2023). Growth and survival of bronze featherback (*Notopterus notopterus*, Pallas 1769) reared on bucket. *Magna Scientia Advanced Research and Reviews*, 9(1), 101-105. <https://doi.org/10.30574/msarr.2023.9.1.0133>
- Muslim, M., & Syaifudin, M. (2022). Biodiversity of freshwater fish in Kelekar Floodplain Ogan Ilir Regency in Indonesia. *Journal of Tropical Biodiversity and Biotechnology*, 7(1), 1-10. <https://doi.org/10.22146/jtbb.67494>
- Muslim, M., Pitriani, E., & Agustina, H. (2023). Growth and survival of bronze featherback (*Notopterus notopterus*) adapted on box container and aquarium. *GSC Advanced Research and Reviews*, 16(3), 133-137. <https://doi.org/10.30574/gscarr.2023.16.3.0368>

- Mustafa MG, Singha S, Islam M, Mallick N. Population dynamics of *Notopterus notopterus* (Pallas, 1769) from the Kaptai Reservoir of Bangladesh. *SAARC Journal of Agriculture* 2014; 12(2): 112–122. <https://doi.org/10.3329/sja.v12i2.21920>
- Mustafa, M. G., Singha, S., Islam, M. R., & Nayan Mallick, N. M. (2014). Population dynamics of *Notopterus notopterus* (Pallas, 1769) from the Kaptai Reservoir of Bangladesh. *SAARC Journal of Agriculture*, 12(2), 112–122. <https://doi.org/10.3329/sja.v12i2.21920>
- Naeem, M., Salam, A., Gillani, Q., & Ishtiaq, A. (2010). Length-weight relationships of *Notopterus notopterus* and introduced *Oreochromis niloticus* from the Indus River, southern Punjab, Pakistan. *Journal of Applied Ichthyology*, 26(4), 620–620. <https://doi.org/10.1111/j.1439-0426.2010.01480.x>
- Rapita, R., Susiana, S., & Kurniawan, D. (2021). Food habits of belida fish (*Notopterus notopterus*, Pallas 1769) in Sei Gesek Reservoir, Bintan Regency, Riau Island, Indonesia. *IOP Conference Series. Earth Environmental Sci*, 919, 012003. <https://doi.org/10.1088/1755-1315/919/1/012003>
- Riche, M. A., Weirich, C. R., Wills, P. S., & Baptiste, R. M. (2013). Stocking density effects on production characteristics and body composition of market size cobia, *Rachycentron canadum*, reared in recirculating aquaculture systems. *Journal of the World Aquaculture Society*, 44(2), 259–266. <https://doi.org/10.1111/jwas.12023>
- Srivastava, S. M., Singh, S. P., & Pandey, A. K. (2012). Food and feeding habits of threatened *Notopterus notopterus* in Gomti river, Lucknow (India). *Journal of Experimental Zoology*, 15(2), 395–402.
- Sukendi, S., Thamrin, T., Putra, R. M., & Yulindra, A. (2020). Cultivation technology of bronze featherback (*Notopterus notopterus*, Pallas 1769) at different stocking densities and types of feed. *IOP Conference Series: Earth and Environmental Science*, 430(1), 012027. <https://doi.org/10.1088/1755-1315/430/1/012027>
- Ullah, K., Emmanuel, A., & Anjum, M. Z. (2018). Effect of stocking density on growth performance of Indus mahseer (*Tor macrotepis*). *International Journal of Fisheries and Aquatic Studies*, 6(3), 49–52.
- Winn, N. A., Sandi, P., Khaing, T., Nyunt, K. T., Kyaw, H. T., Sabai, M., & Aung, T. T. N. (2021). Length weight relationship of twelve freshwater fish species from Sunye Lake, Mandalay Region, Myanmar. *Greener Journal of Biological Sciences*, 11(2), 74–80.
- Xu, W., Lutz, C. G., Taylor, C. M., & Ortega, M. C. (2022). Improvement of fish growth and metabolism by oligosaccharide prebiotic supplement. *Aquaculture Nutrition*, 28, 5715649. <https://doi.org/10.1155/2022/5715649>
- Yang, Q., Guo, L., Liu, B. S., Guo, H. Y., Zhu, K. C., Zhang, N., ... & Zhang, D. C. (2020). Effects of stocking density on the growth performance, serum biochemistry, muscle composition and HSP70 gene expression of juvenile golden

pompano *Trachinotus ovatus*  
(Linnaeus,  
1758). *Aquaculture*, 518, 734841.  
<https://doi.org/10.1016/j.aquaculture.2019.734841>

Zlaugotne, B., Pubule, J., & Blumberga,  
D. (2022). Advantages and  
disadvantages of using more  
sustainable ingredients in fish  
feed. *Heliyon*, 8(9), e10527.  
<https://doi.org/10.1016/j.heliyon.2022.e10527>

**Comment [WU22]:** Double article

**Comment [WU23]:** 2014?

**Comment [WU24]:**

## Growth and survival of bronze featherback (*Notopterus notopterus*, Pallas 1769) reared on net cages

Pertumbuhan dan Kelangsungan Hidup Ikan Putak (*Notopterus notopterus*, Pallas 1769) yang Dipelihara dalam Waring

Disubmit: xx November 2023, Direvisi: xx Februari 2024, Diterima: xx Februari 2024

### ABSTRAK

Ikan putak (*Notopterus notopterus*, Pallas 1769) dipelihara pada tiga kepadatan yang berbeda di waring untuk mengevaluasi efek dampak kepadatan tebar yang berbeda terhadap pertumbuhan, tingkat kelangsungan hidup, dan efisiensi pakan. *N. notopterus* ditebar pada tiga kepadatan yang berbeda yaitu 1, 3, dan 5 ekor.m<sup>-2</sup> dengan tiga kali ulangan. Bobot awal ikan putak (rata-rata ± std) masing-masing perlakuan yakni 55,40±3,40; 55,41±2,90; dan 56,07±1,06 g. Uji coba pertumbuhan berlangsung selama 60 hari, dari bulan Mei hingga Juli 2023. Bobot individu ikan di setiap waring dicatat setiap bulan. Bobot akhir (rata-rata ± std) ikan yang ditebar pada kepadatan 1, 3, dan 5 ekor.m<sup>-2</sup> mencapai 66,87±2,17, 67,63±3,06, dan 64,93±0,72 g, masing-masing. Nilai laju pertumbuhan spesifik adalah 0,45, 0,48, dan 0,34% hari<sup>-1</sup> pada suhu antara 25,1 dan 30,3°C. Nilai efisiensi pakan adalah 8,36, 9,04, dan 6,30%, dan tingkat kelangsungan hidup dihitung masing-masing 100, 100, dan 93,3%. Hasil penelitian menunjukkan bahwa kepadatan tebar memiliki pengaruh yang signifikan terhadap pertumbuhan dan efisiensi pakan *N. notopterus*.

**Kata kunci:** Asli Indonesia, budidaya ikan, domestikasi, ikan-pisau ????, komoditi ikan

### ABSTRACT

The bronze featherback (*Notopterus notopterus*, Pallas 1769) were reared at three different densities in the net cages to evaluate the effects of stocking density on growth, survival rate, and feed efficiency. The *N. notopterus* were stocked in three different densities of 1, 3, and 5 fish.m<sup>-2</sup> with three replicates and an initial body weight (mean ± std) of 55.40±3.40, 55.41±2.90, and 56.07±1.06 g, respectively. The growth trial lasted for 60 days, from May to July 2023. The individual weights of fish in each net cage were recorded monthly. The final body weights (mean ± std) of the fish stocked at densities of 1, 3, and 5 fish.m<sup>-2</sup> reached 66.87±2.17, 67.63±3.06, and 64.93±0.72 g, respectively. The specific growth rate values were 0.45, 0.48, and 0.34% day<sup>-1</sup> at temperatures ranging between 25.1 and 30.3°C. The feed efficiency values were 8.36, 9.04, and 6.30%, and the survival rates were calculated at 100, 100, and 93.3%, respectively. The results revealed that stocking density had a significant effect on the growth and feed efficiency of *N. notopterus*.

**Keywords:** domestication, native Indonesia, fish farming, fishery commodities, knifefish.

### INTRODUCTION

Density is an important aquaculture parameter. It has a significant impact on the variable growth

performance of fish (Liu et al., 2017; Li et al., 2021). In fisheries production, stocking density refers to the number of fishes initially cultivated per unit area (Battisti et al., 2020). The stocking

**Comment [L1]:** Say rasa tidak perlu ditampilkan

**Comment [L2]:** Abstrak yang baik terdiri dari pendahuluan berisi permasalahan yang menjadi tujuan penelitian, metode singkat se parameter yang diukur, hasil dan kesimpulan abstrak yang ada belum menunjukkan hal ini

**Comment [L3]:** Kata kunci yang baik, seharusnya ada di dalam narasi abstrak dan menjadi hal yang penting, dan sebaiknya tidak ada dalam judul, kata kunci yg ada tidak sama sekali didalam abstrak !

**Comment [L4]:** I think it doesn't need to be written



density of the fish is one of the most important factors to reflect and evaluate the level of physiological stress (Barton, 2002). According to Riche *et al.*, (2013), Stocking density has a significant impact on growth, water quality, and fish welfare in aquaculture systems (Riche *et al.*, 2013). ~~all of this impact which are have~~ related to production efficiency (Yang *et al.*, 2020). The welfare and growth of fish can be affected positively or negatively by extreme low or high densities, depending on the species (Millán-Cubillo *et al.*, 2016; Liu *et al.*, 2019; Yang *et al.*, 2020). Fish wellbeing is especially affected by stocking density, which is a significant factor affecting fish health in commercial production (Li *et al.*, 2021). In aquaculture, stocking density is a crucial factor and a topic of frequent debate. It can cause chronic stress, leading to physiological changes such as stress responses, reduced growth, and health impairment. It is important to carefully consider stocking density to ensure the well-being of aquatic animals (Carbonara *et al.*, 2020). Determining optimal stocking densities is difficult due to the complex relationship between fish welfare and stocking density (Kozłowski & Piotrowska, 2023).

The bronze featherback (*Notopterus notopterus*, Pallas 1769) is one of the freshwater fish species. Its distribution in Asia includes Indonesia, Malaysia, Philippines, Vietnam, Thailand, Cambodia, Laos, Myanmar, Pakistan, India, and Bangladesh (Naeem *et al.*, 2010; Gupta *et al.*, 2013; Mustafa *et al.*, 2014; Achakzai *et al.*, 2015; Mohanty & Samanta, 2016; Conallin *et al.*, 2023; Winn *et al.*, 2021). In Indonesia, *N. notopterus* is found in South Sumatra, Lampung, Riau, Riau Island Jambi, Bangka Islands, and Kalimantan (Ammar *et al.*, 2014; Mulyani & Budijono, 2014; Rapita *et al.*, 2021; Muslim & Syaifudin, 2022; Muslim, 2023). *N. notopterus* belongs to the omnivorous fish group; its diet consists of insects, small fish, insect larvae, crustaceans, nematodes, shrimps, annelids, and detritus (Srivastava *et al.*,

2012; Achakzai *et al.*, 2015). *N. notopterus* is a nocturnal fish species. ~~It~~ has unique ~~body with~~ shape ~~body and~~ has the potential to be used as an ornamental fish commodity (Muslim *et al.*, 2023). Several studies on *N. notopterus* in Indonesia ~~has done, include like~~ ~~aspects~~ aspects of reproductive biology (Gustomi *et al.*, 2016), feeding habits (Rapita *et al.*, 2021), rearing with different stocking densities and different feeds (Sukendi *et al.*, 2020), rearing in box container and aquarium (Muslim *et al.*, 2023), rearing in buckets (Muslim & Simanjuntak, 2023), and length-weight relationships and condition factors (Muslim, 2023).

In South Sumatra, Indonesia, *N. notopterus*, is highly valued. The price of fresh fish ranges from IDR 50.000 to 100.000 kg<sup>-1</sup>, depending on the size of the fish. As an ornamental fish, *N. notopterus* with a total length of 10 cm, is sold for IDR 20.000 to 30.000 fish<sup>-1</sup>. In addition, the price of pulverized *N. notopterus* meat ranges from IDR 100.000 to ~~IDR~~ 150.000 kg<sup>-1</sup>. This fish meat paste is used to make pempek and crackers, a culinary specialty of South Sumatra (Muslim *et al.*, 2023). The increasing demand for this fish has led to increased exploitation of this commodity. This increase in exploitation has caused the population of *N. notopterus* in the wild to decline dramatically. The production of *N. notopterus* ~~still from is~~ wild-caught, while aquaculture is not yet available. ~~Studies on the rearing of~~ ~~N. notopterus~~ into net cages are very important as a basic piece of information. This is the first step towards providing *N. notopterus* that has adapted to the cultural environment. This study focused on the growth of *N. notopterus* reared in net cages in the Kelekar River, South Sumatra, Indonesia. The objective of this study was to evaluate the effects of different stocking densities on the growth performance, survival rate, and feed efficiency of *N. notopterus*.

## MATERIAL AND METHODS

**Comment [L5]:** Sebaiknya langsung fokus pada belum ada kajian tentang padat tebar, atau perlu informasi berapa padat tebar yang optimum dalam pemeliharaan ikan platak di keramba sehingga ruh pentignya penelitian ini terlihat



### 1. Study site and time

This study was conducted in Kelekar River, Tanjung Pering Village, North Indralaya District, Ogan Ilir Regency, South Sumatra, at ordinate point 3°14'48.0012 "S 104°38'43.674 "E (Figure 1) in May-July 2023.




Figure 1. Map of the research location of the bronze featherback (*Notopterus notopterus*, Pallas 1769) in Kelekar River, Ogan Ilir Regency, South Sumatra

### 2. Experimental Design

This study used nine net cages measuring 1 m in length, 1 m in width, and 1 m in height or depth. A net cage is made of high-density polyethylene by weaving it using a flat loom. Bamboos were anchored to the riverbank to serve as pillars for the net cage. Each corner of the net cage was strapped to the bamboo using a strap, and the bottom of the net cage was tied to a stone so that the net cage would not float. The water level in the net cage ranged from 70–80 cm from the water surface to the bottom. The experimental design used in the study was a complete randomized design. The treatments had different stocking densities, i.e., 1, 3, and 5 fish.m<sup>-2</sup>, and each treatment was repeated three times.

### 3. Fish Stocking

The *N. notopterus* used were caught by local fishermen. The body weight of the fish used ranged from 55 to 58 g, and the total length ranged from 19 to 20 cm. The number of fishes was stocked in net cages according to the

treatment (1, 3, and 5 fishes.m<sup>-2</sup>). The body weight and total length of the fish were measured individually before being stocked into the net cages. The fish were stocked into the net cage in the afternoon at 4:00 p.m. for what? Why must at 4 pm?.

### 4. Fish Rearing and Feeding

The *N. notopterus* were reared for 60 days (May–July 2023). During the rearing period, the fish were fed with seluang (international name?) fish (*Rasbora* sp.) at a feeding rate of 3% (daily? weekly? Monthly?). The frequency of feeding was twice a day, at 8:00 a.m. and 4:00 p.m. The seluang (international name?) fish used as feed are live fish, not dead ones. The protein content of seluang (international name?) fish is 15.25%, fat is 4.46%, ash is 15.2%, and fiber is 0.8% (sumber data dari mana?).

### 5. Measured Parameters

The growth in body weight was calculated as follows:  $W = W_t - W_0$  (where  $W$  is growth in body weight (g),  $W_t$  is the final body weight of fish (g), and  $W_0$  is the initial body weight of fish (g)). The growth in total length was calculated as follows:  $L = L_t - L_0$  (where  $L$  is growth in total length (cm),  $L_t$  is the final total length of fish (cm), and  $L_0$  is the initial total length of fish (cm)). The specific growth rate (SGR) and feed efficiency (FE) were calculated as follows:

$$SGR = \frac{\ln W_t - \ln W_0}{t} \times 100$$

$$FE = \frac{(W_t + D) - W_0}{F} \times 100\%$$

where  $t$  is the day of feeding,  $W_0$  is the initial body weight of fish (g),  $W_t$  is the final body weight of fish (g),  $D$  is the body weight of the dead fish (g), and  $F$  is the amount of feed given (g).

The survival rate (SR) was calculated as follows:  $SR = [N_t/N_0] \times 100$  (where  $N_0$  is the number of fishes at the beginning of the experiment (fish), and  $N_t$  is the number of fishes at the end of the experiment (fish)). The water

**Comment [L6]:** Jika 1 ekor m<sup>2</sup> jadi P1, 3 ekor jadi P2, dan 5 ekor jadi P3 bagaimana? Sehingga dalam proses penulisan hasil dan pembahasan akan lebih efisien dan mudah dipahami....

quality parameters measured included temperature using a thermometer, pH using a pH-meter, dissolved oxygen using a DO-meter, and ammonia using a spectrophotometer. Water temperature and pH measurements were taken daily at 7:00 a.m. and 5:00 p.m., while dissolved oxygen and ammonia were measured weekly.

### 7. Statistical Analysis

Data were represented as arithmetic means of individual weight (mean±std). Data on body weight growth, total length growth, specific growth rate, survival, and feed efficiency were analyzed with one-way analyses of variance (ANOVA) and the Duncan's multiple range test using SPSS version 20.0. Water quality data were analyzed descriptively.

## RESULTS AND DISCUSSION

This study successfully reared *N. notopterus* in net cages in the Kelekar River for 60 days. The growth performance, feed efficiency, and survival of *N. notopterus* from this study are presented in Table 1, Figure 2, and Figure 3. The growth of body weight, total length, and specific growth rate was best in the 3 fish.m<sup>-2</sup> treatment. There was no significant difference ( $P > 0.05$ ) between the 1 fish.m<sup>-2</sup> treatment and the 3 fish.m<sup>-2</sup> treatment. Feed efficiency in the three treatments was significantly different ( $P < 0.05$ ). The survival rate of fish was not significantly different for all treatments ( $P > 0.05$ ). Water quality parameters during rearing were within the tolerance range for the growth and survival of *N. notopterus*. The water temperature ranged from 25.1 to 30.3 °C, the pH value ranged from 5.63 to 7.15, and the dissolved oxygen ranged from 3.6 to 5.5 mg. L<sup>-1</sup>, the ammonia ranged from 0.09 to 0.017 mg. L<sup>-1</sup>, and the transparency of water ranged from 50 to 57 cm.




Figure 2. The final body weight and total length of the bronze featherback (*Notopterus notopterus*, Pallas 1769)




Figure 3. The feed efficiency of the bronze featherback (*Notopterus notopterus*, Pallas 1769)

The results of this study showed that different stocking density treatments had a significant effect on the growth of body weight and total length of *N. notopterus*. The best treatment for growth in body weight and total length of *N. notopterus* was a stocking density of 3 fish.m<sup>-2</sup>. A significant determining factor in fish productivity in cages is stocking density, which is based on the volume of water or surface area per fish. The higher stocking density causes more stress, which increases energy requirements and slows growth and food consumption

Comment [L7]: I think it doesn't need to be written

Comment [L9]: I think just delete pictures 2 and 3 because they display the same data as table 1

Comment [L8]: I think water quality will be best to show in table...

(Aksungur *et al.*, 2007; Jia *et al.*, 2022). In teleost fish, growth processes follow steps that are characteristic for each species and ~~are more or less directly~~ under the control of environmental factors (Xu *et al.*, 2022; Abdel-Latif *et al.*, 2023). The fish depends on internal and external factors, which control or synchronize many activities or functions, including growth performance (Bøuf & Payan, 2001). Internal factors include health, stress, and reproductive status, while external factors include food quality and quantity, water temperature, and some water quality parameters. In the 3 fish.m<sup>-2</sup> treatment, feed utilization by fish is more efficient than other treatments. Competition for food and niches is low, so food is fully utilized for the growth process. Fish reared in low densities will be more aggressive (Andersson *et al.*, 2022). The body weight growth of *N. notopterus* in this study was higher than the results of Sukendi *et al.*, (2020).

The specific growth rate of *N. notopterus* in this study ranged from 0.34% to 0.48%. Specific growth rate is closely related to body weight growth. The growth of body weight is in line with the specific growth rate; the higher the growth of body weight, the higher the specific growth rate. In this study, the highest specific growth rate was observed in the stocking density treatment of 3 fish.m<sup>-2</sup>. According to Li *et al.*, (2021), the specific growth rate of fish increases significantly at low densities. A low stocking density provides more space for movement than a high stocking density, resulting in less competition for niches and food (Ullah *et al.*, 2018). The results of this study showed that different stocking density treatments significantly affected the feed efficiency of *N. notopterus* reared. The highest feed

efficiency was 9.04% in the stocking density treatment of 3 fish.m<sup>-2</sup>. Feed efficiency value is closely related to the growth value of fish. The feed efficiency is an essential indicator, as high-quality feed and effective aquaculture management can reduce the amount of feed used and result in better fish growth with less feed consumed (Zlaugotne *et al.*, 2022).

The survival of *N. notopterus* in the treatment of 1 and 3 fish.m<sup>-2</sup> was 100.00%, and in the treatment of 5 fish.m<sup>-2</sup>, it was 93.33%. These results show that stocking density has no significant effect on survival. According to Budiardi *et al.*, (2007), the optimal stocking density is when a high number of fishes are stocked, but competition for food and space is still tolerated by the fish, resulting in high survival rates, fish growth rates, and low size variations. Water quality plays a critical role as it has a direct impact on the overall health status of the reared fish (Abdel-Aziz *et al.*, 2021). The results of this study indicate that water quality was favorable for the growth and survival of *N. notopterus*. The water quality of the *N. notopterus* habitat in the Kelekar Floodplain was as follows: water temperature ranged from 24 to 31°C, pH ranged from 4 to 7, and dissolved oxygen ranged from 4.32 to 7.61 mg. L<sup>-1</sup> (Muslim & Syaifudin, 2022). The water quality of *N. notopterus* rearing media in the box container was as follows: water temperature ranged from 27 to 29.7°C, pH ranged from 6 to 7.92, dissolved oxygen ranged from 4.5 to 7.8 mg. L<sup>-1</sup>, and ammonia ranged from 0.041 to 0.055 mg. L<sup>-1</sup> (Muslim *et al.*, 2023).

**Comment [L10]:** discussions of water quality should focus on whether there is an impact of stocking density on water quality? Is the water quality still within the tolerance range for research object fish?

**Tabel 1.** The growth performance, feed efficiency, and survival of the bronze featherback (*Notopterus notopterus*, Pallas 1769)

Parameters	Treatments		
	1 fish.m <sup>-2</sup>	3 fish.m <sup>-2</sup>	5 fish.m <sup>-2</sup>
Initial body weight (g)	55.40±3.40 <sup>a</sup>	55.41±2.90 <sup>a</sup>	56.07±1.06 <sup>a</sup>
Final body weight (g)	66.87±2.17 <sup>b</sup>	67.63±3.06 <sup>b</sup>	64.93±0.72 <sup>a</sup>
Growth in body weight (g)	11.47±1.29 <sup>b</sup>	12.22±1.09 <sup>b</sup>	8.86±0.57 <sup>a</sup>
Initial total length (cm)	20.03±0.42 <sup>a</sup>	20.16±0.40 <sup>a</sup>	19.87±0.09 <sup>a</sup>
Final total length (cm)	22.30±0.70 <sup>b</sup>	22.52±0.33 <sup>b</sup>	21.33±0.06 <sup>a</sup>
Growth in total length (cm)	2.27±0.32 <sup>b</sup>	2.37±0.12 <sup>b</sup>	1.46±0.10 <sup>a</sup>
Specific growth rate (% day <sup>-1</sup> )	0.45±0.07 <sup>b</sup>	0.48±0.05 <sup>b</sup>	0.34±0.03 <sup>a</sup>
Feed efficiency (%)	8.36±1.03 <sup>b</sup>	9.04±0.81 <sup>c</sup>	6.30±0.05 <sup>a</sup>
Survival rate (%)	100.00±0.00 <sup>a</sup>	100.00±0.00 <sup>a</sup>	93.33±11.55 <sup>a</sup>

Note:  $\text{mean} \pm \text{std}$  Value in the same column with different superscript (a, b, c) are significantly different (P < 0.05)

## CONCLUSION

In conclusion, the results of this study showed that stocking density affected the growth performance and feed efficiency of *N. notopterus* but not its survival. This study indicated that *N. notopterus* can be adapted to a net cage. This study provides a valuable reference for the domestication of *N. notopterus* in the future.

## ACKNOWLEDGEMENT

We are grateful to Mr. Nurrohman, a fisherman from the village of Tanjung Pering, who has been a great help in the technical work in the field.

## REFERENCES

Abdel-Aziz, M.F.A., Hassan, H.U., Yones, A.M., Abdel-Tawwab, Y.A., & Metwalli, A.A.A.T. (2021). Assessing the effect of different feeding frequencies combined with stocking density, initial weight, and dietary protein ratio on the growth performance of tilapia, catfish and carp. *Scientific African*, 12, e00806. <https://doi.org/10.1016/j.sciaf.2021.e00806>

Abdel-Latif, H.M.R., Ahmed, H.A., Shukry, M., Khafaga, A.F.,

Elkhayat, B.K., Abdel-Tawwab, M., & Abd-elaziz, R.A. (2023). Growth performance, physiological responses, and histoarchitectural changes in juvenile *Pangasianodon hypophthalmus* under different environmental salinities. *Fishes*, 8(6), 282. <https://doi.org/10.3390/fishes8060282>

Achakzai, W. M. (salah satu bentuk inkonsistensi, berjarak seperti ini? karena yang awal mepet semuanya), Saddozai, S., Baloch, W. A., Soomro, A. N., & Memon, N. (2015). Length-weight relationship and condition factor of *Notopterus notopterus* (Pallas, 1769) from Manchar Lake Sindh, Pakistan. *Sindh University Research Journal-SURJ (Science Series)*, 47(3), 515-518.

Aksungur, N., Aksungur, M., Akbulut, B., & Kutlu, İ. (2007). Effects of stocking density on growth performance, survival and food conversion ratio of Turbot (*Psetta maxima*) in the net cages on the southeastern coast of the Black Sea. *Turkish Journal of Fisheries and Aquatic Sciences*, 7(2), 147-152.

- Ammar, J. A., & Kamal, M. M. (2014). Keragaman ikan di Danau Cala, Kabupaten Musi Banyuasin Sumatera Selatan. *Depik*, 3(3), 216-220.  
<https://doi.org/10.13170/depik.3.3.2147>
- Andersson, M., Roques, J. A., Aliti, G. M., Ademar, K., Sundh, H., Sundell, K., Ericson, M. & Kettunen, P. (2022). Low holding densities increase stress response and aggression in zebrafish. *Biology*, 11(5), 725.  
<https://doi.org/10.3390/biology11050725>
- Barton, B. A. (2002). Stress in fishes: a diversity of responses with particular reference to changes in circulating corticosteroids. *Integrative and comparative biology*, 42(3), 517-525.  
<https://doi.org/10.1093/icb/42.3.517>
- Battisti, E. K., Rabaioli, A., Uczay, J., Sutili, F. J., & Lazzari, R. (2020). Effect of stocking density on growth, hematological and biochemical parameters and antioxidant status of silver catfish (*Rhamdia quelen*) cultured in a biofloc system. *Aquaculture*, 524, 735213.  
<https://doi.org/10.1016/j.aquaculture.2020.735213>
- Bœuf, G., & Payan, P. (2001). How should salinity influence fish growth?. *Comparative Biochemistry and Physiology Part C: Toxicology and Pharmacology*, 130(4), 411-423.  
[https://doi.org/10.1016/s1532-0456\(01\)00268-x](https://doi.org/10.1016/s1532-0456(01)00268-x)
- Budiardi, T., Gemawaty, N., & Wahjuningrum, D. (2007). Production of *Paracheirodon innesi* on different densities in recirculating system. *Jurnal Akuakultur Indonesia*, 6(2), 211-215.  
<https://journal.ipb.ac.id/index.php/jai/article/view/4030>
- Carbonara, P., Alfonso, S., Gai, F., Gasco, L., Palmegiano, G., Spedicato, M. T., Zupa, W., & Lembo, G. (2020). Moderate stocking density does not influence the behavioural and physiological responses of rainbow trout (*Oncorhynchus mykiss*) in organic aquaculture. *Aquaculture Research*, 51(7), 3007-3016.  
<https://doi.org/10.1111/are.14640>
- Conallin, J., Tun, N. N., Swe, A. M., Baumgartner, L. J., Lunn, Z., Mallen-Cooper, M., ... (???) & Thew, P. (2023). Using fish swimming ability to refine criteria for fishway construction in Myanmar. *Fisheries Research*, 262, 106680.  
<https://doi.org/10.1016/j.fishres.2023.106680>
- Gupta, A., Lal, K. K., Punia, P., Singh, R. K., Mohindra, V., Sah, R. S., Kumar, J., Luhariya, R.K., Dwivedi, A.K., Masih, P., Mishra, R.M., & Jena, J. K. (2013). Characterization of polymorphic microsatellite markers and genetic diversity in wild bronze featherback, *Notopterus notopterus* (Pallas, 1769). *Molecular Biology Reports*, 40, 6625-6631.  
<https://doi.org/10.1007/s11033-013-2776-z>
- Gustomi, A., Sulistiono, S., & Yonvitner, Y. (2016). Reproductive biology feather back (*Notopterus notopterus*, Pallas, 1769) in Simpbur Reservoir, Bangka Island. *Indonesian Journal of Agricultural Sciences*, 21(1), 56-62.  
<https://doi.org/10.18343/jipi.21.1.56>



- Jia, R., Wang, L., Hou, Y., Feng, W., Li, B., & Zhu, J. (2022). Effects of stocking density on the growth performance, physiological parameters, redox status and lipid metabolism of *Micropterus salmoides* in integrated rice–fish farming systems. *Antioxidants*, 11(7), 1215. <https://doi.org/10.3390/antiox11071215>
- Kozłowski, M., & Piotrowska, I. (2023). Effect of stocking density on growth, survival and cannibalism of juvenile pikeperch, *Sander lucioperca* (L.), in a recirculating aquaculture system. *Aquaculture International*, 1-9. <https://doi.org/10.1007/s10499-023-01339-6>
- Li, L., Shen, Y., Yang, W., Xu, X., & Li, J. (2021). Effect of different stocking densities on fish growth performance: A meta-analysis. *Aquaculture*, 544, 737152. <https://doi.org/10.1016/j.aquaculture.2021.737152>
- Liu, B., Jia, R., Zhao, K., Wang, G., Lei, J., & Huang, B. (2017). Stocking density effects on growth and stress response of juvenile turbot (*Scophthalmus maximus*) reared in land-based recirculating aquaculture system. *Acta Oceanologica Sinica*, 36, 31-38. <https://doi.org/10.1007/s13131-017-0976-4>
- Liu, Y., Liu, H., Wu, W., Yin, J., Mou, Z., & Hao, F. (2019). Effects of stocking density on growth performance and metabolism of juvenile Lenok (*Brachymystax lenok*). *Aquaculture*, 504, 107-113. <https://doi.org/10.1016/j.aquaculture.2019.01.058>
- Millán-Cubillo, A. F., Martos-Sitcha, J. A., Ruiz-Jarabo, I., Cárdenas, S., & Mancera, J. M. (2016). Low stocking density negatively affects growth, metabolism and stress pathways in juvenile specimens of meagre (*Argyrosomus regius*, Asso 1801). *Aquaculture*, 451, 87-92. <https://doi.org/10.1016/j.aquaculture.2015.08.034>
- Mohanty, D., & Samanta, L. (2016). Multivariate analysis of potential biomarkers of oxidative stress in *Notopterus notopterus* tissues from Mahanadi River as a function of concentration of heavy metals. *Chemosphere*, 155, 28-38. <https://doi.org/10.1016/j.chemosphere.2016.04.035>
- Mulyani, I., & Budijono, B. (2020). Morphometric and meristic analysis of Asian knifefish (*Notopterus notopterus*) in Sail River, Pekanbaru Riau Province. *Jurnal Ilmiah Biologi Eksperimen dan Keanekaragaman Hayati*, 7(2), 59-64.
- Muslim, M. (2023). Length-weight relationship and condition factor of *Notopterus notopterus* (Pallas, 1769) from East Pedamaran Floodplain, Ogan Komering Ilir, South Sumatra, Indonesia. *International Journal of Advanced Multidisciplinary Research and Studies*, 3(5), 1227-1231.
- Muslim, M., & Simanjuntak, W. J. (2023). Growth and survival of bronze featherback (*Notopterus notopterus*, Pallas 1769) reared on bucket. *Magna Scientia Advanced Research and Reviews*, 9(1), 101-105. <https://doi.org/10.30574/msarr.2023.9.1.0133>
- Muslim, M., & Syaifudin, M. (2022). Biodiversity of freshwater fish in Kelekar Floodplain Ogan Ilir Regency in Indonesia. *Journal of Tropical Biodiversity and*

- Biotechnology*, 7(1), 1-10.  
<https://doi.org/10.22146/jtbb.67494>
- Muslim, M., Pitriani, E., & Agustina, H. (2023). Growth and survival of bronze featherback (*Notopterus notopterus*) adapted on box container and aquarium. *GSC Advanced Research and Reviews*, 16(3), 133-137.  
<https://doi.org/10.30574/gscarr.2023.16.3.0368>
- Mustafa MG, Singha S, Islam M, Mallick N. Population dynamics of *Notopterus notopterus* (Pallas, 1769) from the Kaptai Reservoir of Bangladesh. *SAARC Journal of Agriculture* 2014; 12(2): 112–122.  
<https://doi.org/10.3329/sja.v12i2.21920>
- Mustafa, M. G., Singha, S., Islam, M. R., & Nayan Mallick, N. M. (2014). Population dynamics of *Notopterus notopterus* (Pallas, 1769) from the Kaptai Reservoir of Bangladesh. *SAARC Journal of Agriculture*, 12(2), 112–122.  
<https://doi.org/10.3329/sja.v12i2.21920>
- Naem, M., Salam, A., Gillani, Q., & Ishtiaq, A. (2010). Length-weight relationships of *Notopterus notopterus* and introduced *Oreochromis niloticus* from the Indus River, southern Punjab, Pakistan. *Journal of Applied Ichthyology*, 26(4), 620-620.  
<https://doi.org/10.1111/j.1439-0426.2010.01480.x>
- Rapita, R., Susiana, S., & Kurniawan, D. (2021). Food habits of belida fish (*Notopterus notopterus*, Pallas 1769) in Sei Gesek Reservoir, Bintan Regency, Riau Island, Indonesia. *IOP Conference Series. Earth Environmental Sci*, 919, 012003.  
<https://doi.org/10.1088/1755-1315/919/1/012003>
- Riche, M. A., Weirich, C. R., Wills, P. S., & Baptiste, R. M. (2013). Stocking density effects on production characteristics and body composition of market size cobia, *Rachycentron canadum*, reared in recirculating aquaculture systems. *Journal of the World Aquaculture Society*, 44(2), 259-266.  
<https://doi.org/10.1111/jwas.12023>
- Srivastava, S. M., Singh, S. P., & Pandey, A. K. (2012). Food and feeding habits of threatened *Notopterus notopterus* in Gomti river, Lucknow (India). *Journal of Experimental Zoology*, 15(2), 395–402.
- Sukendi, S., Thamrin, T., Putra, R. M., & Yulindra, A. (2020). Cultivation technology of bronze featherback (*Notopterus notopterus*, Pallas 1769) at different stocking densities and types of feed. *IOP Conference Series: Earth and Environmental Science*, 430(1), 012027.  
<https://doi.org/10.1088/1755-1315/430/1/012027>
- Ullah, K., Emmanuel, A., & Anjum, M. Z. (2018). Effect of stocking density on growth performance of Indus mahseer (*Tor macrotepis*). *International Journal of Fisheries and Aquatic Studies*, 6(3), 49-52.
- Winn, N. A., Sandi, P., Khaing, T., Nyunt, K. T., Kyaw, H. T., Sabai, M., & Aung, T. T. N. (2021). Length weight relationship of twelve freshwater fish species from Sunye Lake, Mandalay Region, Myanmar. *Greener Journal of Biological Sciences*, 11(2), 74-80.
- Xu, W., Lutz, C. G., Taylor, C. M., & Ortega, M. C. (2022). Improvement of fish growth and metabolism by oligosaccharide



prebiotic supplement. *Aquaculture Nutrition*, 28, 5715649.  
<https://doi.org/10.1155/2022/5715649>

Yang, Q., Guo, L., Liu, B. S., Guo, H. Y., Zhu, K. C., Zhang, N., ... & Zhang, D. C. (2020). Effects of stocking density on the growth performance, serum biochemistry, muscle composition and HSP70 gene expression of juvenile golden pompano *Trachinotus ovatus* (Linnaeus, 1758). *Aquaculture*, 518, 734841.  
<https://doi.org/10.1016/j.aquaculture.2019.734841>

Zlaugotne, B., Pubule, J., & Blumberga, D. (2022). Advantages and disadvantages of using more sustainable ingredients in fish feed. *Heliyon*, 8(9), e10527.  
<https://doi.org/10.1016/j.heliyon.2022.e10527>

# [JSAI] Editor Decision

2024-05-28 12:39 AM

Dear Muslim Muslim:

We have reached a decision regarding your submission to Jurnal Sumberdaya Akuatik Indopasifik, "Growth and survival of bronze featherback (*Notopterus notopterus*, Pallas 1769) reared on net cages".

Our decision is: Revisions Required

-----  
Reviewer A:

Recommendation: Revisions Required  
-----

**General Preview:**

# [JSAI] Editor Decision

2024-05-28 12:39 AM

Dear Muslim Muslim:

We have reached a decision regarding your submission to Jurnal Sumberdaya Akuatik Indopasifik, "Growth and survival of bronze featherback (*Notopterus notopterus*, Pallas 1769) reared on net cages".

Our decision is: Revisions Required

-----  
Reviewer A:

Recommendation: Revisions Required  
-----

**General Preview:**

# [JSAI] Editor Decision

2024-05-28 12:39 AM

Dear Muslim Muslim:

We have reached a decision regarding your submission to Jurnal Sumberdaya Akuatik Indopasifik, "Growth and survival of bronze featherback (*Notopterus notopterus*, Pallas 1769) reared on net cages".

Our decision is: Revisions Required

-----  
Reviewer A:

Recommendation: Revisions Required  
-----

**General Preview:**

## [JSAI] Editor Decision

2024-05-28 11:29 PM

Dear. Muslim Muslim:

We have reached a decision regarding your submission to Jurnal Sumberdaya Akuatik Indopasifik, "Growth and survival of bronze featherback (Notopterus notopterus, Pallas 1769) reared on net cages ".

Our decision is to: Accept Submission

Editor,

[Jurnal Sumberdaya Akuatik Indopasifik](#)

## [JSAI] Editor Decision

2024-05-28 11:29 PM

Dear. Muslim Muslim:

We have reached a decision regarding your submission to Jurnal Sumberdaya Akuatik Indopasifik, "Growth and survival of bronze featherback (Notopterus notopterus, Pallas 1769) reared on net cages ".

Our decision is to: Accept Submission

Editor,

[Jurnal Sumberdaya Akuatik Indopasifik](#)

## [JSAI] Editor Decision

2024-05-28 11:29 PM

Dear. Muslim Muslim:

We have reached a decision regarding your submission to Jurnal Sumberdaya Akuatik Indopasifik, "Growth and survival of bronze featherback (*Notopterus notopterus*, Pallas 1769) reared on net cages ".

Our decision is to: Accept Submission

Editor,

[Jurnal Sumberdaya Akuatik Indopasifik](#)



## [JSAI] Editor Decision

2024-06-02 01:07 AM

Dear Muslim Muslim:

The editing of your submission, "Growth and survival of bronze featherback (Notopterus notopterus, Pallas 1769) reared on net cages ," is complete. We are now sending it to production.

Submission URL: <https://ejournalfpikunipa.ac.id/index.php/JSAI/authorDashboard/submission/420>

Editor,

[Jurnal Sumberdaya Akuatik Indopasifik](https://ejournalfpikunipa.ac.id/index.php/JSAI/authorDashboard/submission/420)

## [JSAI] Editor Decision

2024-06-02 01:07 AM

Dear Muslim Muslim:

The editing of your submission, "Growth and survival of bronze featherback (Notopterus notopterus, Pallas 1769) reared on net cages ," is complete. We are now sending it to production.

Submission URL: <https://ejournalfpikunipa.ac.id/index.php/JSAI/authorDashboard/submission/420>

Editor,

[Jurnal Sumberdaya Akuatik Indopasifik](https://ejournalfpikunipa.ac.id/index.php/JSAI/authorDashboard/submission/420)

## [JSAI] Editor Decision

2024-06-02 01:07 AM

Dear Muslim Muslim:

The editing of your submission, "Growth and survival of bronze featherback (Notopterus notopterus, Pallas 1769) reared on net cages ," is complete. We are now sending it to production.

Submission URL: <https://ejournalfpikunipa.ac.id/index.php/JSAI/authorDashboard/submission/420>

Editor,

[Jurnal Sumberdaya Akuatik Indopasifik](https://ejournalfpikunipa.ac.id/index.php/JSAI/authorDashboard/submission/420)