


Effectiveness of Biolarvicides of Imperata cylindrica, Saccharum spotaneum and Andropogon aciculatus on Aedes aegypti larval Mortality and Egg-laying Ability in Adults

By Chandra Irsan

Research Article



Effectiveness of Biolarvicides of *Imperata cylindrica*, *Saccharum spontaneum* and *Andropogon aciculatus* on *Aedes aegypti* larval Mortality and Egg-laying Ability in Adults

Heru Listiono¹, Nurhayati Damiri^{2*}, Totong Kamaluddin³, Chandra Irsan², Supli Effendi Rahim⁴

¹Department of Environmental Science, Graduate Program, Universitas Sriwijaya, Palembang 30139, South Sumatra, Indonesia

²Department of Plant Protection, Faculty of Agriculture, Universitas Sriwijaya, Ogan Ilir, South Sumatra Indonesia

³Department of Farmakology, Faculty of Medicine, Universitas Sriwijaya, Ogan Ilir, South Sumatra Indonesia

⁴Master of Agricultural Science, Postgraduate Studi Program, Universitas Muhammadiyah Palembang, South Sumatra, Indonesia

ARTICLE INFO

Article history:

Received January 22, 2024

Received in revised form September 22, 2024

Accepted January 13, 2025

KEYWORDS:

Biopesticides,
Aedes aegypti,
larval mortality,
egg-laying

ABSTRACT

Vector-borne disease such as Dengue Hemorrhagic (DHF), transmitted by *Aedes aegypti* and *Aedes albopictus*, remain a significant public health concern in Indonesia. Controlling these disease often involves insecticides; however, the negative impact of chemical insecticides have prompted interest in organic alternatives derived from plants. Certain weeds, including cogon grass (*Imperata cylindrica*), wild sugarcane (*Saccharum spontaneum*), and needle grass (*Andropogon aciculatus*), have shown potential as botanical insecticides. Research findings showed that weed root extracts significantly affect larval mortality rate of *Ae. Aegypti*. At 1000 ppm, larval mortality was significantly higher compared to 100 ppm and the control, while treatments of 1 ppm and 10 ppm showed similar results to the control. Probit analysis revealed that *I. cylindrica* root extract achieved an LC50 of 974.99 ppm within 24 hours, indicating it could kill 50% of *Ae. Aegypti* larvae. Within 48 hours, the LC50 dropped to 889.20 ppm. Toxicity tests further revealed significant differences in *Ae. Aegypti* egg-laying abilities when treated with extracts. Analysis of variance yielded p-values of 0.000 for egg hatching within 72 and 96 hours, highlighting significant differences across samples. These findings suggest the extracts influence mosquito reproduction, warranting further studies to assess the quality of egg hatched from larvae exposed to these treatments. The potential of botanical insecticides derived from weeds represents a promising step toward sustainable mosquito control in the fight against vector-borne diseases.



Copyright (c) 2025@ author(s).

1. Introduction

Dengue, a mosquito-borne virus, is the leading cause of arthropod-borne viral disease globally, posing a significant health concern. Since early 2023, continuous transmission and an unexpected surge in dengue cases have led to nearly record levels, with over five million cases and more than 5,000 dengue-related deaths reported in over 80 countries and territories across five WHO regions: Africa, the

Americas, Southeast Asia, the Western Pacific, and the Eastern Mediterranean (World Health Organization 2023).

DHF is a disease transmitted by mosquitoes as a vector. The two invasive mosquito species in Indonesia are *Aedes aegypti* and *Ae. Albopictus* is not just a significant nuisance to humans but also the main carrier of various foreign pathogens, including the viruses responsible for dengue, zika, and chikungunya. It's crucial to control these diseases effectively. Are a significant nuisance to humans and serve as the main carriers of various foreign pathogens, including the viruses responsible for dengue, Zika, and chikungunya.

* Corresponding Author

E-mail Address: nurhayati@fp.unsri.ac.id

(Rhida *et al.* 2023). Dengue Hemorrhagic Fever (DHF) has been increasing and spreading throughout Indonesia since its discovery in 1968 (Giroth *et al.* 2019).

Mosquito control is a worldwide public health practice worldwide in tropical and sub-tropical areas (Listiono *et al.* 2023; Duval *et al.* 2023). Mosquitoes are the most dangerous disease vector belonging to the arthropod family. They are significant for public health because of their blood-feeding habit (Siam *et al.* 2022; Nebbak *et al.* 2022; Attaullah *et al.* 2023). Mosquitoes exhibit high species diversity, with approximately 3,490 officially recorded (Kirik *et al.* 2021). Despite ongoing control efforts, the mosquito population and transmitted diseases rapidly expand globally, posing significant health problem (Lim *et al.* 2023). Disease management strategies to reduce the burden of mosquito-borne diseases depend on vector or mosquito control (Ramanti *et al.* 2023). The unregulated and excessive use of insecticides and pesticides to control pests has led to mosquito resistance to these larvicides (Raul *et al.* 2021).

Breaking the vector transmission chain can be achieved during the larval phase. One way to control DHF during the larval stage is using chemicals known as insecticides. However, the use of chemical insecticides has several side effects. The long-term use of synthetic insecticides has made mosquitoes resistant, necessitating new vector control methods. The synthetic chemical insecticides currently in use have some disadvantages. Factors such as vector resistance and toxicity to humans and non-target organisms increase the interest in exploring new control alternatives (Pavela 2016; Billi 2018).

There has been an increase in the use of plants as an alternative to synthetic insecticides due to their broad insecticidal properties, biodegradability, and adaptability to ecological conditions (Demirak and Canpolak 2022; Lim *et al.* 2023). One effort to mitigate these negative impacts is the utilization of organic insecticides derived from natural compounds found in plants (Giroth *et al.* 2019; Pereira *et al.* 2024). Several studies regarding the potential of plant extract in controlling mosquitoes, including the fern *Actinidia chinensis* radiata, were tested with novel mosquitocidal activity against larvae of *Ae. aegypti* and *Anopheles stephensi* (Kamaraj *et al.* 2018). Seed oil extract of *Acacia nilotica* possessed robust larvicidal action against major mosquito vectors (Vivekanandhan *et al.* 2018). Plant essential oils

were also reported to have mosquitocidal potential. The crude volatile oil (CVO) from Piper betle leaves possessed significant larvicidal, ovipositional and repellency effects against *Ae. aegypti* (Vasanthasrinivasan *et al.* 2018; Vivekanandhan *et al.* 2018; Chellappandian *et al.* 2019).

Weed plants contain various active substances, including tannins, phenolics, terpenoids, alkaloids, saponins, and flavonoids. These phytochemical compounds can trigger various reactions in the mosquito larvae, disrupting their growth and development and ultimately leading to larval death. Flavonoids function as potent respiratory inhibitors, causing larval death; saponin compounds can corrode the mucous membrane of the larval digestive tract; tannin compounds act as stomach poisons for mosquito larvae, disrupting their nutrition; and steroid compounds can inhibit the molting process in larvae, causing them to become unconscious (Fitriyani *et al.* 2023). The presence of these bioactive compounds suggests that cogon grass (*Imperata cylindrica*), wild sugarcane (*Saccharum spontaneum*), and needle grass (*Andropogon aciculatus*) can be used as natural larvicides.

Imperata cylindrica is among the ten worst weeds in tropical and subtropical regions. Despite having many benefits, the issues related to its invasive nature outweigh the positive aspects (Rusdy *et al.* 2020). *Imperata cylindrica* contains organic phenolic compounds that are toxic to *Aedes aegypti* mosquito larvae, but the concentration used in experiments is not high enough to kill the larvae. *Imperata cylindrica* can paralyze larvae within the first twenty-four hours, with larval movement observed a few hours later (Calago *et al.* 2015). Furthermore, chemical component analysis of wild sugarcane reveals that the resulting bio-oil is dominated by acetic acid, phenol, and 1-hydroxy-2-propanone (Wibowo *et al.* 2015).

The objective of this research was to determine the effectiveness of bio larvicides derived from weed plants, specifically cogon grass (*I. cylindrica*), wild sugarcane (*S. spontaneum*), and needle grass (*A. aciculatus*) on the mortality of *Ae. aegypti* mosquito larvae.

2. Materials and Methods

This study was pure experimental research with a post-test-only control group design. The purpose of this study was to investigate the effectiveness of secondary

metabolites from weed plants, namely cogon grass (*I. cylindrica*), wild sugarcane (*S. spontaneum*), and needle grass (*A. aciculatus*) on the mortality of *Aedes aegypti* mosquito larvae. The research was conducted at the Health Research and Development Center, Litbangkes Entomology Laboratory Baturaja, South Sumatra, from January to August 2023.

Preparation of larvae. Two thousand mosquito eggs of *Ae. aegypti* were obtained from the Health Research and Development Center in Baturaja, South Sumatra, and colonized until they became larvae. After the larvae hatched and reached the third instar, 500 third-instar larvae were taken to be treated with weed extract.

Preparation of weed extract. Weeds of *I. cylindrica*, *S. spontaneum*, and *A. aciculatus* collected from agricultural and plantation areas were cut into small pieces and dried in the sun. Once completely dry, the weed plants were ground into a fine powder using a blender and then extracted using the maceration method with 96% ethanol, followed by evaporation using a rotary evaporator. The weed extract obtained is then diluted according to the treatment, namely one ppm, ten ppm, 100 ppm, and 1000 ppm. The weed extraction method is carried out separately between the three weeds. This concentration is based on the Guidelines for Laboratory and Field Testing of Mosquito Larvicides (WHO/CDS/WHOPES/GCDPP/2005).

Application of weed extract. This study involved five treatment groups, including one control and four experimental groups, with four repetitions based on the Fecundity formula.

The negative control group was treated with water without the addition of weed extract. In contrast, the experimental groups were exposed to four different concentrations of weed extract, one ppm, ten ppm, 100 ppm, and 1000 ppm. Six different plant parts were used as samples, including leaves and roots of cogon grass (*I. cylindrica*), wild sugarcane (*S. spontaneum*), and needle grass (*A. aciculatus*).

They are testing the effectiveness of weed extract on *Ae. aegypti* mosquito larvae was carried out using a 100 ml plastic cup, where the extract solution was dropped into the plastic cup according to the respective concentration. One plastic cup contains 25 larvae x 5 concentrations and four repetitions = 500 intra three larvae, which have been selected, and healthy larvae are taken after the larvae are put into the plastic cup. Next, the number of dead larvae was counted during the 24-hour and 48-hour test periods.

To test the *Ae. aegypti* mosquito's ability based on the number of hatched eggs, third-instar larvae that had been exposed to the extract treatment and survived were bred to become adult mosquitoes. The research was done by transferring them into mosquito cages. After the mosquitoes had matured in each sample type, specifically for the 1000 ppm treatment, two healthy male and female mosquitoes were taken and bred in one mosquito cage for every kind of sample. They received a blood meal from a shaved-back male guinea pig during breeding. After the mosquitoes had mated, fed on blood, and laid eggs, the number of eggs was counted from 72 hours to 96 hours.

One-way ANOVA or Kruskal-Wallis analysis was used to analyze the data obtained in the research. If the One-Way ANOVA or Kruskal-Wallis test results were significant, a post-hoc analysis would be conducted to determine which groups were significant. For One-Way ANOVA, the post-hoc analysis used was Bonferroni, and for the Kruskal-Wallis test was Mann-Whitney (Hastono 2018). The LC50 was determined by using the probit test (Ekawati *et al.* 2017).

Analysis of secondary metabolite content from the three types of weeds was carried out using GS-MS. Weed extracts of *I. cylindrica*, *S. spontaneum*, and *A. aciculatus* using maceration or soaking method. Then, it was put in a micro tube containing 0.5 g of powder and 1.5 ml of ethanol solvent, vortexed for 1 minute, and centrifuged for 3 minutes at a speed of 90 rpm. The supernatant formed was continued for GC-MS testing. The time was 60 minutes with an injector temperature of 260°C, detector 250°C, and column 325°C. The carrier gas used is a helium gas carrier with a constant flow rate of 1 ml per minute. The identification process using the GC-MS tool produces several bioactive which can be seen from the peaks of the chromatogram as identification data from chromatography and mass spectrometry (MS) results seen from the mass spectrum with each molecular weight of the bioactive compound.

3. Results

3.1. Effect of Extract on The Death of *Ae. aegypti* Larvae

The average larval mortality in the thatch root treatment significantly differs in each concentration. The results show that that of 1000 ppm significantly differs from the concentration of 100 ppm and the control. However, the one ppm and ten ppm treatments

were not significantly different from the control (Table 1).

Based on the results of this research, it is clear that the *I. cylindrical* root extract has quite a practical effect on killing *Ae. Aegypti*. The leaf extract of *I. cylindrical*, *S. spontaneum*, and *A. aciculatus* was compared with the roots extract of *S. spontaneum* and *A. aciculatus*.

Based on the probit analysis in Table 1, it can be seen that extracts from *I. cylindrical* leaves, *S. Spontaneum*, and *A. aciculatus* were not toxic to *Ae. Aegypti* larvae within 24 hours, as well as root extracts of *S. spontaneum* and *A. aciculatus* were not toxic to *Ae. aegypti* within 24 hours after application. In contrast, *I. cylindrical* root extract was toxic to *Ae. aegypti* larvae within 24 hours and were able to kill 50% of the *Ae. aegypti* larvae are equivalent to 50 out of 100 larvae.

The average larval mortality after 48 hours of applying the extract significantly differs at each

concentration. The results show that the concentration of 1000 ppm significantly differs from that of 100 ppm and the control. However, the one-ppm and ten-ppm treatments were not significantly different from the control (Table 2).

Based on the probit analysis, *A. aciculatus* leaf extract and *S. spontaneum* leaf extract were non-toxic to *Ae. aegypti* larvae at 48 hours. In contrast, *I. cylindrical* roots extract was toxic to *Ae. aegypti* larvae within 48 hours and could kill 50% of *Ae. aegypti* larvae, which is equivalent to 50 larvae out of 100. For *S. spontaneum* roots, the result showed LC 50>1000 ppm (141579.4), and for *A. aciculatus* roots, the result indicated LC 50>1000 ppm (141579.4), suggesting that both root extracts were non-toxic to *Ae. aegypti* larvae at 48 hours. Symptoms of the effect of the three grass extracts on *Ae. aegypti* larvae were not different from each other (Figure 1)

Table 1. Effect of extract on the death of *Ae. aegypti* larvae in the 24 hour after application

Concentration	Differences in average larval mortality					
	Leaf <i>I. cylindrical</i>	Leaf <i>S. spontaneum</i>	Leaf <i>A. aciculatus</i>	Root <i>I. cylindrical</i>	Root <i>S. spontaneum</i>	Root <i>A. aciculatus</i>
48						
Control	0.00	0.00	0.00	0.00c	0.00	0.00
1 ppm	0.00	0.00	0.00	0.00c	0.00	0.00
10 ppm	0.00	0.00	0.00	0.00c	0.00	0.00
100 ppm	0.25	0.00	0.00	2.00b	0.00	0.00
1000 ppm	6.50	1.25	0.50	13.25a	0.75	1.25
F Value	1.26ns	2.09ns	3.00ns	63.72*	2.65ns	2.88ns
P Value	0.33	0.13	0.06	3.12×10^{-9}	0.07	0.06
BNJ 5%	-	-	-	0.71	-	-
LC 50	3681.2 ppm	141579.4 ppm	141579.4 ppm	974,989 ppm	141579.4 ppm	141579.4 ppm

* significantly different; values in columns followed by the same letter are not significantly different at $P < 0.05$ according to the BNJ test. Original data were transformed using square root transformation

Table 2. Effect of extract on the death of *Ae. aegypti* larvae in the 48 hour after application

Concentration	Differences in average larval mortality					
	Leaf <i>I. cylindrical</i>	Leaf <i>S. spontaneum</i>	Leaf <i>A. aciculatus</i>	Root <i>I. cylindrical</i>	Root <i>S. spontaneum</i>	Root <i>A. aciculatus</i>
65						
Control	0.00	0.00b	0.00	0.00c	0.00	0.00
1 ppm	0.00	0.00b	0.00	0.00c	0.00	0.00
10 ppm	0.00	0.00b	0.00	0.00c	0.00	0.00
100 ppm	0.25	0.00b	0.00	2.50b	0.00	0.00
1000 ppm	7.50	1.50a	0.75	15.00a	0.75	1.25
F Value	2.68ns	4.34*	2.65ns	129.20*	2.65ns	2.88ns
P Value	0.07	0.01	0.07	1.97×10^{-11}	0.07	0.06
BNJ 5%	-	0.57	-	0.54	-	-
LC 50	3169.5 ppm	141579.4 ppm	141579.4 ppm	889,201 ppm	141579.4 ppm	141579.4 ppm

* significantly different; values in columns followed by the same letter are not significantly different at $P < 0.05$ according to the BNJ test. Original data were transformed using square root transformation

3.2. Identification of Molecular Formulas using The Gas Chromatography-Mass Spectrometry (GC-MS) Method

The result of the analysis of secondary metabolites compounds present in the three types of grass used can be seen in Table 3.

Table 3 shows that the chemical compounds in weed leaves that are thought to be used as bioinsecticides are campesterol, sitosterol, -Phenylalanyl-glycine,

and stigmasterol. In *I. cylindrica* leaves, campesterol appeared at 21.64 minutes with an area size of 1.22%, 21.79 minutes with an area size of 2.87%, 21.95 minutes with an area size of 1.93% and sitosterol appeared at 26.40 minutes with an area size of 28.18%. In *S. spontaneum* leaves, there is 1-Phenylalanyl-glycine appearing at 9.98 minutes with an area size of 0.85%, and in *A. aciculatus* leaves, there is








Figure 1. Effect of extracts of *I. cylindrica*, *S. spontaneum*, *A. aciculatus* on *Ae. Aegypti* larvae. (A) Larvae from the extract test whose body shape is incomplete. (B) Larvae from the extract test whose body parts are broken down. (C) Larvae as a result of the extract test whose insides are broken down. (D) Healthy larvae

Table 3. GCMS test results on *I. cylindrica*, *S. spontaneum*, *A. aciculatus* leaves

Weed leaves	Compound name	Chemical compound formulasi	Information
<i>Imperata cylindrica</i>	Campesterol		Campesterol appeared at 21.64 minutes with an area size of 1.22%, 21.79 minutes with an area size of 2.87%, 21.95 minutes with an area size of 1.93%
	Sitosterol		Sitosterol appeared at 26.40 minutes with an area size of 28.18%
<i>Saccarum spontaneum</i>	1-Phenylalanyl-glycine		1-Phenylalanyl-glycine appears at 9.98 minutes with an area of 0.85%
<i>Andropogon aciculatus</i>	Stigmasterol		Stigmasterol appeared at 23.86 minutes with an area size of 2.44%

Stigmasterol appearing at 23.86 minutes with an area size of 2.44%.

It can be seen in Table 4 that chemical compounds in weed roots that are thought to be able to be used as bioinsecticides are Phenol,2,4-bis, Phenol,2,6-bis, Phenol,3,5-bis, and Sitosterol. In the roots of *I. cylindrica* Phenol, 2,4-bis, appears at 13.74 minutes with an area size of 0.58%. In the roots of *S. spontaneum*, there is Phenol, 2,6-bis, appearing at 13.74 minutes with an area size of 0.58% and Sitosterol appearing at 26.40 minutes with an area size of 1.56% as well as in the roots of *A. aciculatus* There is Phenol,3,5-bis appearing at 13.74 minutes with an area size of 1.28%.

Effect of weed extract toxicity test on mosquito larvae in their ability to lay eggs.

The results of observations of the effect of plant extracts on the egg-laying ability of *Ae aegypti* showed that the longer the application period, the more eggs were hatched. The number of eggs that hatched at a concentration of 1000 ppm in 72 hours showed the average number of eggs that hatched: 75 *I. cylindrica* leaves, 77.6 *S. spontaneum* leaves, 80.3 *A. aciculatus* leaves. *I. cylindrica* roots 9.6, *S. spontaneum* roots 27.3, and *A. aciculatus* roots 13.3. For 96 hours, the average number of eggs hatched was as follows: *I. cylindrica* leaves 76.33, *S. spontaneum* leaves 90, *A. aciculatus* leaves 83. *I. cylindrica* roots 14, *S. spontaneum* roots 19.3, *A. aciculatus* roots 17.6 (Figure 2).





Analysis of the effect of weed extract on *Ae aegypti* mosquito larvae at 72 hours and 96 showed a very


significant effect on the egg-laying ability of adult mosquitoes-the results of further tests on the effect of 74ed extract on the ability to lay of *Ae. Aegypti* egg can be seen in Table 5 and 6.

Table 5 shows significant differences in the number of eggs produced by mosquitoes subjected to various treatments. Notably, differences were observed between mosquitoes treated with *I. cylindrica* leaf extract and those treated with *I. cylindrica* root extract ($p = 0.009$), *I. cylindrica* leaf extract and *A. aciculatus* root extract ($p = 0.015$), and *I. cylindrica* leaf extract and *S. spontaneum* root extract ($p = 0.014$). Additionally, there were differences between mosquitoes treated with *S. spontaneum* leaf extract and *I. cylindrica* root extract ($p = 0.007$), *S. spontaneum* leaf extract and *S. spontaneum* root extract ($p = 0.011$), *S. spontaneum* leaf extract and *A. aciculatus* root extract ($p = 0.010$), *A. aciculatus* leaf extract and *I. cylindrica* root extract ($p = 0.005$), *A. aciculatus* leaf extract and *S. spontaneum* root extract ($p = 0.008$), and *A. aciculatus* leaf extract and *A. aciculatus* root extract ($p = 0.007$).

Table 6 presents Bonferroni's post-hoc test result, revealing significant differences in the number of eggs mosquitoes produce under various treatments. There were notable differences between mosquitoes treated with *I. cylindrica* leaf extract and those treated with *I. cylindrica* root extract ($p = 0.001$), *I. cylindrica* leaf extract and *A. aciculatus* root extract ($p = 0.003$), *I. cylindrica* leaf extract and *S. spontaneum* root extract ($p = 0.002$), *S. spontaneum* leaf extract and *I. cylindrica* root extract ($p = 0.000$), *S. spontaneum* leaf extract and *S. spontaneum* root extract ($p = 0.000$),

Table 4. GCMS test results on roots of *Imperata cylindrica*, *Saccarum spontaneum*, *Andropogon aciculatus*

Weed leaves	Compound name	Chemical compound formulasi	Information
<i>Imperata cylindrica</i>	Phenol,2,4-bis		Phenol,2,4-bis, appears at 13.74 minutes with an area size of 0.58%
<i>Saccarum spontaneum</i>	Phenol,2,6-bis		Phenol,2,6-bis, appears at 13.74 minutes with an area size of 1.17%
	Sitosterol		Sitosterol appears at 26.40 minutes with an area size of 1.56%
<i>Andropogon aciculatus</i>	Phenol,3,5-bis		Phenol,3,5- appears at minute 13.74 with an area size of 1.28%



27
Figure 2. Effect of weed extract on *Ae. aegypti* eggs hatching

27
Table 5 . Effect of weed extract on *Ae. aegypti* larvae and the ability of their adult mosquitoes to lay eggs at 72 hours after application

(I) Jenis sample	(J) Jenis sample	Mean difference (I-J)	Std. error	Sig.	95 % Confidence interval	
					lower bound	upper bound
Leaf <i>I. cylindrica</i>	Leaf <i>S. spontaneum</i>	-2,667	14,184	1.000	-54.42	49.09
	Leaf <i>A. aciculatus</i>	-5,333	14,184	1.000	-57.09	46.42
	Root <i>I. cylindrica</i>	65,000*	14,184	0.009	13.58	117.09
	Root <i>S. spontaneum</i>	61,000*	14,184	0.015	9.58	113.09
	Root <i>A. aciculatus</i>	61,667*	14,184	0.014	9.91	113.42
Leaf <i>S. spontaneum</i>	Leaf <i>I. cylindrica</i>	2,667	14,184	1.000	-49.09	54.42
	Leaf <i>A. aciculatus</i>	-2,667	14,184	1.000	-54.42	49.09
	Root <i>I. cylindrica</i>	68,000*	14,184	0.007	16.24	119.76
	Root <i>S. spontaneum</i>	64,000*	14,184	0.011	12.24	115.76
	Root <i>A. aciculatus</i>	64,333*	14,184	0.010	12.58	116.09
Leaf <i>A. aciculatus</i>	Leaf <i>I. cylindrica</i>	5,333	14,184	1.000	-46.42	57.09
	Leaf <i>S. spontaneum</i>	2,667	14,184	1.000	-49.09	54.42
	Root <i>I. cylindrica</i>	70,667*	14,184	0.005	18.91	122.42
	Root <i>S. spontaneum</i>	66,667*	14,184	0.008	14.91	118.42
	Root <i>A. aciculatus</i>	67,000*	14,184	0.007	15.24	118.76
Root <i>I. cylindrica</i>	Leaf <i>I. cylindrica</i>	-65,333*	14,184	0.009	-113.09	-9.58
	Leaf <i>S. spontaneum</i>	-68,000*	14,184	0.007	-115.76	-12.24
	Leaf <i>A. aciculatus</i>	-70,667*	14,184	0.005	-118.42	-14.91
	Root <i>S. spontaneum</i>	-4,000	14,184	1.000	-47.76	55.76
	Root <i>A. aciculatus</i>	-3,667	14,184	1.000	-51.42	52.09
Root <i>S. spontaneum</i>	Leaf <i>I. cylindrica</i>	-61,333*	14,184	0.015	-113.09	-9.58
	Leaf <i>S. spontaneum</i>	-64,000*	14,184	0.11	-115.76	-12.24
	Leaf <i>A. aciculatus</i>	-66,667*	14,184	0.008	-118.42	-14.91
	Root <i>I. cylindrica</i>	4,000	14,184	1.000	-47.76	55.76
	Root <i>A. aciculatus</i>	0,333	14,184	1.000	-51.42	52.09
Root <i>A. aciculatus</i>	Leaf <i>I. cylindrica</i>	-61,667*	14,184	0.014	-113.42	-9.91
	Leaf <i>S. spontaneum</i>	-64,333*	14,184	0.010	-116.09	-12.58
	Leaf <i>A. aciculatus</i>	-67,000*	14,184	0.007	-118.76	-15.24
	Root <i>I. cylindrica</i>	3,667	14,184	1.000	-48.09	55.42
	Root <i>S. spontaneum</i>	-0,333	14,184	1.000	-52.09	51.42

Table 6. Effect of weed extract on *Ae. aegypti* larvae and the ability of their adult mosquitoes to lay eggs at 72 hours after application

(I) Jenis sample	(J) Jenis sample	Mean difference (I-J)	Std. error	Sig.	95 % Confidence interval	
					lower bound	upper bound
Leaf <i>I. cylindrica</i>	Leaf <i>S. spontaneum</i>	-13,333	10,919	1.000	-53.18	26.51
	Leaf <i>A. aciculatus</i>	-6,333	10,919	1.000	-46.18	33.51
	Root <i>I. cylindrica</i>	60,000*	10,919	0.001	22.82	102.51
	Root <i>S. spontaneum</i>	57,333*	10,919	0.003	17.49	97.18
	Root <i>A. aciculatus</i>	59,000*	10,919	0.002	19.16	98.84
Leaf <i>S. spontaneum</i>	Leaf <i>I. cylindrica</i>	13,333	10,919	1.000	-26.51	53.18
	Leaf <i>A. aciculatus</i>	7,000	10,919	1.000	-32.84	46.84
	Root <i>I. cylindrica</i>	76,000*	10,919	0.000	36.16	115.84
	Root <i>S. spontaneum</i>	70,667*	10,919	0.000	30.82	110.51
	Root <i>A. aciculatus</i>	72,333*	10,919	0.000	32.49	112.18
Leaf <i>A. aciculatus</i>	Leaf <i>I. cylindrica</i>	6,333	10,919	1.000	-33.51	46.18
	Leaf <i>S. spontaneum</i>	-7,000	10,919	1.000	-46.84	32.84
	Root <i>I. cylindrica</i>	69,000*	10,919	0.001	29.16	108.84
	Root <i>S. spontaneum</i>	63,667*	10,919	0.001	23.52	103.51
	Root <i>A. aciculatus</i>	65,333*	10,919	0.001	25.49	105.18
Root <i>I. cylindrica</i>	Leaf <i>I. cylindrica</i>	-62,667*	10,919	0.001	-102.51	-22.82
	Leaf <i>S. spontaneum</i>	-76,000*	10,919	0.000	-115.84	-36.16
	Leaf <i>A. aciculatus</i>	-69,000*	10,919	0.001	-108.84	-29.16
	Root <i>S. spontaneum</i>	-5,333	10,919	1.000	-45.18	34.51
	Root <i>A. aciculatus</i>	-3,667	10,919	1.000	-43.51	36.18
Root <i>S. spontaneum</i>	Leaf <i>I. cylindrica</i>	-57,333*	10,919	0.003	-97.18	-17.49
	Leaf <i>S. spontaneum</i>	-70,667*	10,919	0.000	-110.51	-30.82
	Leaf <i>A. aciculatus</i>	-63,667*	10,919	0.001	-103.51	-23.82
	Root <i>I. cylindrica</i>	5,333	10,919	1.000	-34.51	45.18
	Root <i>A. aciculatus</i>	1,667	10,919	1.000	-38.18	41.51
Root <i>A. aciculatus</i>	Leaf <i>I. cylindrica</i>	-59,000*	10,919	0.002	-98.84	-19.16
	Leaf <i>S. spontaneum</i>	-72,333*	10,919	0.000	-112.18	-32.49
	Leaf <i>A. aciculatus</i>	-65,333*	10,919	0.001	-105.18	-25.49
	Root <i>I. cylindrica</i>	3,667	10,919	1.000	-36.18	43.51
	Root <i>S. spontaneum</i>	-1,667	10,919	1.000	-41.51	38.18

S. spontaneum leaf extract and *A. aciculatus* root extract ($p = 0.000$), *A. aciculatus* leaf extract and *A. aciculatus* root extract ($p = 0.001$), *A. aciculatus* leaf extract and *S. spontaneum* root extract ($p = 0.001$), and *A. aciculatus* leaf extract and *A. aciculatus* root extract ($p = 0.001$).

4. Discussion

4.1. Effect of Extract on The Death of *Ae. aegypti* Larvae

The average larval mortality in the thatch root treatment significantly differs in each concentration. The results show that the concentration of 1000 ppm significantly differs from the concentration of 100 ppm and the control. However, the 1 ppm and 10 ppm treatments were not significantly different from the control. These results showed two sample types with significant differences in the average among the 5

concentration series, namely the *I. cylindrica* root and *S. spontaneum* root extracts. Furthermore, based on the probit analysis, for the *I. cylindrica* root extract, $LC_{50} < 1000$ ppm (5874.989), indicating that *I. cylindrica* root extract was toxic to *Ae. aegypti* larvae within 24 hours. At a concentration of 974.989 ppm, the *I. cylindrica* root extract could kill 50% of *Ae. aegypti* larvae, which is equivalent to 50 eggs out of 100 larvae.

The potency of various concentrations of *I. cylindrica* root extract in killing *Aedes* sp. dengue mosquito larvae. There was an increase in the average mortality. The higher the concentration and treatment time applied, the higher the average mortality of *Aedes* sp. dengue mosquito larvae. (Antou *et al.* 2023). In Figure 1, it can be seen that an increase in concentration indicates a toxic compound that enters the body of the test larvae, resulting in disrupted larval life and causing mortality. When treated with concentrations of stolon extract from weeds, the death of the larvae caused a

change in activity, where when the larvae were about to die, there was movement up and down on the surface. The death of the larvae if the larvae do not move and do not respond, and float above the surface indicates that the larvae are dead.

Plants are rich in biologically active substances with the potential to control *Ae. aegypti* are considered attractive alternatives to traditional chemical insecticides (Uangmoon *et al.* 2018; Azevedo *et al.* 2019). The phytochemical analysis of the *I. cylindrica* root extract revealed the presence of flavonoid compounds, known for their phenolic content (Tungmunthum *et al.* 2018). Flavonoids have been shown to act as potent respiratory inhibitors that can lead to larval mortality (Fitriyani *et al.* 2022). Several chemical factors can influence the extraction results, including internal factors like the type of active compounds in the material, qualitative composition of active compounds, quantitative composition of active compounds, and average total content of active compounds. External factors, on the other hand, include the extraction method, the type of extraction equipment, the characteristics of the material, and the solvents used in the extraction process (Zhang *et al.* 2023). Another factor is believed to affect the LC50 of the *I. cylindrica* root extract on *Ae. aegypti* larval mortality is the difference in flavonoid content between *I. cylindrica* root, *S. spontaneum* root, and *A. aciculatus* root. *I. cylindrica* root is thought to contain more flavonoids than wild sugarcane and *A. aciculatus* roots. *I. cylindrica* roots are obtained from terrestrial plantation areas, while *S. spontaneum* and *I. cylindrica* roots are collected from aquatic areas. Environmental factors, such as temperature stress (Utomo *et al.* 2020), can also affect the production of secondary metabolites. In this context, terrestrial plantation areas have higher temperatures than aquatic areas.

Based on the observations of *Ae. aegypti* mosquito larvae, signs of mortality were evident, with the larvae displaying disrupted body integrity and visible deterioration in their internal structure. The researchers continued this observation until the fifth day, indicating that most larvae had not yet transformed into pupae or adult mosquitoes. Additionally, the larvae exhibited reduced activity. These observations were believed to result from inhibiting larval growth by secondary metabolites found in the *I. cylindrica* root extract. The mortality of *Ae. aegypti* larvae, particularly at a concentration of 1000 ppm of *I. cylindrica* root extract, was suspected to be caused by the active compounds coming into direct contact with the larvae. In the ethanol

extract of *I. cylindrica* roots, active compounds such as flavonoids, alkaloids, and tannins are present (Maharani *et al.* 2018). Flavonoids, secondary metabolites derived from plants and other microorganisms, influence insect development and physiology. These compounds can act as larvicides against *Ae. aegypti*. For instance, several plants' flavonoid extracts or purified flavonoids have shown larvicidal activity against *Ae. aegypti* and other vector mosquitoes (Pessoa *et al.* 2018; Pavela *et al.* 2019; Al-Massarani *et al.* 2019; Swargiary *et al.* 2019).

4.2. Effects of Weed Extract Toxicity on The Ability of *Ae. Aegypti* Larvae to Lay Eggs

The results of the *Ae. aegypti* larva breeding test regarding their ability to lay eggs at a concentration of 1000 ppm within 72 hours showed variations in the number of eggs hatched among the different weed extract samples. The average number of eggs hatched for each sample was as follows: *I. cylindrica* leaves 75, *S. spontaneum* leaves 77.6, *A. aciculatus* leaves 80.3, *I. cylindrica* roots 9.6, *S. spontaneum* roots 27.3, *A. aciculatus* roots 13.3. At 96 hours, the average number of eggs hatched for each sample was as follows: *I. cylindrica* leaves 77.3, wild sugarcane leaves 70.3, needle grass leaves 66, *I. cylindrica* roots 14, *S. spontaneum* roots 19.3, *A. aciculatus* roots 17.6.

Based on the Analysis of Variance for the Effects of Weed Extract Toxicity Testing on *Ae. aegypti* mosquito larvae's egg-laying ability at 72 hours, the average number of eggs hatched in the larval test treatment at a concentration of 1000 ppm was as follows: for cogon grass leaves (75.00), wild sugarcane leaves (77.67), needle grass leaves (80.33), *I. cylindrica* roots (9.67), *S. spontaneum* roots (27.33), and *A. aciculatus* roots (13.33). The obtained p-value was 0.000, indicating a significant difference in the average number of eggs hatched among the six samples.

It is similarly based on the variance analysis of weed extract toxicity testing effects on *Ae. aegypti* mosquito larvae's egg-laying ability at 96 hours, the average number of eggs hatched in the larval test treatment at a concentration of 1000 ppm was as follows: for *I. cylindrica* leaves (76.67), *S. spontaneum* leaves (90.00), *A. aciculatus* leaves (83.00), *I. cylindrica* roots (14.00), *S. spontaneum* roots (19.33), and *A. aciculatus* roots (17.67). The obtained p-value was 0.000, indicating a significant difference in the average number of eggs hatched among the six samples.

Based on the analyses above, it was evident that there was a significant difference in the average number of

eggs hatched among the six testing samples, both at the 72-hour and 96-hour time intervals. From these results, it could be concluded that in the testing of *I. cylindrica* root extracts, the number of hatched eggs was lower than the number of eggs in the testing of *I. cylindrica* leaf extracts. Based on phytochemical test results, the *I. cylindrica* root extracts contain flavonoid compounds with phenolic content (Maharani et al. 2018).

The crude extracts of leaves, flowers, fruits, rhizomes, and endosperms of plants exhibited significant larvicidal activity against *Ae. aegypti* was measured by lethal concentration (LC) and larval mortality (Priya et al. 2023). The ethanolic seed extract of *Annona mucosa* (Annonaceae) demonstrated an LC50 value of 2.6 mg/ml against *Ae. aegypti* larvae within 24 hours (Rodrigues et al. 2021). The dichloromethane extract from *Ateleia glazioviana* (Fabaceae) leaves showed significant larvicidal activity at a concentration of 500 µg/ml. *Annona glabra*, the aqueous extract, exhibited LC50 values of 2.43 mg/L and 1.17 mg/L against *Ae. aegypti* larvae at 24 and 48 hours, respectively (Almadiy 2020; Amarasinghe et al. 2020). The steroidal alkaloid solanidine, extracted from the fruit of *Solanum paludosum*, resulted in a 63% mortality rate of fourth instar larvae at a concentration of 150 µg/ml (Cruz et al. 2019).

Based on the research by Alivianti (2021), during a 15-day observation period, mosquito eggs in ovitraps in the control group (aquades) had a small portion (21%) of mosquito eggs with a count of 46 eggs. In contrast, the intervention group with water soaked in *I. cylindrica* extracts showed a smaller portion (16%) of mosquito eggs, with a count of 191 eggs, indicating the influence of using soaked water from *I. cylindrica* on the number of mosquito eggs trapped in the ovitraps.

These findings align with the research conducted by Ridha et al. (2019), which found that using *I. cylindrica*-soaked water resulted in the highest oviposition compared to other soaking methods. For hatching results, rice-soaked water, water from egg residues, and aquades all produced high hatching rates, while soaking in lemon grass and cogon grass resulted in lower hatching rates (Ridha et al. 2019).

Acknowledgements

The authors would like to express their sincere gratitude to the Health Research and Development Center, Baturaja, South Sumatra, Indonesia, for granting permission to

conduct the research, and all parties for their support and collaboration throughout the research process.

References

- Alivianti, A., Atmaja, B.P., Udiyani, R., 2021. The effect of *Imperata cylindrica* soaking water on the number of *Aedes spp* mosquito eggs trapped in ovitraps in efforts to prevent dengue Hemorrhagic Fever (DHF). *Nursing Science Journal*. 5. <https://doi.org/10.30737/nsj.v5i2.1925>
- Al-Massarani, S., El-Shaibany, A., Tabanca, N., Ali, A., Estep, A.S., Becnell, J.J., Goger, F., Demirci, B., El-Gamal, A., Baser, K.H.C., 2019. Assessment of selected Saudi and Yemeni plants for mosquitocida activities against the yellow fever mosquito *Aedes aegypti*. *Saudi Pharm J*. 27, 930-938. DOI: 10.1016/j.jsps.2019.07.001
- Almadiy, A.A., 2020. Chemical composition, insecticidal and biochemical effects of two plant oils, and their significant fractions against *Aedes aegypti*, the common dengue fever vector. *Heliyon*. 6, e04915. DOI:10.1016/j.heliyon.2020.e04915.
- Amarasinghe, L., Wickramarachchi, P., Aberathna, A., Sithara, W., De Silva, C., 2020. Comparative study on the larvicidal activity of green synthesized silver nanoparticles and *Annona glabra* (Annonaceae) aqueous extract to control *Aedes aegypti* and *Aedes albopictus* (Diptera: Culicidae). *Heliyon*. 6, E04322. DOI:10.1016/j.heliyon.2020.e04322
- Antou, R.N., Repi, R.A., Tulu, M.L.S., 2023. Phytochemical test and effectiveness of alang-alang stolon extract (*Imperata cylindrica* L.) against mortality of Dengue Fever mosquito larvae (*Aedes sp.*) *Biodiversitas*. 3, 33-39. DOI:10.53682/ibj.v3i2.5738
- Attaullah, M., Gul, S., Bibi, D., Andaleeb, A., Ilahi, I., Siraj, M., Ahmad, M., Ullah, I., Ali, M., Ahmad, S., Ullah, Z., 2023. Diversity distribution and relative abundance of the mosquito fauna (Diptera: Culicidae) of Malakand and Dir. Lower Pakistan. *Braz. J Biol.* 83, e247374. DOI:10.1590/1519-6984.247374
- Azevedo, F.R., Maciel, G.C., Oliveira, G.B.S., Mesquita, F.O., Alves, A.C.L., 2019. Larvicidal activity of native plant extracts from the Araripe National Forest on *Aedes aegypti*. *The Journal of Agricultural Science*. 11, 105. <https://doi.org/10.5539/jas.v11n7p105>
- Benelli, G., 2018. Mode of action of nanoparticles against insects. *Environmental Science and Pollution Research International*. 25, 12329-12341. DOI:10.1007/s11356-018-1850-4.Pmid:29611126
- Calago, P.M., Dayao, M.R., Roleda, P.K., Saquilabon, C.R., Tecson, J.M., Wee, J.A.M., Arong, E.A., 2015. The effect of cogon grass (*Imperata cylindrica*) and carabao grass (*Paspalum conjugatum*) leaf extract on mortality of *Aedes aegypti* larvae. Herdin Record : R07-CDU-16091009200196.

- 81 Chellappandian, M., Thanigaivel, A., Vasantha-Srinivasan, P., Edwin, E.S., Ponsankar, A., Rani, S., Kalaivani, K., Sengottayan, S.N., 2017. Toxicological effects of *Sphaeranthus indicus* linn (asteraceae) leaf essential oil against human disease vector. *Culex quinquefasciatus* and *Aedes aegypti* linn and impacts on a beneficial mosquito predator. *Environ. Sci. Poll. Res.* 25, 10294-10306. DOI:10.1007/s11356-017-8952-2
- 22 Cruz, I.L.S., Pardai, B.M., Alves, S.P., Medeiros, G.A.M., Barbosa-Filho, J.M., Maleck, M., 2019. *Alcaloide esteroidal*, substancia de *Solanum poludosum*, com atividade larvica sobre *Aedes aegypti*. *Revista de Saude.* 10, 15-19. <https://doi.org/10.21727/rs.v10i02.1812>
- 68 Demirak, M.S.S., Canpolat, E., 2022. Plant-based bioinsekticides for mosquito control: Impact on insecticide resistance and disease transmission. *Insect.* 13, 162. DOI:10.3390/insects13020162
- 25 Duval, P., Aschan-Leygonie, C., Moro, C.V., 2023. A review of knowledge, attitudes and practices regarding mosquitoes and mosquito-borne infectious diseases in nonendemic regions. *Front Public Health.* 8, 11. doi: 10.3389/fpubh.2023.1239874
- 50 Ekawati, E.R., Santoso, S.D., Purwanti, Y.R., 2017. Utilization of lime fruit peel (*Citrus aurantifolia*) as *Aedes aegypti* instar III larvicide. *Biota.* 31-5. DOI:10.19109/Biota.v3i1.926
- Fitriyani, N.L., Ristiawati, Mastutui, D.N.R., 2022. Water filter with galangal extract biolarvicides as an effort for Dengue vector control. *Balaba.* 18 19-26. DOI:10.22435/blb.v18i1.5003
- 39 Giroth, S.J., Bernadus, J.B.B., Sorisi, A.M.H., 2019. Efficacy Test of Lemongrass (*Cymbopogon citratus*) Plant extract on mortality rates of *Aedes sp* larvae. *eBiomedicine.* 9, 13-20 DOI: 10.35790/ebm.9.1.2021.31716
- Hastono., 2018. *Data Analysis in the Health Sector*. Rajawali Press, Depok.
- 71 Kamaraj, C., Deepak, P., Balasubramani, G., Karthi, S., Asrul, D., Aiswarya, D., Vadivel, A., Vimalkumar, E., Mathivanan, D., Suseem, S.R., Pandian, C.K.M., Nathan, S.S., Perumal, P., 2018. Target and non-target toxicity of fern extracts against mosquito vectors and benefecial aquatic organisms. *Ecotoxicol. Environ. Saf.* 161, 221-230. DOI:101016/j.ecoenv.2018.05.062.
- 6 Kirik, H., Burtin, V., Tummeleht, L., Kurina, O., 2021. Friends in all the green spaces: Weather dependent changes in urban mosquito (Diptera: Culicidae) abundance and diversity. *Insects.* 12, 352. <https://doi.org/10.3390/insects12040352>
- 19 Lim, H., Lee, S.Y., Ho, L.Y., Sit, N.W., 2023. Mosquito larvicidal activity and cytotoxicity of the extracts of aromatic plants from Malaysia. *Insects.* 14, 512. DOI:10.3390/insects14060512
- 64 Listiono, H., Damiri, N., Kamaluddin T., Irsan, C., 2023. Diversity od disease-carrying mosquito vectors in potential areas of dengue fever in Palembang City, South Sumatra, Indonesia. *Biodiversitas.* 25, 474-481. DOI: 10.13057/biodiv/d250205
- 38 Maharani, A.R., Restuti, C.T., Sari, E., Wahyuningsih, N.E., Murwani, R., Hapsari, M., 2018. Nutrient intake of dengue hemorrghic fever patients in Semarang city. *J. Phys:Conf.Ser.* 1025, 012059. DOI:10.1088/1742-6596/1025/1/012059
- 57 Muangmoon, R., Junkum, A., Chaithong, U., Jitpakdi, A., Riyan, D., Wannasan, A., Somboom, P., Pitasawat, B., 2018. Natural larvacides of botanical origin against dengue vector *Aedes aegypti* (Diptera:culicidae). *The Southeast Asian Journal of Tropical Medecine and Public Health.* 49, 227-239.
- 47 Nebbak, A., Almeras, L., Parola, P., Bitam, I., 2022. Mosquito vectors (Diptera: Culicidae) and Mosquito-borne diseases in North Africa. *Insect.* 13, 962. <https://doi.org/10.3390/insects13100962>
- 31 Pavela, R., 2016. History, presence and perspective of using plant extract as comercial botanical insecticides and farm products for protection against insects- a review. *Plant Protection Science.* 52, 229-241. DOI:10.17221/31/2016-PPS
- Pavela, R., Maggi, F., Iannarelli, R., Benelli, G., 2019. Plants extracts for developing mosquito larvicides: from laboratory to the field, with insight on the modes of action. *Acta Trop.* 178, 236-271. DOI:10.1016/j.actatrop.2019.01.019
- 78 Pereire, V., Figueira, O., Castilho, P.C., 2024. Flavonoid as insecticides in crop protction: A review of current research and future prospects. *Plants.* 13, 776. DOI:103390/plants13060776.
- 6 Pessoa, L.Z.D.S., Daurte, J.L., Ferreira, R.M.A., Oliveira, A.L.M.F.M., Cruz, R.A.S., Faustino, S.M.M., Carvalho, J. C.T.C., Fernandes, C.P., Sauto, R.N.P., Araujo, R.S., 2018. Nanosuspension of quercetin: Preparation, characterization and effect agains *Aedes aegypti* larvae. *Res Bras Farm.* 28, 618-625. DOI:10.1016/j.bjp.2018.07.003
- 63 Priya, S.S., Vasantha-Srivasan, P., Altermimi, A.B., Keerthana, R., Radhakrishna, N., Senthil-Nathan, S., Kalaivani, N. C., Chandrasekar, N., Karthi, S., Ganesan, R., Alkanan, Z.T., Pal, T., Verma, O.P., Prockow, J., 2023. Bioactive molecules derived from plants in managing dengue vector *Aedes aegypti* (Linn). *J. Molecules.* 28, 2386. DOI:10.3390/molecules28052286
- 45 Ramayanti, I., Herlinda, S., Muslim, A., Hasyim, H., Anwar, C., Suwandi, S., Damiri, N., Irsan, C., Verawaty, M., 2023. Pathogenicity of entomopathogenic fungi to eggs, larvae, and adults and their effects on development of *Aedes albopictus*. *Biodiversitas.* 24, 4766-4774. DOI: 10.13057/biodiv/d240917
- 32 Raul, P.K., Santra, P., Goswami, D., Tyagi, V., Yellappa, C., Mauka, V., Devi, R.R., Chattopadhyay, P., Jayaram, R.V., Dwivedi, S.K., 2021. Green synthesis of carbon dotsilver nanohybrids from fruits and vegetable's peel waste: Applications as potent mosquito larvicide. *Current Research in Green and Sustainable Chemistry.* DOI:10.1016/j.crgsc.2021.100158.

- 66 Ridha, M.R., Fadilly, A., Hairani, B., Sembiring, W.R., Meliyanie, G., 2019. Effectiveness of attractants on hatchability and number of *Aedes albopictus* mosquito eggs in the Laboratory. *ASPIRATOR-Journal of Vector-Borne Disease Studies*. 11, 43-106. <https://doi.org/10.22435/asp.v11i2.1164>
- Ridha, M.R., Marlinae, L., Zubaidah, T., Fadillah, N.A., Widjaya, J., Rosadi, D., Rahayu, N., Ningsih, M., Desimal, I., Sofyandi, A., 2023. Control methods for invasive mosquitoes of *Aedes aegypti* and *Aedes albopictus* (Diptera: Culcidae) in Indonesia. *Veterinary World*. 16, 1952-1963. DOI:10.14202/vetworld.2023.1952-1963
- 26 Rodrigues, A.M., da Silva, A.A., de Freitas, J.C.C., Martin, V.E.P., Ferreira, M.A.P., Ferreira, A.C.S., Cabeca, C.L.S., Rodrigues, A.L.M., Alves, D.R., de Moraes, S.M., 2021. Larvicidal activity of *Annona mucosa* Jacq. Extract and main constituents rolipatin 1 and rolipatin against *Aedes aegypti* and *Aedes albopictus*. *Ind. Crop. Prod.* 169, 113678. <https://doi.org/10.1016/j.indcrop.2021.113678>
- 20 Kusdy, M., 2020. *Imperata cylindrica*: Reproduction, Dispersal, and Controls. *CAB Reviews* 2020. 15, No. 038. doi: 10.1079/PAVSNNR202015038
- Siam, M.A.H., Owareat J.K., Khan A.R., 2022. Mosquito control management using phytochemicals: A review. *International Journal of Mosquito Research*. 9, 10-17. <https://doi.org/10.22271/23487941.2022.v9.i4a.617>
- 49 Swargiary, A., Daimari, M., Roy, M., Haloi, D., Ramchiary, B., 2019. Evaluation of phytochemical properties and larvicidal activities of *Cynodon dactylon*, *Clerodendrum viscosum*, *Spilanthes acmella* and *Terminalia chebula* against *Aedes aegypti*. *Asian Pac. J Trop Med*. 12, 224-231. <https://doi.org/10.4103/1995-7645.259243>
- 87 Tungmunthum, D., Thongboonyou, A., Pholboon, A., Yangsabai, A., 2018. Flavonoid and other phenolic compounds from medicinal plants for pharmaceutical and medical aspects: An overview. *Medecines*. 5, 93. DOI:10.3390/medecines5030093
- Utomo, D.S., Kristianti, E.B.E., Mahardika A., 2020. Effect of growing location on the levels of flavonoids, phenolics, chlorophyll, carotenoids and antioxidant activity in horsehip plants (*Stachytarpheta jamaicensis*). *Biomes*. 22, 143-149.
- 67 Vasantha-Srinivasan, P., Thanigaivel, A., Edwin, E.S., Ponsankar, A., Senthil-Nathan, S., Rani, S., Kalaivani, K., Hunter, W.B., Duraipandiyan, V., Al-Dhabi, N.A., 2018. Toxicological effects of chemical constituents from Piper against the environmental burden *Aedes aegypti* Liston and their impact on non-target toxicity evaluation against biomonitoring aquatic insects. *Environ. Sci. Pollut. Res.* 25, 10434-10446. DOI:10.1007/s11356-017-9714-x
- 13 Vivekanandhan, P., Venkatesan, R., Ramkumar, G., Karthi, S., Senthil-Nathan, S., Shivakumar, M.S., 2018. Comparative analysis of major mosquito vectors response to seed-derived essential oil and seed pod-derived extract from *Acacia nilotica*. *Intern. J. Environ. Res. Public Health*. 15, 388. DOI:10.3390/ijerph15020388
- Wibowo, S., Hendra, D., 2015. Characteristics of bio-oil from reed grass (*Saccharum spontaneum*) using a fast pyrolysis process. *Journal of Forest Research*. 33, 347-365. <https://doi.org/10.20886/jphh.2015.33.4.347-363>
- 1 World Health Organization. 2005. Guidelines for laboratory and field testing of mosquito larvicides. WHO/CDS/WHOPES/GCDPP/2005. <https://www.who.int/publication>
- 40 World Health Organization. 2023. Dengue-Global situation. <https://www.who.int/newroom>
- 7 Zhang, M., Zhao, J., Dai, X., Li, X., 2023. Extraction and analysis of chemical compositions of natural products and plants. *Separations*. 10, 558. DOI:10.3390/separations10120598

Effectiveness of Biolarvicides of Imperata cylindrica, Saccharum spontaneum and Andropogon aciculatus on Aedes aegypti larval Mortality and Egg-laying Ability in Adults

ORIGINALITY REPORT

13%
SIMILARITY INDEX

PRIMARY SOURCES

1	publish.sub7journal.com Internet	30 words — < 1 %
2	sifisherliessciences.com Internet	30 words — < 1 %
3	www.tandfonline.com Internet	30 words — < 1 %
4	Kala Agbo bidi, Luís Almeida, Jean-Michel Coron. "Global Stabilization of a Sterile Insect Technique Model by feedback Laws", Journal of Optimization Theory and Applications, 2025 Crossref	29 words — < 1 %
5	oxfordjournals.org Internet	29 words — < 1 %
6	oup.silverchair-cdn.com Internet	23 words — < 1 %
7	upcommons.upc.edu Internet	23 words — < 1 %
8	ejournal3.undip.ac.id Internet	20 words — < 1 %
9	www.e3s-conferences.org	

20 words — < 1 %

10 etheses.iainkediri.ac.id

Internet

18 words — < 1 %

11 acikerisimarsiv.selcuk.edu.tr:8080

Internet

17 words — < 1 %

12 1library.net

Internet

16 words — < 1 %

13 de.genizer.com

Internet

16 words — < 1 %

14 ir.bdu.edu.et

Internet

16 words — < 1 %

15 www.dovepress.com

Internet

16 words — < 1 %

16 www.scilit.net

Internet

16 words — < 1 %

17 Oscar Hernando Pardo Cuervo, Laura Valentina Barrera Martínez, Ingrid Dayana Jiménez Camacho, Andrea Camila Martínez Gordon et al. "Natural extracts as a sustainable alternative for the control of Aedes aegypti: an updated review", Revista de Ciências Agroveterinárias, 2024

Crossref

15 words — < 1 %

18 Rosanna Lorrane Francisco dos Reis Matos, Núbia Nunes de Souza, Sandro Morais dos Santos, Aline Ferreira Rafael et al. "Larvicidal Activity of Leaf Extract From Mauritiella armata (Aceraceae) on Aedes aegypti and Culex quinquefasciatus (Culicidae)", Journal of Agricultural Science, 2022

Crossref

15 words — < 1 %

19 Roodabeh Bahramsoltani, Roohollah Ahmadian, Maria Daglia, Roja Rahimi. " (Mill.) Fuss (Parsley): An Updated Review of the Traditional Uses, Phytochemistry, and Pharmacology ", Journal of Agricultural and Food Chemistry, 2024

Crossref

20 floram.org 14 words — < 1 %

Internet

21 jurnal.ipb.ac.id 14 words — < 1 %

Internet

22 novapublishers.com 14 words — < 1 %

Internet

23 biomedicineonline.org 13 words — < 1 %

Internet

24 Aarti Sharma, Monika Mishra, Roopa Rani Samal, Vinay Singh Dagar, Manoj Kumar, Anupama Shukla, Sarita Kumar. "Influence of Cassia occidentalis leaf and stem extracts on the life parameters of Aedes aegypti (Linnaeus, 1762)", Journal of Applied and Natural Science, 2024

Crossref

25 Basiliana Emidi. "Establishing and sustaining women-led vector control groups: community perspectives from Misungwi and Ilemela districts, Mwanza, Tanzania", The Journal of Basic and Applied Zoology, 2025

Crossref

26 Bassam S. M. Al Kazman, Joanna E. Harnett, Jane R. Hanrahan. "Traditional Uses, Phytochemistry and Pharmacological Activities of Annonaceae", Molecules, 2022

Crossref

27 Sharif Ahmed, Bhagirath Singh Chauhan. "Performance of Different Herbicides in Dry-Seeded Rice in Bangladesh", The Scientific World Journal, 2014

Crossref

28	jurnal.uns.ac.id Internet	12 words — < 1 %
29	pesquisa.bvsalud.org Internet	12 words — < 1 %
30	sanad.iau.ir Internet	12 words — < 1 %
31	sciendo.com Internet	12 words — < 1 %
32	www.ictmumbai.edu.in Internet	12 words — < 1 %
33	Ashley L. Muller, Luisa Diaz-Arias, Mackenzie C. Cervenka, Tanya J.W. McDonald. "The effect of anti-seizure medications on lipid values in adults with epilepsy", <i>Epilepsy & Behavior</i> , 2023 Crossref	11 words — < 1 %
34	Fatma Benjeddou, Ines Mnif, Marie Rossignol, Dhouha Ghribi, Slim Tounsi, Fabrice Chandre, Raida Zribi Zghal. "Towards sustainable vector control: Innovative substrate identification for developing eco-friendly, cost-effective, and highly potent biopesticides", <i>Biological Control</i> , 2025 Crossref	11 words — < 1 %
35	Qian Chen, Yizhen Yuan, Fangzhou Cai, Zhe Li, Qiang Wei, Wei Wang. "Single-Nucleus and Spatial Transcriptomics Revealing Host Response Differences Triggered by Mutated Virus in Severe Dengue", <i>Viruses</i> , 2024 Crossref	11 words — < 1 %
36	ampllaeditora.com.br Internet	11 words — < 1 %
37	avesis.ankara.edu.tr Internet	11 words — < 1 %

-
- 38 nlist.inflibnet.ac.in 11 words — < 1 %
Internet
-
- 39 repositori.unsil.ac.id 11 words — < 1 %
Internet
-
- 40 www.ajtmh.org 11 words — < 1 %
Internet
-
- 41 Fred Ssempijja, Samuel Sunday Dare, Edmund E. M. Bukenya, Keneth Iceland Kasozi et al. "Attenuation of Seizures, Cognitive Deficits, and Brain Histopathology by Phytochemicals of (Poaceae) in Acute and Chronic Mutant Epilepsy Models ", Journal of Evidence-Based Integrative Medicine, 2023
Crossref
-
- 42 Govindharaj Guru-Pirasanna-Pandi, Totan Adak, Basana Gowda, Naveenkumar Patil, Mahendiran Annamalai, Mayabini Jena. "Toxicological effect of underutilized plant, Cleistanthus collinus leaf extracts against two major stored grain pests, the rice weevil, Sitophilus oryzae and red flour beetle, Tribolium castaneum", Ecotoxicology and Environmental Safety, 2018
Crossref
-
- 43 Hongbin Zou, Haoyi Gu, Jialu Cheng, Chao Tian, Qilong Shu, Peilin Peng, Bing Li. "The Mechanism of Damage to the Midgut by Low Concentration of Bacillus thuringiensis in the Silkworm, Bombyx mori", Insects, 2024
Crossref
-
- 44 Manikandan Vani Raju, Meenakshi Kaniyur Chandrasekaran, Meenakshi Sundari Rajendran, Gopalakrishnan Velliyur Kanniappan et al. "Deciphering the therapeutic, larvicidal, and chemical pollutant degrading properties of leaves-mediated silver nanoparticles obtained from Alpinia purpurata", BioResources, 2024
Crossref

45 S Ananda, R Anwar. " The presence of the entomophthoralean fungus during a outbreak affecting shallot in Brebes and Tegal Regencies ", IOP Conference Series: Earth and Environmental Science, 2025
Crossref 10 words — < 1 %

46 Zainuddin Nawawi, H Ahmad, David Caulker, Annuar Mohd Ramli, Muhammad Abu Bakar Sidik, S.H. Dahlan. "On the interaction of 132 kV transmission system with communication line: Street cabinet resistivity evaluation", 2017 International Conference on Electrical Engineering and Computer Science (ICECOS), 2017
Crossref 10 words — < 1 %

47 aj.tubitak.gov.tr
Internet 10 words — < 1 %

48 etheses.bham.ac.uk
Internet 10 words — < 1 %

49 mzuir.inflibnet.ac.in
Internet 10 words — < 1 %

50 repository.universitas-bth.ac.id
Internet 10 words — < 1 %

51 www.jorae.cn
Internet 10 words — < 1 %

52 Angela H. Valente, Matthew de Roode, Madeleine Ernst, Miguel Peña-Espinoza et al. "Identification of compounds responsible for the anthelmintic effects of chicory (Cichorium intybus) by molecular networking and bio-guided fractionation", International Journal for Parasitology: Drugs and Drug Resistance, 2021
Crossref 9 words — < 1 %

53 Ben-Dov, Eitan, and Yoel Margalith. "Biological Control by Bacillus thuringiensis subsp. 9 words — < 1 %

- 54 Ran Wang, Yuying Li, Lili Guo, Jiao Wu, Sitong Wang, Bixi Li, Dingding Zuo, Qi Guo, Xiaogai Hou. 9 words — < 1%

"Efficient Agrobacterium rhizogenes-mediated transformation system for fleshy roots of Paeonia 'Coral Sunset' without tissue culture", Industrial Crops and Products, 2025

Crossref

- 55 Roman Pavela, Filippo Maggi, Giulio Lupidi, Hélène Mbuntcha et al. "Clausena anisata and Dysphania ambrosioides essential oils: from ethno-medicine to modern uses as effective insecticides", Environmental Science and Pollution Research, 2017 9 words — < 1%

Crossref

- 56 Samuel M. Waititu, Fugui Yin, Rob Patterson, Juan C. Rodriguez-Lecompte, Charles M. Nyachoti. 9 words — < 1%

"Short-term effect of supplemental yeast extract without or with feed enzymes on growth performance, immune status and gut structure of weaned pigs challenged with Escherichia coli lipopolysaccharide", Journal of Animal Science and Biotechnology, 2016

Crossref

- 57 Siriluk Sintupachee, Suppawan Promprao. 9 words — < 1%

"Phytochemical Profiles and Mosquito-Repellent Properties of Indigenous Plants from Pak Panang District, Nakhon Si Thammarat, Thailand", Malaysian Applied Biology, 2023

Crossref

- 58 Tong, Fan, and Jeffrey R. Bloomquist. "Plant Essential Oils Affect the Toxicities of Carbaryl and Permethrin Against <I>Aedes aegypti</I> (Diptera: Culicidae)", 9 words — < 1%

Journal of Medical Entomology, 2013.

Crossref

59 Yousef Abdulwahab, Mohammed Al-Shmery, Ahyar Ahmad, Isra Wahid, Paulina Taba. "Effectiveness of Coffee Plant Extracts and Green Nanoparticles for Mosquito Control of the Dengue Vector *Aedes aegypti*: A Comprehensive Review", Bangladesh Journal of Infectious Diseases, 2025

Crossref

60 assets.researchsquare.com 9 words — < 1 %

Internet

61 clinphytoscience.springeropen.com 9 words — < 1 %

Internet

62 conservancy.umn.edu 9 words — < 1 %

Internet

63 cris.bgu.ac.il 9 words — < 1 %

Internet

64 discovery.researcher.life 9 words — < 1 %

Internet

65 e-sciencecentral.org 9 words — < 1 %

Internet

66 ejournal2.litbang.kemkes.go.id 9 words — < 1 %

Internet

67 ethesis.lib.ku.ac.th 9 words — < 1 %

Internet

68 jebas.org 9 words — < 1 %

Internet

69 journals.plos.org 9 words — < 1 %

Internet

70 moam.info 9 words — < 1 %

Internet

-
- 71 www.apjtm.org 9 words — < 1 %
Internet
-
- 72 www.astmh.org 9 words — < 1 %
Internet
-
- 73 www.scienceopen.com 9 words — < 1 %
Internet
-
- 74 Bhakti Yudho Suprpto, A. Wahyudin, Hera Hikmarika, Suci Dwijayanti. "The Detection System of Helipad for Unmanned Aerial Vehicle Landing Using YOLO Algorithm", Jurnal Ilmiah Teknik Elektro Komputer dan Informatika, 2021 8 words — < 1 %
Crossref
-
- 75 Mathiyazhagan Narayanan, Maha A. Alshiekheid, Mythili Saravanan. "Antibacterial, mosquito larvicidal, and cytotoxicity potential of AgNPs synthesized using *Pittosporum undulatum* under in vitro conditions", Environmental Research, 2024 8 words — < 1 %
Crossref
-
- 76 Roman Pavela. "Essential oils from *Foeniculum vulgare* Miller as a safe environmental insecticide against the aphid *Myzus persicae* Sulzer", Environmental Science and Pollution Research, 2018 8 words — < 1 %
Crossref
-
- 77 Tutut Indria Permana, Habibatul Khoiriyyah, Atok Miftachul Hudha, Moh. Mirza Nuryady, Muhammad Yusril Ihya Maksum. " Mortality Effect of Bintaro (*L.*) Seed Extract Against (Diptera: Culicidae) Mosquitoes ", BIO Web of Conferences, 2024 8 words — < 1 %
Crossref
-
- 78 Ying Zhang, Xinhang Wang, Zhipeng Bian, Chenchen Se, Guoqing Yang, Yanhui Lu. "Application of flavonoid compounds suppresses the cotton aphid, *Aphis gossypii*", Frontiers in Plant Science, 2025 8 words — < 1 %
Crossref

79	ejournal.undip.ac.id Internet	8 words — < 1 %
80	journal3.uin-alauddin.ac.id Internet	8 words — < 1 %
81	pubblicazioni.unicam.it Internet	8 words — < 1 %
82	El-Mehdi El-Assri, Anouar Hmamou, Youssef El-Assri, Asmae Baghouz et al. "Molecular characterization, GC-MS analysis, and biological activities of <i>Matricaria chamomilla</i> var. <i>recutita</i> L. essential oils from Taounate, Morocco: In vitro and in silico investigations of antioxidant, antimicrobial, and insecticidal effects", Springer Science and Business Media LLC, 2025 Crossref Posted Content	7 words — < 1 %
83	Karin Bakran-Lebl, Stefanie Pree, Thomas Brenner, Eleni Daroglou et al. "First Nationwide Monitoring Program for the Detection of Potentially Invasive Mosquito Species in Austria", Insects, 2022 Crossref	7 words — < 1 %
84	Mona Piplani, Deepak P. Bhagwat, Gautam Singhvi, Murugesan Sankaranarayanan, Rafael Balana-Fouce, Tarini Vats, Subhash Chander. "Plant-based larvicidal agents: An overview from 2000 to 2018", Experimental Parasitology, 2019 Crossref	7 words — < 1 %
85	Rohini Keshava, Nagesh Muniyappa, Rajalakshmi Gope, Ananthanarayana Saligrama Ramaswamaiah. "Anti-Cancer Effects of <i>Imperata cylindrica</i> Leaf Extract on Human Oral Squamous Carcinoma cell line SCC-9 in Vitro", Asian Pacific Journal of Cancer Prevention, 2016 Crossref	7 words — < 1 %
86	ijphs.iaescore.com Internet	7 words — < 1 %

EXCLUDE QUOTES ON
EXCLUDE BIBLIOGRAPHY OFF

EXCLUDE SOURCES OFF
EXCLUDE MATCHES OFF