



Figure 4. In Vitro Rumen Degradation of Cellulose

Although many factors determine the quality of forages, based on the chemical composition, *in vitro* degradation and *in vitro* gas production, it could be concluded that in part, nutritive value of *Leucaena leucocephala* followed by *Centrosema pubescens* was higher than those of the other legumes grown in South Sumatra, Indonesia.

References

- A.O.A.C., 1984. Official Methods of Analysis (13 th Ed.). Association of Official Analytical Chemists: Washington, D.C.
- Goering, H.K. and P.J. Van Soest, 1970. Forage fiber analysis (apparatus, reagents, procedures and some application). Agric. Hand-book 379, ARS, USDA, Washington, D.C.
- Humphreys, L.R., 1995. Diversity and productivity of tropical legumes. In: Tropical Legumes in Animal Nutrition. Edited by J.P.F. D'Mello and C. Devendra. Cal, International. pp. 1-21.
- Menke, K.H., L. Raab, A. Salewski, H. Steingass, D. Fritz and W. Schneider, 1979. The estimation of the digestibility and metabolizable energy content of ruminant feeding stuffs from the gas production when they are incubated with rumen

liquor *in vitro*. J. Agric. Sci. Camb., 93 : 217-222.

- Norton, B.W. and D.P. Poppi, 1995. Comparison and nutritional attributes of pasture legumes. In: Tropical Legumes in Animal Nutrition. Edited by J.P.F. D'Mello and C. Devendra. Cab International. pp. 23-47.
- Skerman, P.J., Cameron, D.G. and F. Riveros, 1988. Tropical Forage Legumes. 2nd Edition. Food and Agriculture Organization of the United Nations.
- Steel, R.G.D. K.I. JR Tomse, 1981. Principles and Procedures of Statistics. A Biometrical Approach (2nd Ed). Mc Graw-Hill International Book Company.
- Tilley, J.M.A and Terry, M.A, 1963. A two stages technique for *in vitro* digestion of forage crops. J. Br. Grassld. Soc. Vol.18: 104-111.
- Van Soest, P.J. 1982. Nutritional Ecology of the Ruminant. Ruminant metabolism, nutritional strategies, the cellulolytic fermentation and the chemistry of forages and plant fibers. O&B Books Inc.