

Characteristics of chitinase isolated from different part of snakehead fish (*Channa striata*) digestive tract

by Ace Baehaki

Submission date: 31-Mar-2019 07:27PM (UTC+0700)

Submission ID: 1102938626

File name: aki_2018_IOP_Conf._Ser._3A_Earth_Environ._Sci._102_012057_1.pdf (607.92K)

Word count: 1421

Character count: 7765

PAPER • OPEN ACCESS

Characteristics of chitinase isolated from different part of snakehead fish (*Channa striata*) digestive tract

To cite this article: A Baehaki *et al* 2018 *IOP Conf. Ser.: Earth Environ. Sci.* **102** 012057

View the [article online](#) for updates and enhancements.

Characteristics of chitinase isolated from different part of snakehead fish (*Channa striata*) digestive tract

A Bachaki^{1*}, S D Lestari¹, Y Wahidman¹, N Gofar²

¹Fisheries Product Technology Study Program, Faculty of Agriculture, Sriwijaya University, Indralaya, South Sumatera, Indonesia

²Soil Sciences study Program, Faculty of Agriculture, Sriwijaya University, Indralaya, South Sumatera, Indonesia.

*E-mail: ace76_none@yahoo.com

Abstract. Naturally, snakehead fish (*Channa striata*) is a prodigious carnivore feeding mainly on live animals, including small shrimp. Based on its feeding habits, the digestive tracts of snakehead is considered as auspicious source of various enzymes including chitinase. The purpose of this study was to partially characterize chitinase enzyme isolated from digestive tract of snakehead fish. Two parts of digestive tract, stomach and intestine were used as enzymes' source. The results showed that chitinase activity from the stomach was higher than chitinase activity from the intestine. The pH and temperature optimum of chitinase activity from digestive tract (the stomach and the intestine) were 6.0 and 70 °C, respectively.

Keywords: *Channa striata*, chitinase, digestive tract

1. Introduction

Chitin is a polysaccharide created by β -1,4 bonding of N-acetyl glucosamine residues. Chitinases are glycosyl hydrolases, characterized for hydrolyzing β -1,4 linkage of N-acetyl glucosamine present in chitin chains [1]. Chitinases have been divided into two main categories: exochitinases and endochitinases [2]. Chitinase is used to hydrolyze chitin to produce chitin oligomers. Functional properties and physiological activities of chitin oligomers depend on molecular weight and chain length mostly [3]. The digestive tract of fish produces many enzymes and there are many reports describing the enzymes present in digestive tract, including protease [4], chitinase [5], and chitosanase [6]. However, reports on chitinase activity in different part of the digestive tract (stomach and intestine) of snakehead fish (*Channa striata*) are still limited. Therefore, in this research we studied the activity of chitinase isolated from the digestive tract of snakehead fish (*Channa striata*) and partial characterization of this enzyme.

2. Materials And Methods

2.1 Chitinase activity

The chitinase activity was measured according to the Spindler method [7].

2.2 Effect of pH enzyme activity

Optimum pH was determined by assaying in buffer with pH values of 6-9 using citric acid buffer (pH 3-4), citric-phosphate buffer (pH 4-6), phosphate buffer (pH 6-8) and boric buffer (pH 8-9), in the presence of soluble chitin substrate (colloidal chitin).



2.3 Effect of temperature on enzyme activity

Optimum temperature was measured at 30, 40, 50, 60, 70 and 80 °C at pH 7, soluble chitin (colloidal chitin) was used as the substrate.

3. Result And Discussion

3.1 Chitinase Activity

The activity of chitinase isolated from digestive tract of snakehead fish (*Channa striata*) as effected by part difference can be seen in Figure 1.




Figure 1. Chitinase activity from digestive tract of snakehead fish (*Channa striata*)

The results showed that the activity of chitinase isolated from the stomach was higher than the activity from the intestine. Chitinase activity from the stomach was 0.41 U/ml while from the intestine was 0.32 U/ml.

3.2 Effect of pH on chitinase activity

Effect of pH on the activity of chitinase isolated from digestive tract of snakehead fish (*Channa striata*) is presented in Figure 2. Optimum activity of chitinase isolated from the stomach and the intestine was achieved at pH 6. The optimum pH ranges of chitinase activity in the stomach and the intestine of Cod (*Gadus morhua*) were 4.5–5.1 and 5.1–6.5, respectively [5]. The ranges were lower than the activity of the same enzyme isolated from the intestine of discus fish (*Symphysodon aequifasciata*) which was optimum at pH 8 [4] but it were higher than the pH 2 reported for enzyme of discus fish (*Symphysodon aequifasciata*) stomach [4], and digestive proteases (acidic protease) of juvenile *Cichlasoma beani* [8].




Figure 2. Effect of pH on chitinase activity in the digestive tract of *Channa striata* (a=the Stomach, b= the intestine)

3.3 Effect of temperature on chitinase activity

A temperature range between 30 °C and 80 °C were used in this study. Figure 3 showed the effect of temperature on chitinase activity from digestive tract of snakehead fish (*Channa striata*).




Figure 3. Effect of temperature on chitinase activity in the digestive tract of *Channa striata* (a= stomach; b=intestine)

The optimum temperature of 70°C were recorded for both chitinase (the stomach and the intestine) in this study. This were higher than optimum temperature of 55 °C reported for digestive proteases (acidic protease) of juvenile *Cichlasoma beani* [8], 65 °C for *Atractosteus tropicus* digestive proteases [9] and 50 °C for purified trypsins A, B and C from skipjack tuna spleen [10].

4. Conclusion

The chitinase activity from the stomach was higher than chitinase activity from the intestine. The pH and temperature optimum of chitinase activity from digestive tract (the stomach and the intestine) were 6.0 and 70 °C respectively.

Acknowledgments

This research was supported by Proffesion Grant 2016 from Ministry of Research, Technology and Higher Education of the Republic of Indonesia (Kementristekdikti) awarded to Prof. Nuni Gofar, Sriwijaya University.

References

- [1] Bhattacharya D, Nagpure A and Gupta R K, 2007 Bacterial chitinases: properties and potential, *Critical Rev. Biotechnol* **27** 21
- [2] Sahai S A and Manocha M S, Chitinases of fungi and plants: their involvement in morphogenesis and host-parasite interaction, 1993 *FEMSMicrobiol Rev* **11** 317
- [3] Shahidi F, Arachchi J K V and Jeon YJ, 1999 Food applications of chitin and chitosans, *Trends Food Sci Tech* **10** 37
- [4] Chong S C A, Hashim R, Chow-Yang L and Ali BA, 2002 Partial characterization and activities of proteases from the digestive tract of discus fish, *Aquaculture* **203** 321
- [5] Danulat E and Kausch H, 1984 Chitinase activity in the digestive tract of Cod, *Gadus morhua* (L.), *J Fish Biol* **24** 125
- [6] Baehaki A, Lestari S D, Gofar N and Wahidman W, 2016 Partial characterization of chitosanase from digestive tract of *Channa striata*, *Oriental J Chem* **32** 3231
- [7] Spindler KD, Chitinase and chitosanase assays. In: Muzzarelli RAA and Peter MG (eds), 1997 Grottammare European Chitin Society, Italy 229
- [8] Martínez-Cárdenas L, Álvarez-González CA, Hernández-Almeida OU, Frías-Quintana CA, Ponce-Palafox CT, and Castillo-Vargasmachuca S, 2017 Partial Characterization of Digestive Proteases in the Green Cichlid, *Cichlasoma beani*, *Fishes* **2**
- [9] Guerrero-Zarate R, Alvarez-Gonzalez CA, Perales-Garcia MA, Olvera-Novoa N, Frías-Quintana CA, Martinez-Garcia R, Contreras-Sanchez WM, 2014 Partial characterization of digestive proteases in tropicalgar *Atractosteus tropicus* juveniles, *Fish Physiol Biochem* **40** 1021
- [10] Klomkloa S, 2008 Digestive proteinases from marine organisms and their applications, *Songklanakarinn J. Sci. Technol* **30** 37

Characteristics of chitinase isolated from different part of snakehead fish (Channa striata) digestive tract

ORIGINALITY REPORT

86%

SIMILARITY INDEX

45%

INTERNET SOURCES

28%

PUBLICATIONS

86%

STUDENT PAPERS

PRIMARY SOURCES

1

Submitted to Sriwijaya University

Student Paper

86%

Exclude quotes On

Exclude matches < 1%

Exclude bibliography On

PENJELASAN HASIL SIMILARITY DI ATAS:

Hasil uji kemiripan (similarity) artikel yang berjudul: Characteristics of chitinase isolated from different part of snakehead fish (Channa striata) digestive tract terhadap dokumen (primary source) menggunakan program turnitin, hasil kemiripan yang paling tinggi adalah 86%. Karena diuji dengan artikel tersebut.