



Medical Journal of Indonesia



<http://mji.ui.ac.id>

Published by the Faculty of Medicine Universitas Indonesia
Official Scientific Journal of the
Faculty of Medicine Universitas Indonesia
in collaboration with
German-Indonesia Medical Association (DIGM)
Jl. Salemba Raya No. 6 Jakarta 10430
Tel./fax +62-21-2302178

- ✓ Year of transformation
- ✓ Polymorphisms in *pfcr1* and *pfmdr1* genes
- ✓ Allogenic mesenchymal stem cells for bone defect
- ✓ In silico study of estrogen receptor inhibitors
- ✓ Single rod implant monoplant® on serum levonorgestrel cervical mucus viscosity
- ✓ Effect of equiosmolar solutions in traumatic brain injury
- ✓ Venous thromboembolism after major surgery
- ✓ Effect of treadmill vs ergocycle exercise on COPD patients
- ✓ Hepatocellular carcinoma in situs inversus totalis
- ✓ Delayed diagnosis of NPC
- ✓ Challenges on HF management: GP's perspective

Basic Medical Research

Polymorphisms in the *pfert* and *pfmdr1* genes in *Plasmodium falciparum* isolates from South Sumatera, Indonesia

Irsan Saleh,¹ Dwi Handayani,² Chairil Anwar²

¹ Department of Pharmacology, Faculty of Medicine, Universitas Sriwijaya, Palembang, Indonesia

² Department of Parasitology, Faculty of Medicine, Universitas Sriwijaya, Palembang, Indonesia

Abstrak

Latar belakang: Selama lebih dari 10 tahun terakhir, resistensi terhadap obat antimalaria telah menjadi masalah utama bagi kesehatan masyarakat di Asia Tenggara termasuk Sumatera Selatan. Studi ini bertujuan untuk mengidentifikasi adanya polimorfisme pada gen *Plasmodium falciparum* yang berhubungan dengan resistensi klorokuin pada isolat *P. falciparum* di Lahat, Sekayu, Baturaja, dan Palembang.

Metode: Studi molekuler dilakukan untuk mengidentifikasi alel mutan dua gen yang berhubungan dengan resistensi klorokuin pada isolat *P. falciparum* di Sumatera Selatan. Sebanyak 25 pasien diambil darahnya, kemudian dilakukan isolasi DNA. Susunan dari kedua gen (*Plasmodium falciparum* chloroquine resistance transporter/*pfert* dan *Plasmodium falciparum* multidrug resistance/*pfmdr1*) dianalisis dengan menggunakan polymerase chain reaction (PCR) dan restriction fragment length polymorphism (RFLP).

Hasil: Polimorfisme pada *pfert* 76-Thr dan *pfmdr1* 86-Tyr ditemukan pada semua isolat. Temuan ini menjelaskan terjadinya kegagalan pengobatan dengan klorokuin dalam beberapa tahun terakhir di Sumatera Selatan.

Kesimpulan: Penelitian ini menguatkan bahwa PCR-RFLP merupakan cara sederhana dan cepat untuk mendeteksi polimorfisme pada gen yang dapat memprediksi chloroquine resistance (CQR). Walaupun adanya polimorfisme pada gen *pfert* dan *pfmdr1* merupakan penanda CQR yang bermakna, perlu penelitian lebih lanjut mengenai peran polimorfisme ini pada respons obat *in vivo* dan *in vitro*.

Abstract

Background: Over the past decade, antimalarial drug resistance has rapidly become a major public health problem in South East Asia region including South Sumatra. This study aimed to determine the extent of gene polymorphisms associated with chloroquine resistance (CQR) in *P. falciparum* isolates from Lahat, Sekayu, Baturaja and Palembang district.

Methods: A molecular study was conducted to identify the mutant alleles of the genes associated with the resistance to chloroquine among the isolates of *Plasmodium falciparum* from South Sumatera. Blood from 25 patients was collected, DNA was isolated, and the sequences of two different genes (*Plasmodium falciparum* chloroquine resistance transporter/*pfert* and *Plasmodium falciparum* multidrug resistance/*pfmdr1*) were analyzed using polymerase chain reaction (PCR) and restriction fragment length polymorphism (RFLP).

Results: This study identified polymorphism in the *pfert* 76-Thr in all isolates and *pfmdr1* 86-Tyr. These findings may reflect the failure of treatment with the standard dose of chloroquine within the last few years in South Sumatera.

Conclusion: PCR-RFLP technique provide a simple and rapid method of detecting polymorphisms in genes that may predict chloroquine resistance (CQR). Although the identification of the polymorphism in the *pfert* and *pfmdr1* genes provides a significant indicator of CQR, further studies are needed to determine the role of these polymorphisms in the *in vivo* and *in vitro* responses to drug treatment.

Keywords: chloroquine, *Plasmodium falciparum*, *pfmdr1*, *pfert*

pISSN: 0853-1773 • eISSN: 2252-8083 • <http://dx.doi.org/10.13181/mji.v23i1.679> • Med J Indones. 2014 ;23:3-8
Correspondence author: Irsan Saleh, irsan_saleh_hazani@yahoo.com

M. Adham
D. Rohdiana
I.D. Mayangsari
Z. Musa

Delayed diagnosis of nasopharyngeal carcinoma in a patient with early signs of unilateral ear disorder

52

Nasopharyngeal carcinoma is frequently overlooked due to nonspecific manifestation. This fact often lead to delayed diagnosis and treatment.

Brief Communication

Rizki
B.B. Siswanto

Challenges on management of heart failure in Indonesia: a general practitioner's perspective

58

Heart failure patients are still incorrectly diagnosed and inadequately treated in Indonesian primary care, despite the availability of current guidelines.

Medical Journal of Indonesia

TABLE OF CONTENTS

Volume 23, Number 1, February 2014, page 1-62, pISSN 0853-1773 - eISSN 2252-8083

Nafrialdi	Editor note Year of transformation	1
<i>Basic Medical Research</i>		
I. Saleh D. Handayani C. Anwar	Polymorphisms in the <i>pfprt</i> and <i>pfmdr1</i> genes in <i>Plasmodium falciparum</i> isolates from South Sumatera, Indonesia All <i>P. falciparum</i> isolates in South Sumatera carry both polymorphism in the <i>pfprt</i> and <i>pfmdr1</i> genes.	3
H.D. Ismail Phedy E. Kholinne A.A. Jusuf N.D. Yulisa	Role of allogenic mesenchymal stem cells in the reconstruction of bone defect in rabbits This study evaluated the transplantation of mesenchymal stem cells, particularly in combination with HA-CaSO ₄ pellets, towards the callus thickness and osteocyte index in bone defects.	9
R. Mustarichie J. Levitas J. Arpina	In silico study of curcuminol, curcumenol, isocurcumenol, and β-sitosterol as potential inhibitors of estrogen receptor alpha of breast cancer This report focus on the in silico studies of curcuminol, curcumenol, isocurcumenol, and β -sitosterol as inhibitors of estrogen receptor alpha of breast cancer.	15
<i>Clinical Research</i>		
E.R. Gunardi B. Affandi	Serum levonorgestrel concentration and cervical mucus viscosity after six months of monoplant® implantation Levonorgestrel concentration is still above therapeutic level until the sixth month of implantation and viscosity of cervical mucus increased immediately following implant insertion.	25
M.R. Ahmad Hanna	Effect of equiosmolar solutions of hypertonic sodium lactate versus mannitol in craniectomy patients with moderate traumatic brain injury Half-molar HSL was as effective as 20% mannitol to produce brain relaxation, with better hemodynamic stability and gave a significant increase in blood glucose level.	30
A.P. Susanto C. Krisnanda D.S-Y. Tan H-Y. Ong D. Pratama R. Soeparwata	Incidence of venous thromboembolism among patients who underwent major surgery in a public hospital in Singapore VTE is a serious complication of major surgery, administration of pharmacologic thromboprophylaxis is advisable	36
H. Turnip A. Ratnawati A. Tulaar F. Yunus A. Kekalih	Comparison of the effects of treadmill and ergocycle exercise on the functional capacity and quality of life of patients with chronic obstructive pulmonary disease Treadmill and ergocycle exercises are some of the reconditioning exercise program to increase the physical capacity and ability to perform daily activities in stable COPD patients.	42
<i>Case Report</i>		
T. Sareo Y.S. Devi L.J. Singh	Hepatocellular carcinoma in situs inversus totalis-a case report Development of hepatocellular carcinoma (HCC) in situs inversus totalis is rare and such condition should be kept in mind while discussing left hypochondrial mass, as diagnostic dilemma may arises in a patient with undiagnosed situs inversus totalis.	48

Editorial Board

Editor-in-chief	Nafrialdi Universitas Indonesia, Indonesia	
Editor-in-chief Emeritus	Isnani A.S. Suryono Universitas Indonesia, Indonesia	
Deputy Editor	Melva Louisa Universitas Indonesia, Indonesia	
Editorial Board Members	Abraham Simatupang DIGM Indonesia, Indonesia Agnes Kurniawan Universitas Indonesia, Indonesia Aria Kekalih Universitas Indonesia, Indonesia Bambang B. Siswanto Universitas Indonesia, Indonesia David H. Garabrant USA Farrokh Habibzadeh Iran Hak Hotta Japan Hans-Joachim Freisleben DIGM Germany, Germany Hans-Jürgen Mägert Germany Harrina E. Rahardjo Universitas Indonesia, Indonesia Inge Sutanto Universitas Indonesia, Indonesia Jeanne A. Pawitan Universitas Indonesia, Indonesia Knut Adermann Germany	Laurentius A. Pramono Universitas Indonesia, Indonesia Markus Meyer Germany Nia Kurniati Universitas Indonesia, Indonesia Pradana Soewondo Universitas Indonesia, Indonesia Rianto Setiabudy Universitas Indonesia, Indonesia Saleha Sungkar Universitas Indonesia, Indonesia Sentot Santoso Germany Septelia I. Wanandi Universitas Indonesia, Indonesia Sri W.A. Jusman Universitas Indonesia, Indonesia Theddeus O.H. Prasetyono Universitas Indonesia, Indonesia Vivian Soetikno Universitas Indonesia, Indonesia Wilfred C.G Peh Singapore Yuditiya Purwosunu Universitas Indonesia, Indonesia
Editorial Assistants	Felix F. Widjaja Universitas Indonesia, Indonesia	Wulan Sari Universitas Indonesia, Indonesia
Business Manager	Jose R.L. Batubara Universitas Indonesia, Indonesia	
Language Editors	Hans-Joachim Freisleben DIGM Germany, Germany	Elizabeth Muliawan Universitas Indonesia, Indonesia
Secretary	Khadijah Buyoyok Indonesia	
Layout Editor	Yudi Tarmizi Indonesia	
Editorial office	Medical Journal of Indonesia Jl. Salemba Raya 5, Jakarta Pusat 10430, Indonesia Telp/fax: +62-21-2302178 E-mail: mji@ui.ac.id	
Publisher	Faculty of Medicine Universitas Indonesia Jl. Salemba Raya 5, Jakarta Pusat 10430, Indonesia Telp/fax: +62 21-3912477 E-mail: humas@fku.ac.id	

Published by the Faculty of Medicine, Universitas Indonesia, Jakarta, Indonesia
Official Scientific Journal of the Faculty of Medicine, Universitas Indonesia in Collaboration
with German-Indonesian Medical Association (DIGM)
pISSN 0853-1773 – eISSN 2252-8063
<http://mji.ui.ac.id>

Focus and Scope

Medical Journal of Indonesia (abbr: Med J Indones) accepts manuscript in basic medical research, clinical research, community research, case report, review article, and brief communication. The journal publishes articles in health sciences (medicine; public health); biology and life sciences (biomedical sciences).

About Us

The Medical Journal of Indonesia was founded in 1991 as the Medical Journal of the University of Indonesia (abbr: Med J Univ Indon). It has been published quarterly consistently and continuously ever since, covering a wide range of medical subject and issues from every medical specialist aspect. In 1995 the name was changed to Medical Journal of Indonesia which reflected the widening of its coverage beyond. The mission of this journal is to provide biomedical scientists clinician researchers, public health researchers and other health care professional with the media to publish their research works.

Peer Review Process

The submitted manuscript is first reviewed by an editor. It will be evaluated in the office whether it is suitable with our focus and scope or has a major methodological flaw. Every submitted manuscript which pass this step will be reviewed by two reviewers. One of the reviewer is appointed from other institutions (national or international). These manuscript will be sent to the reviewers anonymously. Reviewers' comments are then sent to corresponding author to take the necessary actions and responses. The decision of the revised manuscript will then be evaluated in editorial board meeting. The final decision of whom is sent to the corresponding author.

Publication Frequency

This journal is published quarterly.

Open Access Policy

This journal is an open access journal which provides immediate, worldwide, barrier-free access to the full text of all published articles without charge readers or their institutions for access. Readers have right to read, download, copy, distribute, print, search, or link to the full texts of all articles in Medical Journal of Indonesia.

Abstracting and Indexing

Index Medicus for South-East Asia Region (IMSEAR); CAB Abstracts; Global Health; HINARI; Directory of Open Access Journals (DOAJ); Directory of Research Journal Indexing (DRJI); Google Scholar; JournalTOCs; Ulrichsweb Global Serial Directory; WorldCat; New Jour; Electronic Journals Library.

Advertising Policy

Editorial materials will not be influenced by advertisement. Readers can criticize the advertisement by sending it to the office. Advertisement will appear in the print or online version depending on request. For all inquiries, contact the Medical Journal of Indonesia editorial office at Faculty of Medicine Universitas Indonesia, Jalan Salemba Raya 6, Jakarta Pusat 10430, Indonesia; telp/fax: +62-21-2302178; e-mail: mji@ui.ac.id.

Copyright Notice

Faculty of Medicine Universitas Indonesia as publisher reserves the right of first publication of all published material and licensed under a Creative Commons Attribution-NonCommercial-ShareAlike 4.0 International License (<http://creativecommons.org/licenses/by-nc-sa/4.0/>). All statements in articles are the responsibility of the authors.

Subscription

The printed issue should be subscribed for a full calendar year. The price per volume (four issues) and surface postage included: international USD 160, for ASEAN countries USD 100, for Indonesia: institutional IDR 200,000, individual IDR 160,000, and student IDR 100,000.

III cuts the mutant allele into two fragments 248 bp and 124 bp, while the wild-type allele remains undigested. Five of the 25 isolates did not give interpretable results. Of the remaining 20 isolates, all of them (100%) carried the mutant allele (Figure 2).



Figure 2. RFLP for detecting Y86 polymorphism. All sample are mutant alleles. Amplicon 372 bp digested by Afl III into 248 and 124 bp when polymorphism N86Y is present

DISCUSSION

Sample collection of this research was done in dry season, from April to July 2012. Transmission of malaria in dry season is low with the result a few number of sample could be collected. In addition, some isolates may fail to be amplified due to the low density of parasitemia. According to Scopel et al.,¹³ the low mean of parasitemia cause negative PCR result. Besides, sensitivity of malaria parasite to be detected by PCR was influenced by the method of sampling and storage. Improper blood storage cause DNA fragmentation, thus making it difficult for amplification.¹³

The isolates of *P. falciparum* examined in this study were found to carry multiple genetic polymorphisms associated with resistance to chloroquine. Although the molecular basis for the *P. falciparum* resistance to chloroquine remains uncertain, evidences indicate that resistance is multigenic.^{6,10}

The *pfprt* gene is located on chromosome 7 and it has been shown to associate with the inheritance of chloroquine resistance through genetic crossing.¹⁴ This gene encodes a 425-amino acid-long putative transporter protein that localizes to the digestive vacuole membrane of the parasite and effectively neutralize the drug via a mechanism that efflux chloroquine from the digestive vacuole and or pH regulation.¹⁵ Eight point mutations have been identified in *pfprt* gene, i.e. M74I, N75I, K76T, A220S, Q271E, N326S, I356T, and R371I, which

have been associated with chloroquine resistance.^{16,18} The K76T mutation is strongly associated with the chloroquine resistance phenotypes in field and clinical studies.¹⁵⁻¹⁷ Mutation in codon 76 have been found in CQR *P. falciparum* strains all over the world and become the principal determinant of CQR,^{16,18} although it was also present to a lesser frequency in chloroquine-sensitive strain.¹⁰ This evidence suggests that additional mutations in other genes are necessary for conferring CQR, or other mechanism of resistance also appear to be involved.^{10,19}

In addition the role of mutations in *pfmdr1* in the modulation of CQR was shown.²⁰ Mutations in the *pfmdr1* gene, i.e. N86Y, S1034C, N1042D, and D1246Y have been associated with CQR.²¹ The *pfmdr1* gene is a member of the ATP-binding cassette (ABC) transporter family that encodes Pgh-1 protein. The *pfmdr1* gene is located on chromosome 5 and may have a modulatory effect in parasite susceptibility to CQ.¹⁶ Although the 86Y allele is widespread in Asia and Africa, its association with CQR is unclear.²² Nevertheless, the role of *pfmdr1* in this regard could not be excluded.

Previous field-based studies in Indonesia have reported that the 76T polymorphism of *pfprt* is associated with CQR *in vivo* and *in vitro*, and the allele has the potential to be used as a marker for chloroquine treatment failure.^{6,17,23} Other studies in Indonesia have associated the 86Y allele of the *pfmdr1* gene to CQR both *in vivo* and *in vitro*.²⁴ Our interpretable findings showed that all *P. falciparum* isolates in South Sumatera carry both polymorphism in the *pfprt* and *pfmdr1* genes simultaneously. The high prevalence of the *pfprt* T76 allele found in this study is consistent with rates of 65% to 100% reported previously from different geographic regions.¹⁹ In western Indonesia, such as North Sumatera, Lampung, Central Java, East Kalimantan, all asymptomatic and mildly malaria patients were carrying polymorphism in both *pfprt* 76T and *pfmdr1* 86Y genes.⁶ Meanwhile in eastern Indonesia the situation was more varied. Northern Sulawesi had a resistant profile at these two codons, whereas southern Sulawesi had a lower frequency of *pfmdr1* 86Y polymorphism, but *pfprt* 76T was found in all parasite population.⁶ This situation was similar in Papua, Indonesia.⁶ Our present results may reflect the failure of treatment with the standard dose of chloroquine within the last few years in South Sumatera. This finding strengthen the previous researches, which stated that resistance to

mixtures containing ddH₂O 9 µL, Green go taq 10 µL (Promega USA), and a pair of primers. Five microliters of DNA was used as template in the first reaction and 2 µL of first round PCR product was used as template for secondary PCR. Positive (FCR3 Pf strain DNA) and negative (water) controls were used in all PCR. The primers and condition of PCR were as previously described by Duraisigh et al.¹²

Restriction fragment length polymorphism (RFLP)

Restriction enzyme *Aff* III and *Apo* I (New England Biolabs, Beverly, MA) were used to determine the presence of polymorphism N86Y *pfmdr1* and K76T *pfprt* gene. *Aff* III enzyme digested PCR product would show the presence of polymorphism at codon 86, while *Apo* I restriction enzyme would not cut amplicon when polymorphism 76T was present. Five microliters of each PCR product was digested with the restriction enzyme *Aff* III at 37°C and *Apo* I at 50°C for one hour. Digested products were electrophoresed on 1.5-3% agarose gels (Promega, USA) and visualized under UV transillumination after staining with ethidium bromide.

RESULTS

A total of 30 patients were enrolled in the study. Twenty five of them showed positive results in microscopic examination of blood smear stained with Giemsa. The mean age of the falciparum-infected persons was 27 years old (range 6-55 years old). Of these, 76% were male. Analysis of *pfmdr1* and *pfprt* gene PCR products indicated that mutant alleles of these genes have spread to all samples examined in these district (Table 1).

A 145-bp region surrounding the *pfprt* K76T mutation was amplified by PCR, and the mutation was detected using the *Apo* I restriction enzyme. *Apo* I digestion produces two fragments i.e. 125 bp and 20 bp in wild-type alleles, whereas the mutant alleles remain undigested. All isolates (25 of 25) of the amplified samples carried the 76T polymorphism, but there were two heterozygous cases, where the mutant was mixed with wild-type allele (76K) (Figure 1).

The *pfmdr1* N86Y mutation was similarly detected by RFLP analysis. A 372 bp region surrounding the mutation at position 86 was amplified by PCR and digested with the restriction enzyme *Aff* III. *Aff*

Table 1. Genotype profile of *P. falciparum* isolates from Lahat, Sekayu, Baturaja, and Palembang district

Isolate no.	<i>Pfmdr1</i> (86Y)	<i>Pfprt</i> (76T)
1.	Y	T
2.	Y	T
3.	Y	T
4.	Y	K/T
5.	Y	T
6.	Y	K/T
7.	Y	T
8.	Y	T
9.	Y	T
10.	Y	T
11.	Y	T
12.	Y	T
13.	Y	T
14.	Y	T
15.	Y	T
16.	Y	T
17.	Y	T
18.	-	T
19.	-	T
20.	Y	T
21.	Y	T
22.	Y	T
23.	-	T
24.	-	T
25.	-	T

K: wildtype (Lysine); T: mutant (Threonine); Y: mutant (Tyrosine); K/T: heterozygote mutation




Figure 1. RFLP for detecting T76 polymorphism. Sample no. 4 and 6 are heterozygous mutant alleles, others are homozygote mutant. Polymorphism K76T show a single 145 bp, whereas wild type allele digested by *Apo* I into 125 and 20 bp

Malaria *falciparum* is a serious health problem in society, especially in tropical country, and a global threat for the inhabitants of the earth. This situation is aggravated by the increasing antimalarial drug resistance. Chloroquine (CQ) has been used worldwide as a first line drug for acute malaria treatment.¹ Although the policy of malaria treatment in Indonesia has used artemisinin combination therapy (ACT) as the first line since 2001,² CQ remains the first line antimalarial agent in some regions in South Sumatera.³

P. falciparum resistance to chloroquine is a big problem and continuously develops. Resistance to antimalarial drugs adds the disease burden, increases the transmission, and causes epidemics.⁴ Resistance to chloroquine was first reported in East Kalimantan & Papua in 1975.⁵ Since that, chloroquine resistance (CQR) has spread and observed all over provinces in Indonesia.⁶

Molecular studies over the last few decades have identified some mutations in *P. falciparum* genes that are associated with CQR. Mutation in *Plasmodium falciparum* multidrug resistance 1 (*pfmdr1*), especially in kodon 86, where asparagin was changed into tyrosin, have been identified to modulate higher levels of CQR.^{6,7} However, mutation in *pfmdr1* alone is not enough to mediate CQR phenotype and that the trait is multigenic.⁸ A 76-Ser to Thr polymorphism in the *Plasmodium falciparum* chloroquine resistance transporter (*pfprt*) gene, which is located on chromosome 7, is known to be an important key of CQR phenotype.⁹ *pfmdr1* gene codes for P-glycoprotein homologues 1 (*Pgh 1*) and *pfprt* gene codes for transporter protein. Mutation in these genes causes chloroquine efflux into the cytoplasm and modification of acid degree, which have important roles in CQR.⁹

Epidemiologic study in all malaria endemic areas throughout the world have been conducted looking for polymorphisms in the genes and their relationships with treatment failure or resistance to chloroquine.^{6,10} *In vitro* and *in vivo* sensitivity tests to chloroquine in various malaria endemic area showed the existence of CQR and most of the resistant isolate carried mutant allele from those two genes.^{6,10}

The aim of this study is to complement the existing knowledge of *in vivo* & *in vitro* antimalarial drug responses by determining the extent of CQR associated gene polymorphisms in *P. falciparum*

isolates in South Sumatera. Identification of these mutations is expected to provide information about malaria treatment failures in South Sumatera.

METHODS

This study was carried out with the approval of the Ethics Committees at the Medical Faculty of Sriwijaya University (Palembang, Indonesia) No. 059/kepkrsmhfkunsri/2012.

Study sites

Three district hospitals from malaria-endemic area (Lahat, Baturaja, Sekayu) and one center of referral hospital Mohammad Hoesin Palembang in South Sumatra were selected for sample collection. Malaria in this region is mesoendemic with intense transmission between August and Desember. In this area, CQ is still used as first line antimalarial drug except in Baturaja and Sekayu.

Sample collection

Subjects were recruited from the local outpatient hospital. Criteria for participation in this study were age of 5 years or more and symptoms of malaria (e.g.: fever, chills, headache). Exclusion criteria were pregnancy, history of recent treatment with antimalarials, and severe or complicated malaria.

After obtaining informed consent, 5 mL of venous blood was drawn from each patient. Blood samples were collected in edetic acid (EDTA)-coated vacutainer tubes. Sample collection (all) was performed during April through July 2012. *P. falciparum*-infected samples as revealed by microscopic examination of a slide smear were used for DNA isolation. In addition, demographic data (age and sex) of all recruited subjects were noted.

Extraction of DNA

Parasite DNA was extracted from the blood samples using Chelex-100 ion exchanger (Bio-Rad USA) according to the procedure described previously.¹¹ The DNA was either used immediately for polymerase chain reaction (PCR) or stored at -20°C for later analysis.

Polymerase chain reaction amplification

Nested PCRs were performed for *pfprt* and *pfmdr1* genes. All reactions were carried out in 25 µL reaction

- analysis of *pfprt* and *pfmdr1* polymorphisms in *Plasmodium falciparum* isolates from Senegal. *Am J Trop Med Hyg.* 2002;66(5):474-80.
20. Foote SJ, Thompson JK, Cowman AF, Kemp DJ. Amplification of multidrug resistance gene in some chloroquine resistant isolates of *Plasmodium falciparum*. *Cell.* 1989; 57(6):921-30.
21. Foote SJ, Kyle DE, Martin RK, Oduola AMJ, Forsyth K, Kemp DJ, et al. Several alleles of the multidrug-resistance gene are closely linked to chloroquine resistance in *Plasmodium falciparum*. *Nature.* 1990; 345:255-8.
22. Basco LK, Ndounga M, Ngane VF, Soula G. Molecular epidemiology of malaria in Cameroon. XIV. *Plasmodium falciparum* chloroquine resistance transporter (PFCRT) gene sequences of isolates before and after chloroquine treatment. *Am J Trop Med Hyg.* 2002;67(4): 392-5.
23. Nagesha HS, Syafruddin D, Casey GJ, Susanti AI, Fryauff DJ, Reeder JC, et al. Mutations in the *pfmdr1*, *dhfr* and *dhps* genes of *Plasmodium falciparum* are associated with in-vivo drug resistance in West Papua, Indonesia. *Trans R Soc Trop Med Hyg.* 2001;95(1):43-9.
24. Sutanto I, Supriyanto S, Ruckert P, Purnomo, Maguire JD, Bangs MJ. Comparative efficacy of chloroquine and sulfadoxine-pyrimethamine for uncomplicated *Plasmodium falciparum* malaria and impact on gametocyte carriage rates in the East Nusatenggara province of Indonesia. *Am J Trop Med Hyg.* 2004;70(5):467-73.

chloroquine has spread to all malaria endemic areas in Indonesia,^{10,25} including South Sumatera.

It is generally accepted that *pfert* is the principal determinant of CQR. However, it is not possible to predict the degree of CQR based on *pfert* genotype alone or even in combination with *pfmdr1* genotype.¹⁶ It is clear from the data that parasite isolates with very low IC₅₀ levels indicating *in vitro* sensitivity to chloroquine usually carry the *pfert* T76 allele.¹⁹ Other recent studies revealed that many patients with apparently sensitive response to chloroquine therapy were infected with mutant parasites. It means that other factors, including host immunity, may have influence on clinical outcomes after administration of chloroquine. In some areas with high transmission, some patients seem to be able to clear their parasitemia even in the presence of the *pfert* K76T mutation.¹⁸

In conclusion, our results confirm that PCR-RFLP technique provide a simple and rapid method of detecting polymorphisms in genes that may predict CQR. Although the identification of the polymorphism in the *pfert* and *pfmdr1* genes provides a significant indicator of CQR, further studies are needed to determine the role of these polymorphisms in the *in vivo* and *in vitro* responses to drug treatment.

Acknowledgments

We thank to Directorate General of Higher Education Ministry of Education and Culture; Rektor Universitas Sriwijaya Prof. Dr. Badiah Perizade, MBA and Prof. Dr. Muhammad Said, MSc as Director of Unsri Research Institute for financial support.

Conflict of interest

This work was supported by Hibah Fundamental contract no. 004.c/UN9.3.1/PL/2012. The authors declare that this study is free of conflict of interest.

REFERENCES

1. World Health Organization. Guidelines for the treatment of malaria. Geneva: World Health Organization; 2006.
2. World Health Organization. World Malaria Report. Geneva: World Health Organization; 2010.
3. Dinas Kesehatan Provinsi Sumatera Selatan. Pencegahan penyakit dan penyebaran lingkungan. Profil Kesehatan Provinsi Sumatera Selatan. 2010. Indonesian.
4. Fairhurst RM, Nanyar GML, Breman JG, Hallet R, Vennerstorm JL, Duong S, et al. Artemisinin-

- resistant malaria: research challenges, opportunities, and public health implications. *Am J Trop Med Hyg.* 2012;87(2):231-41.
5. Ebisawa I, Fukuyama T. Chloroquine-resistant of *Plasmodium falciparum* malaria from West Irian and East Kalimantan. *Ann Trop Med Parasitol.* 1975;69(3):275-9.
6. Syafruddin D, Asih PB, Casey GJ, Maguire J, Baird JK, Nagesha HS, et al. Molecular epidemiology of *Plasmodium falciparum* resistance to antimalarial drugs in Indonesia. *Am J Trop Med Hyg.* 2005;72(2):174-81.
7. Sutar SK, Gupta B, Ranjit M, Kar SK, Das A. Sequence analysis of coding DNA fragments of *pfert* and *pfmdr1* genes in *Plasmodium falciparum* isolates from Odisha, India. *Mem Inst Oswaldo Cruz.* 2011;106(1): 78-84.
8. Wellems TE. Molecular genetics of drug resistance in *Plasmodium falciparum* malaria. *Parasitol Today.* 1991;7(5):110-2.
9. Carlton J, Fidock DA, Djimde A, Plowe CV, Baker J, Peters JM, et al. Conservation of a novel vacuolar transporter in *Plasmodium* species and its role in chloroquine resistance in *falciparum* but not *vivax* malaria. *Curr Opin Microbiol.* 2001;4(4):415-20.
10. Syafruddin D, Asih PBS, Aggarwal SL, Shankar A. Frequency distribution of antimalarial drug-resistant alleles among isolates of *Plasmodium falciparum* in Purworejo district, Central Java Province, Indonesia. *Am J Trop Med Hyg.* 2003;69(6):614-20.
11. Wooden J, Kyes S, Sibley CH. PCR and strain identification in *Plasmodium falciparum*. *Parasitol Today.* 1993;9(8):303-5.
12. Duraisingh MT, Drakeley CJ, Muller O, Bailey R, Snounou G, Targett GA, et al. Evidence for selection for the tyrosine-86 allele of the *pfmdr1* gene of *Plasmodium falciparum* by chloroquine and amodiaquine. *Parasitology.* 1997;114:205-11.
13. Scopel KK, Fontes CJ, Nunes AC, Horta MF, Braga EM. Low sensitivity of nested PCR using *Plasmodium* DNA extracted from stained thick blood smears: an epidemiological retrospective study among subjects with low parasitaemia in an endemic area of the Brazilian Amazon region. *Malar J.* 2004;3(8):1-6.
14. Wellems TE, Plowe CV. Chloroquine-resistant malaria. *J Infect Dis.* 2001;184(6):770-6.
15. Fidock DA, Nomura T, Talley AK, Cooper RA, Dzekunov SM, Ferdig MT, et al. Mutations in the *P. falciparum* digestive vacuole transmembrane protein PfCRT and evidence for their role in chloroquine resistance. *Mol Cell.* 2000;6(4):861-71.
16. Chaijaroenkul W, Ward SA, Mungthin M, Johnson D, Owen A, Bray PG, et al. Sequence and gene expression of chloroquine resistance transporter (*pfert*) in the association of *in vitro* drugs resistance of *Plasmodium falciparum*. *Malaria J.* 2011; 10(42):1-9.
17. Djimdé A, Doumbo OK, Cortese JF, Kayentao K, Doumbo S, Diouré Y, et al. A molecular marker for chloroquine resistant *falciparum* malaria. *N Engl J Med.* 2001;344(4):257-63.
18. Dorsey G, Kamya MR, Singh A, Rosenthal PJ. Polymorphisms in the *Plasmodium falciparum* *pfert* and *pfmdr1* genes and clinical response to chloroquine in Kampala, Uganda. *J Infect Dis.* 2001;183(9):1417-20.
19. Thomas SM, Ndir O, Dieng T, Mboup S, Wypij D, Maguire JH, et al. *In vitro* chloroquine susceptibility and PCR