

5. Belum-Antioxidant Activity of Methanol Extract of Halodule uninervis Seagrass the.

by Ace Baehaki

Submission date: 23-Oct-2018 12:09PM (UTC+0800)

Submission ID: 1025065473

File name: ivity_of_Methanol_Extract_of_Halodule_uninervis_Seagrass_the.pdf (688.47K)

Word count: 1589

Character count: 8737



Research Journal of Pharmaceutical, Biological and Chemical Sciences

Antioxidant Activity of Methanol Extract of *Halodule uninervis* Seagrass from the Coastal of Lampung, Indonesia.

Ace Baehaki*, Agus Supriadi, Muhammad Cahya Pratama.

Program Studi Teknologi Hasil Perikanan, Fakultas Pertanian Universitas Sriwijaya Indralaya Oganllir Sumatera Selatan, Indonesia.

ABSTRACT

The purpose of this study was to analyze phytochemistry contents and antioxidant activity of the methanol extracts from seagrass of *Halodule uninervis*. The samples were collected from the coastal of Lampung. Parameters of research included phytochemical content, DPPH scavenging activity and reducing power. The result showed content of phytochemical compounds of methanol extract seagrass are flavonoids, alkaloids, steroid and phenols. Antioxidant activity with DPPH method (IC₅₀) of *H.uninervis* was 1.575 ppm. The highest of reducing power of *H.uninervis* was 1.381.

Keywords :Seagrass, *Halodule uninervis*, antioxidant.

*Corresponding author



INTRODUCTION

Seagrasses are the marine flowering plants. Seagrasses such as *Amphibolis*, *Heterzostera*, *Phyllospadi*, *Posidonia*, *Pseudalthenia* and *Zostera* are mostly restricted to temperate seas and seven genera of seagrasses such as *Cymodocea*, *Enhalus*, *Halodule*, *Halophila*, *Syringodium*, *Thalasia* and *Thalassodendron* are distributed in tropical seas [1]. Sea grass biomass is used as human food especially by coastal populations [2].

Numerous seagrasses have been shown to have antibacterial activities. *Halophila stipulacea*, *Cymodocea serrulata* and *Halodule pinifolia* [3], *Enhalus acoroides* [4] and *Enhalus acoroides*, *Thalassia hemprichii*, *Halodule pinifolia*, *Syringodium isoetifolium*, and *Cymodocea rotundata* [5] have been reported to exhibit antibacterial activity. Moreover, preliminary data suggest that seagrasses could represent an interesting source of antilarvicidal [6] and antioxidant [7]. However, there is a little information regarding seagrass from *Halodule uninervis* and their antioxidative activity.

MATERIALS AND METHOD

Preparation of Sample

Sea grass of *Halodule uninervis* was collected from Coastal of Lampung and immediately brought to the laboratory in sterile plastic bags containing water to prevent evaporation. Seagrasses were washed thoroughly with water to remove extraneous materials and shade-dried for 10 days at room temperature until constant weight obtained. The dried Seagrasses were powdered and stored in refrigerator for future use.

Preparation of Sea grass extract

Seagrass powder were soaked in 2 L with methanol (1:4 w/v), and kept for 10 days in a shaker. The extraction was repeated thrice and pooled. The dry aqueous extracts were stored in a refrigerator until further analysis.

Phytochemical Screening of *Halodule uninervis*

Flavonoids

Test of flavonoids were determined by Wadood et al method [8]. A sample was heated with 10 ml of ethylacetate over a steam bath for 3 min. The mixture was filtered and 4 ml of the filtrate was shaken with 1 ml of dilute ammonia solution. A yellow colouration indicated the presence of flavonoid.

Alkaloids

Test of alkaloids were determined by Trease & Evan method [9].

Saponins

Test of saponin were determined by Wall et al method [10]. A sample was shaken with water in a test tube. Frothing which persists on warning confirmed the presence of saponins.

Steroids

Test of steroids were determined by Edeoga et al method [11]. 1 ml of acetic anhydride was added twice to 0.5 g of aqueous extract with 2 ml H₂SO₄. The colour changed from violet to blue or green in sample indicating the presence of steroids.

DPPH radical scavenging activity

DPPH (1,1-diphenyl-2-picrylhydrazyl) radical scavenging activity was determined by Hanani *et al.* method [12].

The antioxidant activity of each sample was expressed in percentage inhibition of free radicals which is calculated by the formula:

$$\% \text{ Inhibition} = \frac{\text{blanko absorbance} - \text{sample absorbance}}{\text{blanko absorbance}} \times 100\%$$

Concentration and barriers to samples the value of each plotted on the x axis and y. Obtained equation in the form $[y = b(x) + a]$ is used to find the value of IC (Inhibitory concentration) with a stated value of y is 50 and the value of x as IC50. The data obtained analyzed descriptively.

Reducing power

8 Reducing power was determined by Oyaiza method [13]. The methanol extract of *Halodule uninervis* was mixed with 2.5 ml of 0.2 M phosphate buffer (pH 6.6) and 2.5 ml of 1% potassium ferricyanide. The mixture was incubated at 50°C for 20 min. 2.5 ml of 10% trichoroacetic acid (TCA) was added to the mixture, followed by centrifugation at 3000 rpm for 10 min. The upper layer of solution (2.5 ml) was mixed with 2.5 ml of distilled water and 2.5 ml of 0.1% ferric chloride and the absorbance was read at 700 nm.

RESULT AND DISCUSSION

The phytochemical screening

As seen as Table 1 showed content of phytochemical compounds of methanol extract of seagrass were flavonoids, alkaloids, steriod, tannins and phenols.

Table 1. Phytochemical compound of methanol extract of *H.uninervis* Seagrass

Extract of	Flavonoid	Alkaloid	Saponin	Steroid
<i>Halodule uninervis</i>	+	++	-	++
(+) indicates present while (-) indicates absent				

Flavonoid compounds have been found in methanol extract of *H.uninervis* seagrasses. Earlier study by Anwariyah [14] reported flavonoids have been found in seagrasses *C. rotundata*. Further screening of seagrass of *H.uninervis* showed the result of assayed of saponin compounds from seagrass of *H.uninervis* have not been found, due to the extracts not form foam. Saponin compounds also have been found in methanol extract of *H.uninervis* seagrasses. Saponin is a strong active compounds and cause foam when homogenized with water. Saponins are glycosides triterpena and sterols that have been detected in more than 90 tribes plants [15]. Steroids is typical of higher plants. We also detected a number of steroids in the methanol extract of *H.uninervis*, in a previous study of seagrass, steroid detected in extracts of *C. rotundata* with methanol, n-hexane and ethyl acetat [14]

DPPH radical scavenging activity

Antioxidant activities of extract of *Halodule uninervis* depicted in Figure 1. Increasing of concentration of *Halodule uninervis* indicates an increase in DPPH radical scavenging activity. The stable organic radical DPPH has been widely used in plant extracts [16], and foods[17], etc. The method is based on the reduction of alcoholic DPPH solutions at 517 nm in the presence of an hydrogen donating antioxidant (AH) due to the formation of the non-radical form DPPH-H.




Figure 1. Scavenging radical DPPH activity of methanol extract of *H.uninervis* Seagrass

6

The IC₅₀ of extract was 1.575 ppm. Antioxidants react with DPPH, reducing a number of DPPH molecules equal to the number of their available hydroxyl groups. Therefore, the absorption at 515 nm was proportional to the amount of residual DPPH [18]. Free radical scavenging activity of extract of *Halodule uninervis* could be due to their higher content of flavonoids components. Flavonoids have been interrupt the propagation of autoxidation by donating hydrogen atom of some hydroxyl (OH) bases that are attached outside the benzene rings, resulting in the formation of stable free radical [19]. These groups of polyphenolic compounds are very important in plant because they make their defense mechanisms [20].

Reducing power

Reducing power of extract of *Halodule uninervis* depicted in Figure 2. Increasing of concentration of *Halodule uninervis* indicates an increase in reducing power




Figure 2. Reduction power of methanol extract of *H.uninervis* Seagrass

3

The reducing capacity of the extract is another significant indicator of antioxidant activity. In the reducing power assay, the presence of antioxidants in the sample would result in the reduction of Fe³⁺ to Fe²⁺ by donating an electron. Reducing power of a methanol extract of *H.uninervis* is probably due to the presence of group hydroxyl in the phenol compounds that can serve as an electron donor. Therefore, antioxidants are considered reducers and inactivating oxidants

REFERENCES

- [1] Vijayakumar S, Manakha M 1, Anderson A, Lingakumar K. Int J Microbiol & Parasit 2014; 1(1):1-10
- [2] Hemminga MA, Duarte CM. Seagrass ecology. Cambridge University Press, New York, 2000, pp.298.



- [3] Kannan RRR, Arumugam R, Anantharaman P. Asian Pacific Journal of Tropical Medicine 2010; 890-893
- [4] Qi S, Zhang S, Qian P, Wang B. Botanica Marina 2008;51: 1-7
- [5] Kannan RRR, Arumugam R, Iyapparaj P, Thangaradjou T, Anantharaman P. Food Chem 2013; 136 1484–1489
- [6] Ali MS, Ravikumar S, Beu JM. Int J of Pharm Bio Sci 2013; 4(2): (P) 62 - 67
- [7] Ramah S, Etwarising L, Auckloo N, Gopeechund A, Bhagooli R, Bahorun T. Internet Medical Update 2014; 9(1):27-37
- [8] Wadood A, Ghufuran M, Jamal M, Naeem SB, Khan A. Biochem Anal Biochem 2013; 2: 144.
- [9] Trease GE, Evan WC. A Textbook of pharmacognosy 11th edition. Bailliere Tindall, London, 1978, pp.530
- [10] Wall ME, Krider MM, Krewson CR, Wilaman JJ, Correll S, Gentry HS. Agr. Res service circ. Aic1954; 363,17
- [11] Edeoga HO, Okwu DE, Mbabie BO. Afr. J. Biotechnol 2005;4, 685-688
- [12] Hanani E, Moneim B, Sekarini R. Magazine Pharm. Sci 2005; 2: 127-133.
- [13] Oyaizu, M. Japanese J. Nutr 1986; 44: 307–315.
- [14] Anwariyah, R. 2011. Skripsi. Bogor Agricultural University (IPB). Bogor. pp.79.
- [15] Suradikusumah, E. Plant Chemistry. Inter-University Centers of Biological Sciences, Bogor Agricultural University (IPB), Bogor, 1989. pp.166
- [16] Duh PD, Yen GC. Food Chem 1997; 60:639-645
- [17] Yamaguchi T, Takamura H, Matoba T, Terao J. Biosci Biotechnol Biochem 1998; 62:1201-1204.
- [18] Xu J, Chen S, Hu Q. Food Chem 2005; 91(1):79-83.
- [19] Bahramikia S, Ardestani A, Yazdanparast R. Food Chem 2009; 115(1):37–42.
- [20] Wallace G, Fry SC. Int Rev Cytol 1994; 151(2):229-267.

5. Belum-Antioxidant Activity of Methanol Extract of Halodule uninervis Seagrass the.

ORIGINALITY REPORT

23%

SIMILARITY INDEX

%

INTERNET SOURCES

%

PUBLICATIONS

23%

STUDENT PAPERS

PRIMARY SOURCES

1

Submitted to Northern Caribbean University

Student Paper

3%

2

Submitted to Mahidol University

Student Paper

3%

3

Submitted to Universiti Malaysia Perlis

Student Paper

3%

4

Submitted to University of Kufa

Student Paper

3%

5

Submitted to Federal University of Technology

Student Paper

3%

6

Submitted to Universiti Sains Malaysia

Student Paper

2%

7

Submitted to Higher Education Commission
Pakistan

Student Paper

2%

8

Submitted to University of Zululand

Student Paper

2%

Exclude quotes On Exclude matches < 15 words
Exclude bibliography On

PENJELASAN HASIL SIMILARITY DI ATAS:

1. Hasil uji kemiripan (similarity) artikel yang berjudul: Antioxidant Activity of Methanol Extract of *Halodule uninervis* Seagrass from the Coastal of Lampung, Indonesia terhadap dokumen (primary source) menggunakan program turnitin, hasil kemiripan yang paling tinggi adalah 3%.
2. Menurut pedoman operasional penilaian angka kredit kenaikan jabatan akademik/pangkat dosen tahun 2019 Dirjen Sumberdaya Iptek dan Dikti Kemenristekdikti halaman 25, bila hasil uji kemiripan melebihi 25% (duapuluh lima persen) terhadap 1 (satu) dokumen/primary source (tidak termasuk daftar pustaka, kemiripan kalimat yang kurang dari 3% (tiga persen)), maka peer review secara substansi harus memberikan pendapat ada tidaknya indikasi plagiasi.
3. Berdasarkan pedoman penilaian angka kredit kenaikan jabatan akademik/pangkat tahun 2019, hasil uji kemiripan artikel ini tidak terindikasi plagiasi.