

intern symposium2011

by Melki Melki

Submission date: 25-Jan-2020 10:59AM (UTC+0700)

Submission ID: 1246158637

File name: Melki_2011_International_symposium.doc (514K)

Word count: 2407

Character count: 12396

18
**BIO-POTENTIALS ACTIVITY OF *Sonneratia caseolaris* (MANGROVE)
EXTRACT AS ANTIBACTERIAL COLLECTED FROM SOUTH SUMATERA**

Melki (1), A Zaenal Mustopa (2), Hefni Effendi (3)

17

¹ (M.Si) (Department of Marine Science, Sriwijaya University) – (Jalan Raya Palembang Prabumulih Km. 32 Inderalaya, 30662) – (South Sumatera-Indonesia) – (Telp/fax. +62711581118) – (www.unsri.ac.id) – (melki@mipa.unsri.ac.id)

16

² (M.Si) (Research Center for Biotechnology, Indonesian Institut of Sciences) – (Jalan Raya Bogor Km 46, Cibinong, 16911) – (Cibonong-Bogor, Indonesia) – (Telp. +6221-8754587, Fax. +6221-8754588) – (www.biotech.lipi.go.id) – (azmustopa@yahoo.com)

10

³ (Dr.rer.nat) (Department of Aquatic Resources Management, Faculty of Fisheries and Marine Science, Bogor Agricultural University) – (Kampus IPB Darmaga, Bogor 16680) – (Bogor, Indonesia) – (Telp. +622518621262, Fax. +62251 8622134) – (www.ipb.ac.id) – (hefni_effendi@yahoo.com)

Summary

19

Crude extracts of four mangrove species (leaf, fruit, bark and root), i.e. *Avicennia alba*, *A. marina*, *Rhizophora mucronata*, and *Sonneratia caseolaris* collected from Teluk Payo, Banyuasin, South Sumatera was extracted in methanol, ethyl acetate, hexane and tested for antibacterial (*Escherichia coli* and *Staphylococcus aureus* pathogen), and brine shrimp cytotoxic assay. The highest activity recorded was methanol extract of *S. caseolaris* in *E. coli* isolates (18 mm inhibition) and in *S. aureus* isolates (19 mm inhibition), exhibiting relatively high biopotency. Brine Shrimp Lethality Test showed that leaf of *S. caseolaris* methanol extract was not toxic to *Artemia salina*. The high bioactive mangrove extract evaluated further by HPLC showed that mangrove extracts likely contains flavonoid.

Keywords: Extract of *Sonneratia caseolaris*, antibacterial, BSLT, column chromatography, TLC, and HPLC

Introduction

4

Microorganisms have potential to cause ² human diseases. Most of the time viruses, bacteria and fungi act as major pathogenic organisms. The discovery of antibiotics in the early twentieth century provided an increasingly important tool to combat bacterial diseases. As antibiotics are increasingly used and misused, the bacterial strains become resistant to antibiotic ³s rapidly. Therefore, antibacterial activity of medicinal plants is very important since vast number of medicinal plants have been used for centuries as remedies for human diseases. Among them extracts from different parts of mangroves and mangrove associates are widely used throughout the world. For instance, stem of *Avicennia marina* is used for ulcers and bark of *Bruguiera sexangula* is used for antitumors. Mangrove and mangrove associates contain biologically active antiviral, antibacterial and antifungal. They provide a rich source of steroids, triterpenes, saponins, flavonoids, alkaloids and tannins [1]. Therefore, it is worth to screen mangrove plants for the presence of new antibacterial compounds to combat the normal pathogenic bacterial strains and hospital acquired antibiotic resistant bacterial strains.

9

Mangroves are one of the easiest tropical forest types to generate. They have the ability to grow where no other vascular plants can. The mangroves exist under stressful conditions such as violent environments, high concentration of moisture, high and low tides of water, and abundant living microorganisms and insects. They thrive in a very peculiar environment and serve as a bridging ecosystem between freshwater and marine systems. They possess an unusual morphology ¹ and physiognomy and the path of photosynthesis in mangroves is different from other glycophytes. They possess modifications to establish water and salt economy. There are modifications or alterations in other physiological processes such as carbohydrate metabolism or polyphenol synthesis and due to these reasons, they may have chemical compounds, which protect them from these destructive elements [2].

Mangroves from the coast of Teluk Payo, South Sumatra province were collected. Their extract of leaf, fruit, bark and root was scrutinized to determine their bioactive using *Escherichia coli* and *Staphylococcus aureus* and brine shrimp cytotoxic as target organism.

Materials and Methods

Collection and Extraction of Mangrove Bioactive

Four species of mangroves i.e. *Avicenna marina*, *A. alba*, *Rhizophora mucronata* and *Sonneratia caseolaris* were collected and identified from the mangrove forest in Teluk Payo, South Sumatera during December 2009 (Figure 1). Prior to the extraction, leaves, fruits, barks and roots of respective species were cleaned, shade dried in order to prevent photolysis and thermal degradation, then chopped into small pieces and ground coarsely in a mechanical grinder.




Figure 1. Map showing the study area, Teluk Payo, South Sumatra (Indonesia)

Extraction of Bioactive

Around 100 g of powdered mangrove material was extracted for 3 x 24 hour using methanol 80%, ethyl acetate 80% and hexane 80% as much 250 (12). The extracts were filtered using Whatman no. 1 filter paper. The fraction was evaporated at rotary evaporator at 40 - 50°C, then collected in air-tight plastic vials and stored in the refrigerator for further studies.

Bioassays

Antimicrobial assay was carried out as described by Abesinghe and Wanigatunge [3] against *Escherichia coli* NBRC 13276 and *Staphylococcus aureus* NBRC 14237 pathogen. The cytotoxic activity of mangrove extracts was tested against freshly hatched free-swimming nauplii of *Artemia salina* Leach. The assay system was prepared with 3 ml of filtered seawater containing chosen concentration of mangroves extract in cavity blocks (embryo cup) and 10 nauplii each was transferred in experimental, vehicle control and negative control wells. Invariably the concentration of the experimental systems was determined on the basis of exploratory experiments. The percentage of mortality was determined by comparing the mean surviving larvae of the test and control tubes. The LC₅₀ value was determined using probit scale [4].

Column Chromatography and Thin Layer Chromatography (TLC) 5

The mangroves extract (2 ml) was loaded on a silica gel 60 Fe₂₅₄ (Merck) column packed with chloroform and eluted with chloroform and methanol (9:1 to 1:9) to yield fractions. The potential fractions were then examined by TLC 5 on 25 TLC alluminium silica gel 60 Fe₂₅₄ (Merck) using same solvent with column chromatography. Individual fractions were collected and tested for bioactivity.

High Performance Liquid Chromatography (HPLC)

Mangrove extracts derived from column chromatography were injected to HPLC KNAUER, column Eurospher100-5C₁₈ (4.6 x 150 mm, 5 µm), injection volume 20 µl, PDA detector with 200-400 nm, flow rate of gradient 1 ml/minute, methanol:water (v/v) (at 0 minute 0% water, at 22 minute 100% methanol, at 30 minute 100% methanol, at 33 minute 100% water, at 40 minute 100% water).

Results and Discussion

Extraction of Component Bioactive

Methanol solvent produced more weight extract percentage than that of ethyl acetate and hexane solvent (**Table 1**).

Table 1. Mangrove Extracted with MeOH, EtOAc, and Hexane

Species of Mangrove	Part	Original weight (g)	Weight Extract (g)			Crude Extract (% b/v)		
			MeOH	EtOAc	Hexane	MeOH	EtOAc	Hexane
<i>A. marina</i>	Leaf	100	7.71	6.80	3.34	7.71	6.80	3.34
	Bark	100	4.22	4.08	2.21	4.22	4.08	2.21
	Fruit	100	11.8	10.89	2.23	11.80	10.89	2.23
	Root	100	5.22	4.72	2.56	5.22	4.72	2.56
<i>A. alba</i>	Leaf	100	7.93	7.23	4.23	7.93	7.23	4.23
	Bark	100	9.34	8.67	4.21	9.34	8.67	4.21
	Fruit	100	10.31	9.98	4.22	10.31	9.98	4.22
	Root	100	8.30	7.56	3.67	8.30	7.56	3.67
<i>R. mucronata</i>	Leaf	100	10.89	9.79	2.58	10.89	9.79	2.58
	Bark	100	6.73	5.56	2.78	6.73	5.56	2.78
	Root	100	6.75	6.34	3.53	6.75	6.34	3.53
<i>S. caseolaris</i>	Leaf	100	8.67	6.65	3.45	8.67	6.65	3.45
	Bark	100	6.89	6.34	3.12	6.89	6.34	3.12
	Fruit	100	8.90	7.54	3.45	8.90	7.54	3.45
	Root	100	9.78	8.67	4.34	9.78	8.67	4.34

Antibacterial Activity

Crude extracts of mangrove (leaf, fruit, bark and root) had the ability to inhibit bacterial growth of *E. coli* and *S. aureus*. The invitro antibacterial activity revealed that methanol extract of mangroves had remarkable antibacterial activity. Among four species tested, methanol extract of the leaf *S. caseolaris* exhibited wide spectrum of activity which suppress the growth of all tested bacteria, produced a mean zones of inhibition in *E. coli* culture (18 mm) and in *S. aureus* culture (19 mm) (**Table 2** and **Table 3**).

Table 2. Antibacterial activity screening of mangrove extracts towards *E. coli*

Species of Mangroves	Part	Diameter of inhibition (mm)*							
		MeOH (20µl/5µg)	EtOAc (20µl/5µg)	Hexane (20µl/5µg)	A (-)	B (-)	C (-)	D (+)	E (+)
<i>A.marina</i>	Leaf	10	8	0	0	0	0	10	15
	Bark	12	0	0	0	0	0	10	15
	Fruit	16	12	0	0	0	0	10	15
	Root	9	9	0	0	0	0	10	15
<i>A.alba</i>	Leaf	9	7	0	0	0	0	10	15
	Bark	10	8	0	0	0	0	10	15
	Fruit	13	9	0	0	0	0	10	15
	Root	9	8	0	0	0	0	10	15
<i>R.mucronata</i>	Leaf	11	9	0	0	0	0	10	15
	Bark	11	9	0	0	0	0	10	15
	Root	9	8	0	0	0	0	10	15
<i>S.caseolaris</i>	Leaf	18	17	0	0	0	0	10	15
	Bark	12	11	0	0	0	0	10	15
	Fruit	14	11	0	0	0	0	10	15
	Root	9	8	0	0	0	0	10	15

info: * optimization hour observation at the 16th, A (-) MeOH, B (-) EtOAc, C (-) Hexane, D (+) Penicillin 10 µg, E (+) Chloramphenicol 30 µg

Table 3. Antibacterial activity screening of mangrove extract towards *S. aureus*

Species of Mangroves	Part	Diameter of inhibition (mm)*							
		MeOH (20µl/5µg)	EtOAc (20µl/5µg)	Hexane (20µl/5µg)	A (-)	B (-)	C (-)	D (+)	E (+)
<i>A.marina</i>	Leaf	10	7	0	0	0	0	10	15
	Bark	12	11	0	0	0	0	10	15
	Fruit	12	9	0	0	0	0	10	15
	Root	7	10	0	0	0	0	10	15
<i>A.alba</i>	Leaf	9	7	0	0	0	0	10	15
	Bark	8	7	0	0	0	0	10	15
	Fruit	10	8	0	0	0	0	10	15
	Root	9	8	0	0	0	0	10	15
<i>R.mucronata</i>	Leaf	10	8	0	0	0	0	10	15
	Bark	10	8	0	0	0	0	10	15
	Root	9	8	0	0	0	0	10	15
<i>S.caseolaris</i>	Leaf	19	16	0	0	0	0	10	15
	Bark	16	11	0	0	0	0	10	15
	Fruit	11	11	0	0	0	0	10	15
	Root	10	8	0	0	0	0	10	15

info: * optimization hour observation at the 16th, A (-) MeOH, B (-) EtOAc, C (-) Hexane, D (+) Penicillin 10 µg, E (+) Chloramphenicol 30 µg

The result of bioactivity tests showed that mangrove extract bioactive compounds have strong power. According to Davidstout [5], that the power of antibacterial is pointed out by growth inhibition zone. The growth inhibition of >20 mm means very strong, inhibition zone of 10-20 mm means strong, inhibition zone of 5-10 mm means medium, and inhibition zone of <5 mm means weak.

The highest activity recorded was methanol extract of the leaf *S. caseolaris* inhibiting the growth of *E. coli* isolates (18 mm inhibition) and the growth of *S. aureus* isolates (19 mm inhibition).

The difference between the antibacterial activities of mangrove extract could be due to the type and quantity of antimicrobial substances present in each form.

Brine Shrimp Assay

15

The brine shrimp assay is considered as a reliable indicator for the preliminary assessment of toxicity [6]. This assay is widely employed in the screening process of botanical for the isolation of bioactive metabolites.

The extract of four mangrove species showed different mortality rate at different concentrations (Table 4). The mortality rate increased with the increase of concentration of each sample. The crude extracts of the leaf *S. caseolaris* indicated the highest LC₅₀ value of 34914.03 µg/ml meaning not toxic.

Table 4. *Artemia* cytotoxicity profile of mangrove extracts

Concentration µg/ml	Percentage of Mortality			
	Fruit of <i>A. marina</i>	Fruit of <i>A. alba</i>	Leaf of <i>R. mucronata</i>	Leaf of <i>S. caseolaris</i>
0 (Control)	0.00	0.00	0.00	0.00
10	3.33	3.33	3.33	13.33
100	3.33	6.67	10.00	16.67
200	3.33	30.00	13.33	23.33
500	6.67	30.00	23.33	33.33
1000	10.00	36.67	33.33	30.00
LC ₅₀ (µg/ml)	532108.08	6714.288	1043.019	34914.03
Category of toxicity [7]	Not toxic	Not toxic	Not toxic	Not toxic

Purification with Column Chromatography

First step of identification of bioactive compound of mangroves plant is by usage of column chromatography. Solvent applied is chloroform:methanol (9:1 to 1:9). For extract of the leaf *S. caseolaris* as much 20 fractions (yield extract) are produced. Then all fractions were antibacterial tested (*S. aureus* and *E. coli*).

Purification with TLC

Fraction showing bioactivity then underwent thin layer chromatography TLC. Its Rf point (Retardation fraction) was measured. Extract of the leaf *S. caseolaris* eluted by chloroform:methanol (9:1 to 1:9) showed Rf points 0.78. (Figure 2). Spot on TLC is taken further and tested by antibacterial (*S. aureus* and *E. coli*). The growth inhibition was 7 mm.




Figure 2. Profile of bioactive fraction of *S. caseolaris* leaf on TLC

HPLC Characteristic of Bioactive Substance

Analysis of HPLC with detector photodiode array (PDA) ($\lambda = 200 - 400$ nm) showed that the dominant peak with its retention time and wavelength characteristic was similar with those of the standard compound characteristic stored in HPLC data base.

HPLC analysis of the leaf *S. caseolaris* showed have 4 peak dominant at retention time (Rt) 14,50 ($\lambda = 234$ nm), Rt 15,92 ($\lambda = 258$ nm), Rt 19,68 ($\lambda = 230$ nm), and Rt 23,13 ($\lambda = 223$ nm and 277 nm) (Figure 3).




Figure 3. HPLC Analysis of leaf *S. caseolaris* extract

Conclusions

Among four mangrove species screened, the broadest activity was showed by *S. caseolaris* extract, therefore this mangrove plant might posses potential source for further study of its bioactive compounds for the purpose of biopharmaceuticals.

Acknowledgements

We wish to thank to all parties supporting this research.

References

- [1] Bandaranayake, W. M. Traditional and medicinal uses of mangroves. *Mangroves and Salt Marshes*, 2: 133-148. 1998.
- [2] Bandaranayake, W. M. Bioactivities, bioactive compounds and chemical constituents of mangrove plants. *Wetlands Ecology and Management* 10: 421-452. 2002.
- [3] Abeysinghe, P, D and Wanigatunge, R. P. Evaluation of antibacterial activity of different mangrove plant extracts. *Ruhuna Journal of Science* 1:104-112. 2006.
- [4] Wardlaw, A.C. Practical statistics for experimental biologists. John Wiley and Sons, Chichester. 1985.
- [5] Davidstout. Disc plate method of microbiological antibiotic assay. *Journal of Microbiology* 4: 659-665. 1971.
- [6] Cyrus, W.G., Daniel, G.W., Nanyangi, M.O., Njonje, F.K. and Mbaria, J.M. Antibacterial and cytotoxic activity of Kenyan medicinal plants, *Memórias do Instituto Oswaldo Cruz*, 103(7): 650-652. 2008.
- [7] Meyer, B. N. Brine Shrimp: a convinient general bioassay for active plant constituents. *Plant Medica* 45: 31-34. 1982.
- [8] Haque, M. E., Islam, M. N., Rahman, M. H., Mohamad, A. U., Antimicrobial and Cytotoxic Activities of the Crude Extracts and Isolated Compounds of *Xylocarpus molluccensis*. *Dhaka Univ. J. Pharm. Sci.* 2: 109-112. 2007.
- [9] Kitamura, S., Anwar, C., Chaniago, A and Baba, S. *Handbook of Mangrove in Indonesia*. ISME. Japan. 1997.
- [10] Wu, J., Xiao, Q., Xu, J., Li, M.Y., Pana, J.Y. and Yang, M. Natural products from true mangrove flora: source, chemistry and bioactivities. *Natural Product Report*, 25: 955-981. 2008.

intern symposium2011

ORIGINALITY REPORT



PRIMARY SOURCES

- | | | |
|---|--|----|
| 1 | Gehan M. Abou-Elela, Nermene A. El-Sersy, Mohamed A. El-Shenawy, Hanan Abd-Elnabi, Hassan A.H. Ibrahim. "Bio-Control of <i>Vibrio fluvialis</i> in Aquaculture by Mangrove (<i>Avicennia marina</i>) Seeds Extracts", Research Journal of Microbiology, 2009 | 2% |
| | Publication | |
| 2 | ajpcrjournal.com
Internet Source | 1% |
| 3 | gujaratdesertecology.com
Internet Source | 1% |
| 4 | www.ijppsjournal.com
Internet Source | 1% |
| 5 | thalassas.webs.uvigo.es
Internet Source | 1% |
| 6 | www.formatex.info
Internet Source | 1% |
| 7 | banglajol.info
Internet Source | 1% |
-

- 8 Pritam N. Dube, Shweta S. Bule, Yogesh V. Ushir, Manoj R. Kumbhare, Pravin R. Dighe. "Synthesis of novel 5-methyl pyrazol-3-one derivatives and their in vitro cytotoxic evaluation", Medicinal Chemistry Research, 2014
Publication
-
- 9 www.grjournals.com 1 %
Internet Source
-
- 10 hrcak.srce.hr 1 %
Internet Source
-
- 11 Submitted to Institute of Graduate Studies, UiTM 1 %
Student Paper
-
- 12 nopr.niscair.res.in 1 %
Internet Source
-
- 13 www.apjtb.com 1 %
Internet Source
-
- 14 pharmacologyonline.silae.it 1 %
Internet Source
-
- 15 Submitted to Griffith University 1 %
Student Paper
-
- 16 journal.ipb.ac.id 1 %
Internet Source
-
- 17 M. S. Imanudin, E. Armanto, R. H. Susant, S. M. Bernas. "The study water table fluctuation in 1 %

tidal lowland for developing agricultural water management strategies: (A case study for corn cultivation after rice)", 2010 International Conference on Chemistry and Chemical Engineering, 2010

Publication

18 ejournal2.unsri.ac.id 1 %

Internet Source

19 Patra, Jayanta Kumar, Sushanto Gouda, Nabin Kumar Dhal, and Hrudayanath Thatoi. "Comparative Evaluation of Antibacterial Properties and Phytochemical Profile of Selected Mangrove Plants from Bhitarkanika, India", Journal of Biologically Active Products from Nature, 2014.

Publication

Exclude quotes

On

Exclude matches

< 1%

Exclude bibliography

On