



Vol 11, No 1 (2016)

Table of Contents

Articles

[Antioxidant Potential Of Lansium Domesticum Corr. Seed Extract In White Male Rat \(Rattus Novergicus\) Induced By Alcohol](#)

PDF
1-8

Subandrate Subandrate, Sadakata Sinulingga, Sri Wahyuni, M Fakhri Altayan, Fatmawati Fatmawati

| DOI : [10.20884/1.jm.2016.11.1.189](https://doi.org/10.20884/1.jm.2016.11.1.189) | This article has been viewed: **1725 times** |

[Bacterial Cellulose From Rice Waste Water With Addition Chitosan, Glycerol, And Silver Nanoparticle](#)

PDF
9-24

Eli Rohaeti, Endang WLFX, Anna Rakhmawati

| DOI : [10.20884/1.jm.2016.11.1.190](https://doi.org/10.20884/1.jm.2016.11.1.190) | This article has been viewed: **1495 times** |

[Morphology-Controlled Synthesis Of SrTiO3 Nanocube By Capping Agent-Assisted Solvothermal Method](#)

PDF
25-32

Yulia Eka Putri, Alvionita Alvionita, Rini Rahma Yanti, Diana Vanda Wellia

| DOI : [10.20884/1.jm.2016.11.1.191](https://doi.org/10.20884/1.jm.2016.11.1.191) | This article has been viewed: **643 times** |

[Effect Of Non-metal Elements \(C, N, S\) As Anionic Dopants On Electronic Structure Of TiO2-Anatase By Density-Functional Theory Approach](#)

PDF
33-41

Hari Sutrisno

| DOI : [10.20884/1.jm.2016.11.1.192](https://doi.org/10.20884/1.jm.2016.11.1.192) | This article has been viewed: **336 times** |

[Fractionation And Environmental Risk Of Trace Metals In Surface Sediment Of The East China Sea By Modified BCR Sequential Extraction Method](#)

PDF
42-52

Roy Andreas, Jing Zhang

| DOI : [10.20884/1.jm.2016.11.1.193](https://doi.org/10.20884/1.jm.2016.11.1.193) | This article has been viewed: **622 times** |

[Oxidation Of Cyclohexane To Cyclohexanol And Cyclohexanone Using H4\[α-SiW12O40\]/Zr As Catalyst](#)

PDF
53-60

Aldes Lesbani, Menik Setyowati, Risfidian Mohadi, Dedi Rohendi

| DOI : [10.20884/1.jm.2016.11.1.194](https://doi.org/10.20884/1.jm.2016.11.1.194) | This article has been viewed: **934 times** |

[Electro-Synthetic Optimization of Host Material Based on MIL-100\(Fe\)](#)

PDF

[Guide for Authors](#)

[View Articles](#)

[Submit your paper](#)

[Editorial Board](#)

[Focus and Scope](#)

[Publication Ethics](#)

[Peer Review Process](#)

[Plagiarism Policy](#)

Abstract Indexed in:

Scopus[®]



DOAJ DIRECTORY OF OPEN ACCESS JOURNALS



Google scholar

Witri Wahyu Lestari, Joni Hartono, Marisa Adreane, Khoirina Dwi Nugrahaningtyas, Candra Purnawan, Sentot Budi Rahardjo
| DOI : [10.20884/1.jm.2016.11.1.195](https://doi.org/10.20884/1.jm.2016.11.1.195) | This article has been viewed: **1110 times** |

[Identification of Differentially Expressed Protein from Electrical Stunning of Broiler Chickens Meat Protein](#)

Sandra Hermanto, Maya Ina Sholaikah, Sri Suci Mulyani
| DOI : [10.20884/1.jm.2016.11.1.196](https://doi.org/10.20884/1.jm.2016.11.1.196) | This article has been viewed: **479 times** |

[Histochemical Changes Liver and Kidney of Mice Exposed to Mercury and Recovery with Nanogold](#)

Titik Taufikurohmah, I Gusti Made Sanjaya, Afaf Baktir, Achmad Syahrani
| DOI : [10.20884/1.jm.2016.11.1.197](https://doi.org/10.20884/1.jm.2016.11.1.197) | This article has been viewed: **865 times** |

[Physical Damages of Wood Fiber in Acacia Mangium due to Biopulping Treatment](#)

Ridwan Yahya, Mucharromah Mucharromah, Devi Silsia, Septiana Septiana
| DOI : [10.20884/1.jm.2016.11.1.198](https://doi.org/10.20884/1.jm.2016.11.1.198) | This article has been viewed: **395 times** |

[Identification of Secondary Metabolites Compounds and Antibacterial Activities on The Extract of Soursop Leaf](#)

Dian Riana Ningsih, Zufahair Zufahair, Dwi Kartika
| DOI : [10.20884/1.jm.2016.11.1.199](https://doi.org/10.20884/1.jm.2016.11.1.199) | This article has been viewed: **1673 times** |

[Optimizing Conditions to Cholesterol Adsorbed with Carboxymethyl Chitosan](#)

Mardiyah Kurniasih, Dwi Kartika, Riyanti Riyanti
| DOI : [10.20884/1.jm.2016.11.1.200](https://doi.org/10.20884/1.jm.2016.11.1.200) | This article has been viewed: **425 times** |

[Secondary Metabolite from Endophytic Fungi Aspergillus Sp. The Leave Of Kunyit Putih \(Curcuma zedoaria \(Berg\) Roscoe\)](#)

Muharni Muharni, Fitriya Fitriya, Widia Purwaningrum, Ahmad Yogi Nugraha
| DOI : [10.20884/1.jm.2016.11.1.201](https://doi.org/10.20884/1.jm.2016.11.1.201) | This article has been viewed: **932 times** |

[Extraction of Cellulose from Kepok Banana Peel \(Musa parasidiaca L.\) for Adsorption Procion Dye](#)

Poedji Loekitowati Hariani, Fahma Riyanti, Riski Dita Asmara
| DOI : [10.20884/1.jm.2016.11.1.202](https://doi.org/10.20884/1.jm.2016.11.1.202) | This article has been viewed: **820 times** |

[Quantitative Structure-Activity Relationship Analysis of Xanthone Derivates as Cytotoxic Agents in Liver Cancer Cell Line HepG2](#)

Isnatin Miladiyah, Iqmal Tahir, Jumina Jumina, Sofia Mubarika, Mustofa Mustofa
| DOI : [10.20884/1.jm.2016.11.1.203](https://doi.org/10.20884/1.jm.2016.11.1.203) | This article has been viewed: **885 times** |

61-70



[PDF](#)

71-79



[Detail Statistic](#)

[PDF](#)

80-91



USER

Username

Password

Remember me



[PDF](#)

92-100

[PDF](#)

101-111

[PDF](#)

112-124

[PDF](#)

125-134

[PDF](#)

135-142

[PDF](#)

143-157



Molekul

**Jurnal Ilmiah Kimia
Department of
Chemistry, Faculty of
Mathematics and
Natural Sciences,
Universitas Jenderal
Soedirman,
Purwokerto,
Indonesia**



This work is licensed
under a [Creative
Commons Attribution
4.0 International
License](https://creativecommons.org/licenses/by/4.0/).

**SECONDARY METABOLITE FROM ENDOPHYTIC FUNGI *Aspergillus* sp.
THE LEAVE OF KUNYIT PUTIH (*Curcuma zedoaria* (BERG) ROSCOE)**

**METABOLIT SEKUNDER DARI JAMUR ENDOFITIK *Aspergillus* sp. DAUN
TUMBUHAN KUNYIT PUTIH (*Curcuma zedoaria* (BERG) ROSCOE)**

Muharni^{1*}, Fitriya², Widia Purwaningrum¹, and Ahmad Yogi Nugraha¹

¹Department of Chemistry, Faculty of Mathematics and Natural Sciences, Sriwijaya University, Palembang, Indonesia

²Department of Pharmacy, Faculty of Mathematics and Natural Sciences, Sriwijaya University, Palembang, Indonesia

*email: muharnimyd@yahoo.co.id

Received 9 February 2016; Accepted 2 May 2016; Available online 16 May 2016

ABSTRACT

The compound from endophytic fungi of *Aspergillus* sp. from leaves of kunyit putih (*Curcuma zedoaria* (Berg.) Roscoe) has been isolated. Isolation begins with cultivation of *Aspergillus* sp. in 18 L PDB's media (*Potato Dextrose Broth*) for 28 days. The liquid cultivation medium was extracted by partitioning method using ethylacetate and then evaporated. The extract was separated and purified by chromatography techniques. Elucidation structure of the isolated compound was analysis by spectroscopic method NMR 1D and 2D. Antibacterial activity of isolated compound was tested using the disc diffusion method at concentrations 2500, 1000, 500, and 125 ppm. The isolated compounds obtained in the form of a yellow oil (24.30 mg). The ¹³C NMR spectrum indicated 24 signals of carbon and base on analysis spectrum DEPT 135 showed 5 signal methynes carbon, 1 signals methylene, 9 signals of methyl and 9 signals quarternary carbon. These signals from ¹H and ¹³C-NMR suggested that this compound contained aromatic group and four carbonyl. The isolated compound show antibacterial activity at concentration 2500 ppm which inhibition zone for *E. coli*, *S. dysenteriae*, *S. aureus*, *B. subtilis* were 10.3 ; 8.3; 8.4; and 7.8 mm, respectively. Based on the analysis result of NMR 1D and 2D, the compound was methyl 6-(5'-(2"-acetoxo-2"-methylpropanoyl)-3"-methyl-2'-(3"-methylbutanoyl)phenyl)-3-methylbutanoate and has weak antibacterial activity.

Keyword: *Aspergillus* sp., *Curcuma zedoaria*, endophytic fungi

ABSTRAK

Telah diisolasi satu senyawa metabolit sekunder dari jamur endofitik *Aspergillus* sp. pada daun tumbuhan kunyit putih (*Curcuma zedoaria* (Berg) Roscoe). Isolasi diawali dengan kultivasi jamur *Aspergillus* sp. dalam 18 L media PDB (*Potato Dextrose Broth*) selama 48 hari. Medium kultur cair diekstraksi secara partisi menggunakan etil asetat dan dievaporasi. Ekstrak pekat etil asetat dipisahkan dan dimurnikan menggunakan teknik kromatografi. Penentuan struktur senyawa hasil isolasi dilakukan dengan cara spektroskopi NMR 1D dan 2D. Aktivitas antibakteri dari senyawa hasil isolasi telah diuji menggunakan metode difusi cakram dengan konsentrasi 2500, 1000, 500, 125 ppm. Dari penelitian ini diperoleh satu senyawa murni berupa minyak berwarna kuning sebanyak 24,30 mg. Spektrum ¹³C NMR menunjukkan adanya 24 sinyal karbon dan berdasarkan spektrum DEPT terdiri dari 5 karbon metin, 1 karbon metilen, 9 karbon metil dan 9 karbon

kwarterner. Senyawa hasil isolasi menunjukkan aktivitas antibakteri pada konsentrasi 2500 ppm dengan diameter zona bening berturut-turut untuk *E.coli*, *S. dysenteriae*, *S.aureus*, *B.subtilis* 10,3 ; 8,3; 8,4; dan 7,8 mm. Berdasarkan analisis data spektroskopi NMR 1D dan 2D maka diusulkan senyawa hasil isolasi adalah metil 6-(5'-(2"-asetoksi-2"-metilpropanoil)-3"-metil-2'-(3"-meilbutanoil)fenil)-3- metilbutanoat dan memiliki sifat aktivitas antibakteri yang lemah.

Kata kunci: *Aspergillus* sp, *Curcuma zedoaria*, jamur endofitik

INTRODUCTION

Endophytic fungi are defined as fungi which spend the whole or part of their lifecycle colonizing inter and intracellularly inside the healthy tissue of the host plants, typically causing no apparent symptoms of disease (Zhang, Song, & Tan, 2006; Rodriguez., White, Arnol, & Reman, 2009). Plant endophytic fungi have been recognized as an important and novel resource of natural bioactive products with potential application in agriculture, medicine and food industry (Gunatilaka, 2006; Verma, Kharmar, & Strobel, 2009). Novel antibiotics, antimycotics, immunosuppressants, and anticancer compounds are rarely founded after the isolation and culturing of individual endophytes which followed by purification and characterization of some of their natural products (Strobel, Daisy, Castillo, & Harper, 2004).

Endophytic fungi have been found in each plant spesies examined, and estimated that there are over one million fungal endophytes existed in the nature. Traditionally *Curcuma zedoaria* known as herbal medicine which possessing many biological activities. Some bioactive compounds that contained in *Curcuma zedoaria* have been reported are furanodiene, furanodienone, zedorone, curzerenone, curzeone, germacrone, 13-hydroxy germacrone, dihydrocurdione, curcumenone and zedoaronedirole which is of sesquiterpenoid. (Makabe, Maru, Kuwabara, Kamo, & Hirota, 2006). *Curcuma zedoaria* also contains epikurzerenone and kurzerene compounds (Mau, *et.al.* 2003) and isocurcumenol

which have antitumour effects (Lakshmi, *et. al.* 2011) and triterpenoid compounds.

Many endophytic bacterial live in association with their host and may play an important biological roles. Sulistiyani, Lisdiyanti, and Lestari, (2014) have been investigate the endophytic bacterial diversity associated with *Curcuma zedoaria* and total of 207 bacterial colonies were isolated from rhizomes, stems, and leaves and 73 endophytic bacteria were selected based on morphological characteristics.

Research about secondary metabolite that contained in endophytic fungi of *Curcuma zedoaria* also has been reported but still limited. Muharni, Fitrya, Ruliza, Susanti and Elfita (2014b) have been reported two compounds bis-(2-ethylhexyl)phtalat and 3-(2,5-diacetyl)-3,4-dihydroxyphenyl)-butan-2-nylpropionate from endophytic fungi *Penicillium* sp. Compound 2 (Pyranon derivated) show antibacterial activity to *S. Aureus* and antioxidant activity by DPPH method with IC₅₀ 16.05 µg/mL (Muharni, Fitrya, Milanti & Elfita, 2014a). In vivo assay as antibacterial from endophytic fungus *Penicillium* sp extract has been done used mice (*Mus musculus*) (Muharni, Heni, Fitrya & Roni, 2015). To complete profile of the chemical constituents of endophytic microbe of *Curcuma zedoaria*, in this paper we will be explained isolation of secondary metabolite compound from endophytic fungi *Aspergillus* sp in leaves of *Curcuma zedoaria*. The antibacterial activity of isolated compound have been done.

EXPERIMENTAL SECTION

Materials and Equipment

The leaves of kunyit putih were collected on Mei 2013 from the Indralaya, Ogan Ilir, South Sumatra. Material for isolation and cultivation endophytic fungi: ethanol 70%, NaOCl, chloramphenicol, potato dextrose broth (PDB), potato dextrose agar (PDA). Material for isolation compound: silica gel 60 (70-230 mesh.), thin layer chromatography (TLC) using Merck (Art.5554), silica gel 60 F₂₅₄, *n*-hexane, ethylacetate, and methanol. The organic solvents were used from distilled technical grade. Material for antibacterial activity assay: nutrient agar (NA) and nutrient broth (NB), and ampicillin.

The apparatus in the research were counter colony, autoclave, incubator, water bath, microscope, magnetic hotplate, UV lamp, column chromatography and generally apparatus in organic and microbiology laboratory, melting point was determined using Fisher John Apparatus. NMR spectra were recorded at 500 MHz (¹H) and 125 MHz (¹³C) on JEOL JNM ECA-500 spectrometer.

Procedure

Isolation of endophytic fungus

The study begins with the isolation of endophytic fungi from leaves *Curcuma zedoaria* plant. The procedure refers to Muharni *et al.* (2014) with slight modifications. The leaves sample were washed and sterilized in 70% ethanol for 5 min and 0.5% NaOCl for 5 min and then washed with sterile distilled water. The segment were placed on petri-plates containing potato dextrose agar medium (PDA). The plates were incubated at 25±2 °C. The plant segment observed every day to see the growth of endophytic fungus. Fungal colony that shows a different characteristic further purified by transferring it in the PDA medium other then some subculture to obtain pure fungal cultures (*Aspergillus* sp.) (Barik, Tayung,

Jagadev, & Duta, 2010; Kour *et al.*, 2008; Eyberger, Dondapati, & Porter, 2006)

Identification of the endophyte

The endophytic fungal strain was identified by the morphological method. The morphological examination was performed by scrutinizing the fungal culture, the mechanism of spore production, and the characteristics of the spores. All experiments and observations were repeated at twice (Guo *et al.*, 2008).

Cultivation of pure fungal strain

The procedure for the cultivation refers to Muharni *et al.*, 2014a with slight modifications. Cultivation of fungus that have been pure (*Aspergillus* sp), (a small park) were transferred into the medium under sterile conditions to the PDB medium. To isolate the secondary metabolites, the fungal strains were static cultivated in 30 flasks each containing 600 mL of PDB medium and incubated for 28 days at room temperature, so the metabolites sekundernya will enter into PDB medium. (Xu *et al.*, 2008). Furthermore filtered to separate filtrate and biomass. The Filtrate containing secondary metabolites, then partitioned with ethylacetate solvent. Then the ethyl acetate phase was further concentrated by rotary vacuum evaporator at 40 °C and obtained ethyl acetate fraction of liquid cultures (5,0 g).

Isolation sekundare metabolite from ethylacetate fraction of endophytic fungus

Concentrated ethylacetate from *Aspergillus* sp (5.0 g) were isolated using column chromatography with silica gel 60 as stationary phase. The solvent system used for chromatography was *n*-hexane with increasing portion of EtOAc (gradient elution system). The ratio of the solvent between *n*-hexane and EtOAc were 100 : 0, 90 : 10, 10 : 90). Fraction were collected every 10 mL and each fraction was tested by TLC. The spot were detected by UV light (254 and 366 nm). Fraction having spots with the same Rf

value were combined and treated as a group. Fraction 2nd (0.9 g) was rechromatography using the same method to give pure compound form yellow oil (24.3 mg) (Hundley, 2005). The molecular structure of pure compound were determined on the basis of spectroscopic analysis (¹H-NMR, ¹³C-NMR, DEPT, HMQC, HMBC, and COSY).

Antibacterial activity

The antibacterial activity assay used: *Escherichia coli*, *Sigella dysentryae*, *Staphylococcus aureus* and *Bacillus subtilis*. The antibacterial activity were determined by disc diffusion method were described previously for the preliminary of antibacterial activity (Lai, Chyau & Mau, 2004). A sterile paper disc was impregnated with test material and the disc was placed on the nutrient agar medium. Plates were then incubated at 37 °C for 72 h under anaerobic conditions. All disc diffusion tests were performed in three separated experiments and the antibacterial activity was expressed as the mean of inhibition diameters (mm). The test material was

prepared in various concentration and as standard used ampicillin 10 ppm.

RESULT AND DISCUSSION

Isolation and Identification The Secondary Metabolits

The fungus strain was identified as *Aspergillus* sp. Base on literature study *Aspergillus* sp species isolated as endophytes were usually obtained from several plant spesies such as, endophytic fungi *Aspergillus niger* from stem bark of *Garcinia griffithii* (Elfita, Muharni, Munawar, & Aryani, 2012), *Aspergillus niger var taxi* from *Taxus cuspidata* (Zhou *et al.*, 2009), *Aspergillus fumigates* from fruit of *G. griffithii* (Elfita, Muharni, & Indah, 2011), and *Podocarpus sp* (Sun, Rang & Wang, 2008), *Aspergillus flavus* from sambiloto (*Andographis peniculata* Nees) (Elfita, Muharni, Munawar, Salmi & Oktasari, 2010).

Fungus *Aspergillus sp* after cultivation in media 18L PDB then extracted with ethyl acetate and obtained 5 g of concentrated ethyl acetate extract. 5.0 g of ethylacetate extract after being separated by column chromatography techniques to yield pure compound in the form of yellow oil (24.3 mg) (**Figure 1**).




Figure 1. Isolation of the compound from ethyl acetate extract of *Aspergillus sp* from the leaves of *Curcuma zedoaria*

Table 1. The NMR data of isolated compound, recorded at ^1H - 500 MHz; ^{13}C -125 MHz, spectral data of recorded in CDCl_3

NO. C	δ_{C} (ppm)	DEPT	δ_{H} ppm (ΣH , multiplicity, J (Hz))	HMBC	COSY
1	174.6	C			
2	39.8	CH	2.46 (1H, <i>m</i>)		
3	27.4	CH	1.70 (2H, <i>m</i>)		0.93; 1.18
4	11.5	CH_3	0.93 (3H, <i>d</i> , $J = 7.75$)	27.4; 40.4	1.70
5	17.9	CH_3	1.18 (3H, <i>d</i> , $J = 7.75$),	27.4; 40.4	1.70
6	57.1	CH_3	3.80 (3H, <i>s</i>)	174.6	
1'	167.5	C			
2'	165.2	C			
3'	98.7	C			
4'	99.1	CH	5.93 (1H, <i>s</i>)	167.1 ; 98.5	
5'	167.1	C			
6'	96.5	CH	5.56 (1H, <i>s</i>)	174.6 ; 196.7	
7'	8.3	CH_3	1.94 (3H, <i>s</i>)	165.2; 167.1; 98.7	
1''	196.7	C			
2''	77.3	C			
3''	175.7	C			
4''	25.2	CH_3	1.65 (3H, <i>s</i>)		
5''	25.3	CH_3	1.61 (3H, <i>s</i>)	77.3 ; 196.7	
6''	26.7	CH_3	1.63 (3H, <i>s</i>)	77.3	
1'''	200.9	C			
2'''	40.4	CH_2	2.46 (2H, <i>m</i>)		
3'''	26.7	CH	1.67 (1H, <i>m</i>)	77.3; 200.9	0.85; 1.17
4'''	16.5	CH_3	1.17 (3H, <i>d</i> , $J = 7.15$)	26.7; 40.4	1.67; 2.46
5'''	11.7	CH_3	0.85 (3H, <i>d</i> , $J = 7.15$)	26.7; 40.4	1.67

The ^1H NMR data (**Table 1**) disclosed the presence of two protons at δ_{H} 5.93 and 5.56 ppm (1H, *s*) were characteristic for vinyl proton. The proton signal at δ_{H} 3.80 ppm (3H, *s*) was assigned to a methoxy group. Furthermore, the presence proton signals at δ_{H} 2.46 each (3H, *m*) for one proton methylene and one proton methine, signals at 1.67 (1H, *m*), and 1.70 (1H, *m*) for two methine groups. The ^1H NMR data also indicated the presence of eight methylprotons at δ_{H} 1.94 ppm, δ_{H} 1.65 ppm, δ_{H} 1.63 ppm and 1.61 ppm, δ_{H} 1.18 (3H, *d*, $J = 7.75$), 0.93 (3H, *d*, $J = 7.75$), 1.17 ppm (3H, *d*, $J = 7.15$), and 0.85 ppm (3H, *d*, $J = 7.15$).

The ^{13}C NMR spectrum (**Figure 2**) indicated 24 signal carbon consist that 10 signals as C sp^2 and 14 signals as C sp^3 . Base on analysis spectrum DEPT 135 showed 5 signal methines carbon at δ_{C}

99.1; 98.5; 38.9; 27.4 ppm and δ_{C} 26.7 ppm, 1 signals methylene carbon at δ_{C} 40.4 ppm, 9 signals methyl carbon at δ_{C} 57.1; 26.7; 25.3; 25.2; 17.9; 16.5; 11.7; 11.5 and δ_{C} 8.3 ppm and 9 signal quaternary carbon at δ_{C} 200.9; 196.7; 175.6; 174.5; 167.3; 167.1; 165.1, 98.7 and δ_{C} 77.3 ppm.

Signals carbon at δ_{C} 196.7 ppm and 200.9 ppm indicated these compound have two carbonyl groups and 175.6 and 174.5 characteristic for ester carbonyl. Signal methyl carbon at δ_{C} 57.1 ppm characteristic for signal methoxy carbon. These signals from ^1H and ^{13}C -NMR suggested that this compound contained aromatic group and four carbonyl group. The presences of the functional groups above were suggested by the long range coupling HMBC and correlation of the chemical H and C shift for all protonated carbons was determined based on the

HMQC spectrum as summarized in **Table 1**.

NMR 2D analysis for HMQC spectrum showed the proton signal at δ_H 5.93 ppm and δ_H 5.56 ppm attached to carbon signal at δ_C 99.1 ppm and 98,5 respectively and proton at δ_H 3.80 attached to carbon at δ_C 57.1 ppm. HMBC spectrum showed proton at δ_H 5.93 was correlated to carbon at δ_C 98.5 and δ_C 167.1 ppm, while proton at δ_H 5.56 showed correlation to carbon at δ_C 174.6 and 196.7 ppm. This data indicated that two proton vinilic was not place at the carbon besides it. Further HMBC spectrum showed correlation proton of δ_H 3.80 ppm (3H, s) to carbon at δ_C 174.6 ppm indicated as proton methoxy from ester group.

HMQC spectrum showed that

proton at δ_H 1.61 attached to carbon at δ_C 25.3 and HMBC spectrum showed that proton at δ_H 1.61 was correlated to carbon δ_C 77.3 dan δ_C 196.7 ppm. While proton at δ_H 1.63 ppm at HMQC spectrum showed attached to carbon at δ_C 26.7 ppm and HMBC spectrum showed correlation to carbon at 77.3 ppm. This data indicated that two proton methyl were bounded at the same carbon that δ_C 77.3 ppm. The proton at δ_H 1.94 at HMQC spectrum showed attached to carbon at δ_C 8.3 ppm and HMBC spectrum showed correlated to carbon at δ_C 98.7. Proton at δ_H 2.46 attached to carbon at 39.8 and 40.4 ppm (**Figure 3**). This data supported that proton at δ_H 2.46 ppm were one signal methine proton and one as signal methylene proton.




Figure 2. ^{13}C NMR spectrum isolated compound




Figure 3. HMQC Spectrum (A, B) and HMBC Spectrum (C, D)




Figure 4. HMBC spectrum (A) and dan COSY spectrum (B)

Furthermore at HMBC spectrum showed correlation proton at δ_H 1.94 ppm to carbon δ_C 165.2; δ_C 167.1; and δ_C 98.7 ppm indicated the proton attached at aromatic ring. HMQC spectrum also showed that proton at δ_H 1.18 and 0.93 ppm attached to carbon at δ_C 17.9 ppm, and δ_C 11.5 ppm, while HMBC spectrum (**Figure 4**) showed both of this proton was correlation to carbon δ_C 27.4 dan 39.8. Proton at δ_H 1.17 and 0.85 ppm at HMQC

spectrum showed attached to carbon at δ_C 16.5 and 11.7 ppm at HMBC spectrum showed both of this proton correlation to carbon δ_C 26.7 ppm 40.4 ppm.

COSY spectrum (**Figure 4**) showed proton at δ_H 1.17 ppm and 0.85 ppm each was correlated to proton at δ_H 1.67 ppm and 2.46, while that proton at δ_H 0.93 ppm and 1.18 correlated to proton at δ_H 1.70 ppm. The presences of the functional groups above were suggested by the long

range coupling HMBC experiment.

These spectroscopic data, therefore suggested that compound was identified as methyl 6-(5'-(2''-acetoxy-2''-methylpropanoyl)-3''-methyl-2'-(3'''-methylbutanoyl)phenyl)-3-methylbutanoate. HMBC correlation and structure this compound showed at **Figure 5**. Exploration of secondary metabolites research needs to be done in order to get the profile of organic compounds produced by endophytic fungus of *Curcuma zedoaria*.

Based on the literature study, the biosynthetic pathways of secondary metabolites produced from endophytic fungus has not been found clearly. The substances isolated from endophytic have different biosynthetic pathways: isoprenoid, polyketide, amino acid derivatives, and belonged to diverse structural groups: terpenoids, steroids, xanthenes, chinones, phenols, isocumarines, benzopyranones, tetralones, cytochalasines, and enniatines (Barbara, Christine, Anne, & Kristen, 2002). Literature survey also showed these compounds have never found either of *Curcuma zedoaria* or host plants of other plants. This compound also not yet been found of other endophytic fungi, but the compounds proposed are similar to

compounds that are reported of fungal endophytic dothiorelon B and dothiorelon C, were isolated from microbial *Dothiorella sp* who live on the leaves of the species *Cynodon dactylon* (L) (Poaceae) (Radji, 2005). Other similar compounds ever discovered was 2-{4-methyl-2-[(2-methylpropanoyl)oxy]phenyl}oxiran-2-yl)methyl-3-methylbutanoic (Yannai, 2004) (**Figure 6**).

Antibacterial Activity

The antibacterial activity of this compound was evaluated according to the method previously described. The antibacterial properties of isolated compound was evaluated according to the method described previously (Lai, 2004). This compound showed inactive antibacterial for all bacterial test until concentration test 1000 ppm. This compound will show activity by concentration 2500 ppm with the mean of inhibition diameters (mm) for *E. coli*, *S. dysenteriae*, *S. aureus*, *B. subtilis* 10,3 ; 8.3, 8.4; and 8.8 mm respectively and standard antibacterial ampicillin at concentration 10 ppm showed inhibition diameters 7.5; 8.5; 7.0; and 9.5 mm respectively. Base on this data the compound show weak activity.




Figure 5. Structure, (A), HMBC Correlation (B) and COSY correlation (C) of isolated compound




Figure 6. Structure of dothiorelon B, dothiorelon C from microbial *Dothiorella sp* and 2-{4-methyl-2-[(2-methylpropanoyl)oxy]phenyl}oxiran-2-yl)methyl-3-methylbutanoic.

CONCLUSION

A new compound have been isolated from the endophytic fungi *Aspergillus* sp from the leaves of kunyit putih (*Curcuma zedoaria*). Based on spectroscopic analysis H-NMR and C-NMR (1D and 2D), was identified as methyl 6-(5'-(2"-acetoxo-2"-methylpropanoyl) -3"-methyl-2'-(3"-methylbutanoyl)phenyl)-3-methylbutanoic. Isolated compound showed weak antibacterial activity.

ACKNOWLEDGEMENT

The authors are grateful to the Directorate General of Higher Education which research grant Fundamental 2014 was supported this research.

REFERENCE

- Barbara, S., Christine, B., Siegfried, D., Anne, K. R., and Kristen, K. (2002). Endophytic fungi: a source of novel biologically active secondary metabolites, *Mycological Research*, 106 (9), 996–1004
- Barik, B.P., Tayung, K., Jagadev, N., and Dutta, S. K. (2010). Phylogenetic placement of an endophytic fungus *Fusarium oxysporum* isolated from *acorus calamus* rhizome with antimicrobial activity. *European Journal Biological Science*. 2(1), 8-16.
- Elfita, Muharni, Munawar, Salni, dan A. Oktasari. (2010). Senyawa antimalaria dari jamur endofitik tumbuhan sambiloto (*Andrographis paniculata* Nees. *Jurnal Natur Indonesia* 13(2) 123-129.
- Elfita, Muharni, and T. Indah. (2011). Secondary metabolite of *Aspergillus fumigatus*, an endophytic fungi of the medicinal plant *Garcinia griffithii*. *Makara Seri Sains*, 15(2),124-128.
- Elfita, Muharni, Munawar, and S. Aryani. (2012). Secondary metabolite from endophytic Fungi *Aspergillus niger* of the stem bark of Kandis Gajah (*Garcinia griffithii*). *Indonesian Journal Chemistry*, 112(2), 195-200
- Eyberger, A. L., Dondapati, R., and Porter, J.R. (2006). Endophytic fungal isolated from *Podophyllum peltatum* produce podophyllotoxin. *Journal of Natural Products*, 69(8), 1121 - 1124.
- Gunatilaka A. A. L. (2006). Natural products from plant-associated microorganisms, distribution, structural diversity, bioactivity, and implications of their occurrence. *Journal of Natural Products*. 69(3), 505-526.
- Guo, L., Wu, J. Z., Han, T., Cao, T., Rahman, K., and Qin L.P. (2008). Chemical composition, antifungal and antitumor properties of ether extracts of *Scapania verrucosa* Heeg. and its endophytic fungus *Chaetomium fusiforme* . *Molecules* , 13(9), 2114-2125
- Hundley, N.J. (2005). *Struktur elucidation of bioactive compounds isolated from endophytes of Alstonia scholaris and Acmena graveolens*. Thesis. Department of Chemistry and Biochemistry. Brigham Young University.
- Kour, A., Shawl A.S., Rehman, S., Sultan p., Qazi, P. H., and Verma. (2008). Isolation and identification of an endophytic strain of *Fikusarium oxysporum* producing podophyllotoxin from *Juniperus recurva*. *World Journal of Microbiology and Biotechnology*, 24(7), 1115-1121
- Lai, E.Y.C., Chyau, C.C., and Mau, J.L. (2004). Antimicrobial activity and cytotoxicity of the essential oil of *Curcuma zedoaria*. *American Journal of Chinese Medicine*. 32(2): 281–290.
- Lakshmi.S., Padmaja.G., and Remani. P. (2011). Antitumor effect of isocurcumenol isolated from *Curcuma zedoaria* rhizomes on human and murine cancer cells.

- International Journal Medicinal Chemistry*, 2011, 1-13.
- Makabe, H., Maru, N., Kuwabara, A., Kamo, T., and Hirota, M. (2006). Anti-inflammatory Sesquiterpenes from *Curcuma zedoaria*. *Journal Natural Product*, 20(7), 680-685.
- Mau, J. L., Eric, C Lai., Nai, P.W, Chien, C.C., and Charng, C. Y.. (2003). Composition and antioxidant activity of the essential oil from *Curcuma zedoaria*. *Food Chemistry*, 82(4), 583–591.
- Muharni, Fitrya, Milanti, O., and Elfita. (2014). Antibacterial and antioxidant activity of pyranon derivated compound from endophytic fungi *Penicillium sp* of kunyit putih (*Curcuma zedoaria*), *Traditional Medicine journal*, 19(3), 107-112.
- Muharni, Fitrya, M.O. Ruliza, D.A. Susanti, and Elfita. (2014b). Di-2-Ethylhexyl phtalate and pyranon derivated from endophytic fungi *Penicillium sp* kunyit putih (*Curcuma zedoaria*) *Indonesian Journal Chemistry*, 14(3), 290-296.
- Muharni, Heni, Y., Fitrya, and Roni. (2015). Evaluation acute toxicity and antibacterial activity of *Penicillium sp* endophitic fungus extract of kunyit putih (*Curcuma zedoaria*) in mice (*Mus musculus* L.). *Journal Chemical and Pharmaceutical Research*, 7(9s), 147-151
- Radji, M. (2005). Peranan bioteknologi dan mikroba endofit dalam pengembangan obat herbal. *Majalah Ilmu Kefarmasian*, 2(3), 113 – 126.
- Rodriguez R.J., White, J. T., Arnol, A. E., and Reman, R.S. (2009). Fungal endophytic diversity and functional roles. *New Phytologist*, 182(2), 314–330.
- Strobel, G and Daisy , B., Castillo, U., and Harper J. (2004). Natural products from endophytic microorganisms. *Journal of Natural products*, 67(2), 257 – 268
- Sun, D., Ran, X., and Wang, J. (2008). Isolation and identification of taxol-producing endophytic fungus from *Podocarpus*. *Acta Microbiologica Sinica*. 48, 589-595.
- Sulistiyani, T. R., Lisdiyanti, P., and Lestari, Y. (2014). Population and diversity of endophytic bacteria associated with medicinal plant *Curcuma zedoaria*. *Microbiology Indonesia*, 8(2), 65 - 72
- Verma V. C., Kharmar, R. N., Strobel G. A. (2009). Chemical and functional diversity of natural products from plant associated endophytic fungi. *Natural Product Communications*, 4(11), 1211-1532
- Xu L., Zhou L., Zhao J., Li J., Li X., and Wang J. (2008). Fungal endophytes from *Dioscorea zingiberensis* and their antibacterial activity. *Letters in Applied Microbiology*. 46(1), 68-72.
- Yannai, S. (2004). Dictionary of food compounds with CD-ROM: Additives, flavors, and ingredients. Boca Raton: Chapman & Hall/CRC.
- Zhang, H. W., Song, Y. C., and Tan, R. X. (2006). Biology and chemistry of endophytes. *Natural Product Reports*, 23(5), 753-771.
- Zhou, K., Ping, W., Li, Q., Hao, S., Zhao, L., and Zhou, D. (2009). *Aspergillus niger* var *taxi*, a new species variant of taxol-producing fungus isolated from *Taxus cuspidata* in China. *Journal of Applied Microbiology*, 107, 1202–1207.