

Journal of
Indonesian Natural History



December 2017 Vol.5 No.1 & 2

EDITORS

Dr. Wilson Novarino

Associate Professor for Biology
Department of Biology
Andalas University, Indonesia

Email: editorjinh@jinh.org

Dr. Carl Traeholt

Programme Director, Southeast Asia
Research and Conservation Division
Copenhagen Zoo, Denmark

Email: ctraeholt@gmail.com

EDITORIAL BOARD

Dr. Ardinis Arbain

University of Andalas, Indonesia

Indra Arinal

National Park Management, Department of Forestry Indonesia

Dr. Ahimsa Campos-Arceiz

Nottingham University Malaysia Campus, Malaysia

Dr. Mads Frost Bertelsen

Research and Conservation Division, Copenhagen Zoo, Denmark

Dr. Susan Cheyne

Oxford University, Wildlife Research Unit, United Kingdom

Bjorn Dahlen

Green Harvest Environmental Sdn. Bhd, Malaysia

Dr. Niel Furey

Centre for Biodiversity Conservation, Royal University of Phnom Penh, Cambodia

Dr. Benoit Goossens

Cardiff University, United Kingdom

Dr. Djoko Iskandar

Bandung Institute of Technology, Indonesia

Dr. Matthew Linkie

Fauna & Flora International, Singapore

Dr. Erik Meijaard

People and Nature Consulting International, Indonesia

Dr. John Payne

Borneo Rhino Alliance, Malaysia

Dr. Ramadhanil Pitopang

Tadulako University, Indonesia

Dr. Lilik Budi Prasetyo

Bogor Institute of Agriculture, Indonesia

Dr. Dewi Malia Prawiradilaga

Indonesia Institute of Science, Indonesia

Dr. Rizaldi

University of Andalas, Indonesia

Dr. Dewi Imelda Roesma

University of Andalas, Indonesia

Dr. Jeffrine Rovie Ryan

Wildlife Forensics Lab, Dept. of Wildlife and National Parks, Malaysia

Boyd Simpson

Research and Conservation Division, Copenhagen Zoo, Denmark

Robert B. Stuebing

Herpetology and Conservation Biology, Indonesia

Dr. Sunarto

WWF-Indonesia

Dr. Jatna Supriatna

University of Indonesia

Dr. Campbell O. Webb

The Arnold Arboretum, Harvard University, USA

Dr. Zainal Z. Zainuddin

Borneo Rhino Alliance, Malaysia

The *Journal of Indonesian Natural History* is published biannually by the Department of Biology at the Andalas University, Padang, Sumatra Barat, Indonesia, in collaboration with Copenhagen Zoo, Denmark. The Department of Biology at Andalas University is dedicated to educating Indonesian biologists in the study and conservation of Indonesia's biodiversity and natural history. Copenhagen Zoo, through its Research and Conservation Division, supports *in-situ* conservation in Southeast Asia by assisting local organizations and individuals who undertake research, capacity building and the implementation of conservation programmes and projects.

The *Journal of Indonesian Natural History* is published by the Department of Biology, Andalas University, Indonesia in collaboration with Copenhagen Zoo, Denmark. It is available for free from www.jinh.net

COVER PHOTO: Jakarta's Pramuka bird market illuminates both rich cultural roots and a monumental conservation challenge
© Carl Traeholt, Copenhagen Zoo.

A rapid bird survey in Bukit Puar, Kerinci Seblat National Park; with notes on different survey methods

Muhammad Iqbal¹, Indra Yustian², Arum Setiawan², Rahmat Pratama Abdullah³ and Muhammad Isa⁴

¹KPB-SOS. Jl. Tanjung api-api km 10 Komplek PDK Blok E 1, Palembang 30152, Indonesia.

²Dept. Biology of FMIPA, University of Sriwijaya, Jl. Raya Palembang-Prabumulih km 32, Indralaya, Indonesia

³Community of Conservation, Dept. Biology of FMIPA, University of Sriwijaya, Indralaya, Indonesia

⁴Kerinci Seblat National Park Section V, Jl. Garuda km 7 Kayu Ara, Lubuk Linggau, Sumatera Selatan, Indonesia

Corresponding author: Muhammad Iqbal, Email: kpbsos26@yahoo.com

ABSTRACT

A rapid bird survey was carried out in Bukit Puar, Kerinci Seblat National Park, South Sumatra Province. Three survey methods were applied to assess bird diversity in this area: transect line, mist-netting and riparian survey. A total of 53 species from 27 families was recorded during survey. Among three methods applied, transect line is the most effective method to detect more bird species (45 species), followed by mist-netting (14 species) and riparian survey (nine species). There are 10 species protected by Indonesia law and five birds are listed as Near Threatened (NT) by International Union for Conservation of Nature or IUCN. These findings show that Bukit Puar still good habitat for various of birds, and the forest remaining should be protected.

ABSTRAK

Sebuah survei cepat untuk mengetahui keragaman burung telah dilakukan di Bukit Puar, Taman Nasional Kerinci Seblat, Provinsi Sumatera Selatan. Tiga metode survey digunakan dalam survey ini, yaitu: jalur transek (transect line), mist-netting dan survei dengan menyusuri sungai (riparian survey). Tercatat 53 jenis jenis burung dari 27 famil dalam survei ini. Dari ketiga metode yang digunakan, metode jalur transek merupakan metode yang paling efektif untuk mendeteksi lebih banyak keragaman burung dengan 45 jenis, diikuti mist-netting dengan 14 jenis dan survei menyusuri sungai dengan 9 jenis. Tercatat juga 10 jenis merupakan jenis dilindungi oleh Pemerintah Indonesia dan lima jenis masuk dalam kategori terancam punah menurut International Union for Conservation of Nature atau IUCN. Hasil dari survei ini menunjukkan bahwa Bukit Puar masih merupakan habitat yang baik untuk keragaman burung, dan hutan yang masih tersisa disini sebaiknya tetap dilindungi.

Keywords: *Bird, Bukit Puar, Kerinci Seblat National Park, transect line, mist-netting, riparian survey*

INTRODUCTION

Kerinci Seblat National Park (KSNP) with a total area of 1.389.509ha (Kepmenhut No.420/Menhut-II/2004) is a largest National Park in Sumatra, spanning over four provinces: West Sumatra, Jambi, Bengkulu and South Sumatra (Anon, 2016). The area has great habitats diversity that spans over continuous forest from lowlands to montane, including some natural stands of tropical pine (MacKinnon and Philipps, 1993). KSNP has

been listed as an Important Bird Area (IBA), an Endemic Bird Area and World Heritage Site in Indonesia (Anon, 2016; Holmes and Rombang, 2001; Stattersfield, 1998). The park shelters several important populations of Sumatra's charismatic wildlife, including many Sumatran endemic birds (Cochrane and Cubitt, 2004; Myers, 2005).

The avian diversity in KSNP and adjacent areas comprises over 380 species, which include 17 of the 20 Sumatran endemics (Anon, 2002). Most ornithological exploration of KSNP have taken place in the central parts of the park or in areas adjacent to Gunung Kerinci (eg. Holden, 1997; Hurrell, 1989; Lambert and Howes, 1989; Simpson,

Submitted 21st October, 2016. Accepted 26th February, 2017




Figure 1. Location of survey at Bukit Puar, Kerinci Seblat National Park, South Sumatra province. Yellow circle with dotted line is transect line, blue star is mist-net stations, and white arrow show riparian survey tracks.

1995), whereas information about avifauna outside these areas remains little known. In South Sumatra, an area of 250.613ha of KNSP, the avifauna remains unstudied (Anon, 2016).

Bukit Puar is a hilly forest in a part of KSNP that is located in the province of South Sumatra, where the habitat remains relatively intact, yet with no information available on the fauna in Bukit Puar, including its bird diversity. This survey is the first bird diversity survey to be undertaken in the Bukit Puar area of KSNP.

METHODS

Study Area

The survey took place around Bukit Puar and its adjacent areas ($02^{\circ} 47'S$, $102^{\circ}23'E$) in KSNP (Fig. 1). The survey took place in approx 500ha of Bukit Puar. The area is under management area of subsection V or SPTN V (SPTN = Seksi Pengelolaan Taman Nasional Wilayah) of KSNP. Administratively, the site is located in Muara Kulam village, South Sumatra province. It consists of secondary dipterocarp forest with little encroachment, apart from low intensity selective illegal logging by local people. The topography is hilly with numerous riparian areas (Fig. 2).

Surveys

From 18-22nd April 2016, three survey were undertaken to rapidly assess the bird diversity in Bukit Puar. We applied standard methods for conducting survey, following protocols for rapid survey of terrestrial birds (eg. Bibby et al., 2000; Herzog et al., 2011). The survey methods used were line transect, mist-netting and survey by canoe along riparian zones. Identification of birds were done with the aid of field guides, mainly of MacKinnon and Phillipps (1993).



Figure 2. Bukit Puar consists of hilly secondary dipterocarp forest with little encroachment ©Muhammad Iqbal.

Five fixed-radius 100 m transect line surveys were conducted between sunrise to mid-morning. We recorded distance from observer (≤ 50 m or > 50 m, and time of observation of all birds that could be identified either based on vocalisation or direct observation.

Five mist-nets (15 x 3m, 3-4mm mesh) were deployed in the survey area. The nets were stretched between two poles and checked every two hours, or four times a day. Birds caught in the mist-nets were removed and placed in a cage for identification and morphometric measures.

Access to Bukit Puar from Muara Kulam village took place in canoe powered by a 15Hp outboard motor. This time was used to record birds and other wildlife along the river. A total of 6km over a duration of two hours was undertaken for riparian surveys around Bukit Puar and surrounding area.

RESULTS AND DISCUSSION

A total of 53 species from 27 families (taxonomy, scientific name and English name follow Sukmantoro

et al., 2007) was recorded during survey (Table 1). Among the three survey methods used, transect line survey resulted in most recordings (45 species), followed by mist-netting (14 species) and riparian zones survey (nine species). Only one observation of the Greater racket-tailed drongo was recorded across all three survey methods.

From 53 species recorded in Bukit Puar, 10 species are protected by Indonesia law (Noerdjito and Maryanto, 2001), and five species are listed as Near Threatened (NT) on the IUCN red-list (Birdlife International, 2016). Rufous-collared kingfisher is the only species recorded that is listed as both protected and NT.

Whereas the transect line resulted in more species of recorded birds (85% of total), mist-netting is an important tool to detect shy illusive species, such as small birds from the under-storey community (Rahman, 2002; Whitman et al., 1997). In this survey, mist-nets captured four species that were not recorded by the other survey methods: Green broadbill, Hooded pitta, Banded pitta and Chestnut-rumped babbler. The Green broadbill is a good sample of why mist-netting is important too.



Table 1. A list of species recorded during 18-22 April 2016, and methods use to detect the occurrence of birds in Bukit Puar, Kerinci Seblat National Park. P = Protected by Indonesian law, NT = Near threatened (IUCN red-list status), 1 = Transect line, 2 = Mist-netting, 3 = Riparian survey with canoe.

Family	Scientific Name	English name	Methods		
			1	2	3
Accipitridae	<i>Unidentified Accipitridae</i> (P)	Eagle	+		
	<i>Spilornis cheela</i> (P)	Crested Serpent Eagle			+
Columbidae	<i>Treron sp</i>	Green Pigeon	+		
	<i>Ducula aenea</i>	Green Imperial Pigeon	+		
Psittacidae	<i>Loriculus galgulus</i>	Blue-crowned Hanging Parrot	+		
Cuculidae	<i>Cuculus sp</i>	Cuckoo	+		
	<i>Cuculus micropterus</i>	Indian Cuckoo	+		
	<i>Centropus bengalensis</i>	Lesser Coucal	+		+
Strigidae	<i>Ketupa ketupu</i>	Buffy Fish Owl			+
Apodidae	<i>Collocalia sp</i>	Swiftlet	+		
Hemiprocnidae	<i>Hemiproctne comata</i>	Whiskered Treeswift	+		
Alcedinidae	<i>Actenoides concretus</i> (P, NT)	Rufous Collared Kingfisher		+	+
	<i>Halcyon smyrnensis</i> (P)	White-throated Kingfisher	+		+
	<i>Alcedo meninting</i> (P)	Blue-eared Kingfisher	+		+
Bucerotidae	<i>Buceros rhinoceros</i> (P)	Rhinoceros Hornbill	+		
	<i>Anthracoceus sp</i> (P)	Hornbill	+		
Meropidae	<i>Nyctornis amictus</i>	Red-bearded Bee-eater	+		+
Capitonidae	<i>Megalaima rafflesii</i> (NT)	Red-crowned Barbet	+	+	
	<i>Megalaima mystacophanos</i> (NT)	Red-throated Barbet	+	+	
	<i>Megalaima australis</i>	Blue-eared Barbet	+		
Picidae	<i>Picus miniaceus</i>	Banded Woodpecker	+		
	<i>Micropternus brachyurus</i>	Rufous Woodpecker	+		
Eurylaimidae	<i>Corydon sumatranus</i>	Dusky Broadbill	+		
	<i>Eurylaimus ochromolus</i>	Black-and-yellow Broadbill	+		
	<i>Calyptomena viridis</i>	Green Broadbill		+	
	<i>Cymbirhynchus macrorhynchus</i>	Black-and-red Broadbill	+		
Pittidae	<i>Pitta sordida</i> (P)	Hooded Pitta		+	
	<i>Pitta guajana</i> (P)	Banded Pitta		+	
Campephagidae	<i>Hemipus sp</i>	Flycatcher-shrike	+		
Pycnonotidae	<i>Pycnonotus atriceps</i>	Black-headed Bulbul	+		
	<i>Pycnonotus aurigaster</i>	Sooty-headed Bulbul	+		
	<i>Pycnonotus melanicterus</i>	Black-crested Bulbul	+	+	
	<i>Pycnonotus brunneus</i>	Red-eyed Bulbul	+	+	
	<i>Criniger phaeocephalus</i>	Yellow-bellied Bulbul	+	+	
Chloropseidae	<i>Chloropsis cyanopogon</i> (NT)	Lesser Green Leafbird	+		
	<i>Chloropsis sonnerati</i>	Greater Green Leafbird	+		
	<i>Chloropsis cochinchinensis</i>	Blue-winged Leafbird	+		
Irenidae	<i>Irena puella</i>	Asian Fairy Bluebird	+	+	
	<i>Macronus gularis</i>	Striped Tit Babbler	+		

Timaliidae	<i>Stachyris maculata</i> NT	Chestnut-rumped Babbler		+	
	<i>Trichastoma bicolor</i>	Ferruginous Babbler	+	+	
	<i>Trichastoma rostratum</i>	White-chested Babbler	+		
Turdidae	<i>Copsychus saularis</i>	Oriental Magpie Robin	+		
Sylviidae	<i>Orthotomus ruficeps</i>	Ashy Tailorbird	+		
	<i>Prinia familiaris</i>	Bar-winged Prinia	+		
Muscicapidae	<i>Muscicapa dauurica</i>	Asian Brown Flycatcher	+		
Monarchidae	<i>Ficedula sp</i>	Flycatcher			+
	<i>Hypothymis azurea</i>	Black-naped Monarch	+		
Nectariniidae	<i>Arachnothera longirostra</i> P	Little Spiderhunter	+	+	
Dicaeidae	<i>Dicaeum sp</i>	Flowerpecker	+		
Dicruridae	<i>Dicrurus aeneus</i>	Bronzed Drongo	+		
	<i>Dicrurus paradiseus</i>	Greater Racquet-tailed Drongo	+	+	+
Corvidae	<i>Corvus enca</i>	Slender-billed Crow	+		
TOTAL			45	14	9

While Green broadbill is a common bird, it is often overlooked due to its behaviour, where it is often found stationary in the middle and high canopy and almost invisible (MacKinnon and Phillipps, 1993). For rapid bird censuses, however, transect line surveys are preferred to get an indication of species richness and abundance, since this method is less costly, less invasive, and less time-consuming (Arizaga et al., 2011).

As tropical landscapes become increasingly deforested, fragmented and human-dominated, riparian zones are becoming disproportionately important in connecting and harbouring populations of tropical forest organisms (Sekercioglu 2009). In South Sumatra, a bird survey was undertaken in Merang peat dome areas and Sembilang National Park (Goenner and Hasudungan, 2001, Iqbal, 2004, Waltert, 2008). Although surveys in riparian zones often result in low number of bird species, it is an important habitat that must be included in general biodiversity assessments. In this survey, we recorded three species Crested serpent eagle, Buffy fish owl and *Ficedula* flycatcher, that were not found elsewhere.

Birds are ideal subjects to assess an area for rapid biodiversity surveys, since they are the best known group of organisms in term of their taxonomy, biology, ecology, biogeography and conservation status (Herzog et al. 2011). Our studies in Bukit Puar suggest that the area is an important habitat for a range of bird species.

ACKNOWLEDGEMENT

We thank GIZ Bioclimate for funding this study. We are very grateful to Dr. Carl Traeholt and Dr. Wilson Novarino for their constructive suggestions and advises. Finally, thanks to our Muara Kulam biodiversity team who have valuable contribution: Doni Setiawan, Catur Prasetyo, Deny Noberio Gusna, Ina Aprilia, Rio Firman and Winda Indriati.

REFERENCES

- Anon. (2002). Management framework for Kerinci Seblat National Park 2002–2006. Kerinci Seblat Integrated Conservation and Development Project, Jambi, Indonesia.
- Anon. (2016). Taman Nasional Kerinci Seblat (TNKS). Seksi Pengelolaan TN. Wilayah V Lubuklinggau, Sumatera Selatan, Indonesia.
- Arizaga, J., Dean, J.I., Vilchels, A., Alonso, D. and A. Mendiburu. (2011). Monitoring communities of small birds: A comparison between mist-netting and counting. *Bird Study* **58**: 291–301
- Bibby, C., Jones, M. and S. Marsden. (2000). Teknik-teknik ekspedisi lapangan survei burung. Birdlife International Indonesia Programme, Bogor, Indonesia.

- BirdLife International. (2016). IUCN Red List for birds. Downloaded from <http://www.birdlife.org> on 04/10/2016.
- Cochrane, J. and G. Cubitt. (2004). The national parks and other wild places of Indonesia. New Holland Publisher, London, UK.
- Goenner, C. and F. Hasudungan. (2001). Sembilang Monitoring Report No. 1 Juli/August 2001. Wetlands International-Asia Pacific Indonesia Programme/Berbak Sembilang Project, Palembang, Indonesia.
- Herzog, S.K., O'Shea, B.J. and T. Pequeno. (2011). Toward a standardized protocol for rapid surveys of terrestrial bird communities. In Larsen, T.H. Core Standardized Methods for Rapid Biological Field Assessment. Conservation International, Arlington, USA.
- Holden, J. (1997). Bird list of Kerinci Seblat National Park, Sumatra and adjoining areas. Fauna and Flora International, Jambi, Indonesia. unpublished
- Holmes, D. and W.M. Rombang. (2001). Daerah penting bagi burung: Sumatera. PKA/Birdlife International-Indonesia Programme, Bogor, Indonesia.
- Hurrell, P. (1989). On the rediscovery of Schneider's Pitta in Sumatra. *Kukila* **4(1-2)**: 53-56.
- Iqbal, M. (2004). Pemantauan Kawasan Sembilang No. 9, Februari 2004. Laporan Teknis No. 82 Proyek Konservasi Terpadu Lahan Basah Pesisir Berbak Sembilang-Wetland International Indonesia Programme, Palembang, Indonesia.
- Lambert, F. and J.R. Howes. (1989). A recent sighting of Salvadori's Pheasant. *Kukila* **4(1-2)**: 56-58.
- Mackinnon, J. and K. Phillips. (1993). A field guide to the birds of Borneo, Sumatra, Java and Bali. Oxford University Press, UK.
- Myers, S. (2005). Kerinci-Seblat National Park, Sumatra, Indonesia. *Birding ASIA* **4**: 52-57.
- Noerdjito, M. and Maryanto, I. (2001). Jenis-jenis hayati yang dilindungi perundang-undangan Indonesia. Museum Zoologicum Bogoriense/LIPI/The Nature Conservancy/USAID, Cibinong, Indonesia.
- Rahman, M.A. (2002). Using mist-nets on canopy walkways in Malaysia to study canopy avifauna. *The Raffles Bulletin of Zoology* **50(2)**: 499-506.
- Serkecioglu, C.H. (2009). Tropical ecology: Riparian corridors connect fragmented forest bird populations. *Current Biology* **19(5)**: 210-213.
- Simpson, B. (1995). Sumatran Cochoa, *Cochoa beccarii*, on Gunung Kerinci, Sumatra. *Oriental Bird Club Bulletin* **21**: 50-52.
- Stattersfield, A.J., Crosby, M.J., Long, A.J. and D.C. Wege. 1998. Endemic Bird Areas of the World. Priorities for biodiversity conservation. BirdLife Conservation Series 7. Cambridge, UK.
- Sukmantoro, W., Irham, W., Novarino, W., Hasudungan, F., Kemp, N. and M. Muchtar. (2007). Daftar Burung Indonesia No. 2. The Indonesian Ornithologist's Union/LIPI/OBC Smythies Fund/Gibbon Foundation, Bogor, Indonesia.
- Waltert, M. 2008. Biodiversity assesment in the Merang peat dome area. South Sumatra Forest Management Project, Palembang, Indonesia.
- Whitman, A.A., Hagan, J.M., and N.V.L. Brokaw. (2007). A comparison of two bird survey techniques used in a subtropical forest. *The Condor* **99**: 955-965.

GUIDELINES FOR AUTHORS

The Journal of Indonesian Natural History will publish original work by:

- Indonesian or foreign scientists on any aspect of Indonesian natural history, including fauna, flora, habitats, management policy and use of natural resources
- Indonesian or foreign scientists on any aspect of regional natural history, including fauna, flora and habitats

Preference is given to material that has the potential to:

- Improve conservation intervention and management in Indonesia
- Enhances understanding of conservation needs in Indonesia
- Enhances the understanding of Indonesia's natural history

The Journal language will be in English with abstracts in Bahasa Indonesia as well as English. Authors of full papers are encouraged to provide a Bahasa Indonesia/English translation of their abstract. Submissions in Bahasa Indonesia will be accepted and translated into English only if accepted for publication.

Papers and Short Communications

Full Research Papers (2,000-7,000 words) and Short Communications (200-2,000 words) are invited on topics relevant to the Journal's focus, including:

- The conservation status, ecology or behaviour of wild species
- Status or ecology of habitats
- Prehistoric and extinct species and new findings
- Checklists of species, nationally or for a specific area
- Discoveries of new species records or range extensions.
- Biodiversity value associated with land use change, plantation development and forestry
- Biodiversity in High Conservation Value areas
- Reviews of conservation policy and legislation
- Conservation management plans for species, habitats or areas
- The nature and results of conservation initiatives, including case studies
- Sustainable use of wild species
- Community use and management of natural resources
- Abstracts of unpublished student theses (Short Communications only)

The Journal will not accept descriptions of new species, subspecies or other taxa.

How to Submit a Manuscript

Manuscripts should be submitted by e-mail to the Editor at:

editorjinh@jinh.org

In the covering e-mail, the Lead Author must confirm that:

- The submitted manuscript has not been published elsewhere,
- All of the authors have read the submitted manuscript and agreed to its submission,
- All research was conducted with the necessary approval and permits from the appropriate authorities

Review and editing

All contributors in English are strongly advised to ensure that their spelling and grammar is checked by a native English speaker before the manuscript is submitted to the Journal. The Editorial Team reserves the right to reject manuscripts that are poorly written.

Submission in Bahasa Indonesia is also accepted and the Journal secretariat will undertake a preliminary review of all manuscripts for suitability of content and format. Manuscripts that are rejected at this stage will not be sent for peer review and translation.

Manuscripts that pass through the preliminary review will be sent for translation and peer review by a minimum of two reviewers. Authors are welcome to suggest appropriate reviewers.

Proofs will be sent to authors as a portable document format (PDF) file attached to an e-mail note. PDF-readers can be downloaded free of charge from <http://www.adobe.com> and <http://www.nitropdf.com> for viewing PDF files. Corrected proofs should be returned to the Editor within 3 days of receipt. Minor corrections can be communicated by e-mail.

The Editorial Team also welcomes contributions to the other sections of the journal:

News

Concise reports (<300 words) on news of general interest to the study and conservation of Indonesia's natural history. News reports may include,

- Announcements of new initiatives; for example, the launch of new projects, conferences or funding opportunities.
- Announcements of important new reports or other publications related to Indonesian natural history.
- Summaries of important news from an authoritative published source; for example, new Indonesian species described in other journals, a new research technique, or a recent development in conservation.
- Summaries and analysis of new policies, decrees and laws relevant to the conservation of Indonesian natural history.

Letters to the Editor

Informative contributions (<650 words) in response to material published in the Journal.

Preparation of manuscripts

Full papers follow the style and format of papers published in the journal *Conservation Biology*. Authors should consult examples in *Conservation Biology* for guidance on general style.

Contributions should be in Bahasa Indonesia and/or UK English, double-spaced and in 'doc', 'rtf' or 'wpd' format, preferably as one file attached to one covering e-mail.

The cover page should contain;

The title and full mailing address, e-mail address and address of the Lead Author and all additional authors.

Contributing Papers should contain the following sections and be arranged in the following order: Abstract, Introduction, Methods, Results, Discussion, Acknowledgments, Literature Cited. Tables,

Figures and Plates (including legends), if included, should follow the Literature Cited.

All pages should be numbered consecutively. Do not number section headings or subheadings.

Title: This should be a succinct description of the work, in no more than 20 words.

Abstract: Abstracts should only be submitted for Full Papers. This should describe, in 100-300 words, the aims, methods, major findings and conclusions. It should be informative and intelligible without reference to the text, and should not contain any references or undefined abbreviations. Authors are encouraged to submit an English translation of Indonesian text and an Indonesian translation of an English text.

Keywords: From five to eight pertinent words, in alphabetical order.

Literature cited in text: Enclose citations in text in parentheses e.g. "Asian tapirs are no elephants when it comes to seed dispersal (Campos-Arceiz et al., 2011)."

Use an ampersand (&) between author surnames when the citation is parenthetical: (Traeholt & Idris, 2011).

When a citation is not parenthetical, use "and": "Our results agree with the predictions of Wolf and Rhymer (2001)."

For citations with more than two authors, use et al.: (Campos-Arceiz et al., 2011). Do not italicize et al.

List parenthetical citations in alphabetical order and chronologically from oldest to most recent and separate entries with a semicolon: (Campos-Arceiz et al., 2011; Geissman, 2009, 2010).

Separate the years with commas when citing multiple papers by the same author: (Corlett, 2007, 2010; Geissman, 1984, 1995, 1999, 2000).

"**In press**" means the cited paper has been accepted unconditionally for publication. Provide the year of publication in the text (Bird, 2010) and in Literature Cited section provide the volume number, and substitute "in press" for page numbers (Bird, I.M. 2010. Nesting success in arid lands. *Conservation Biology* 24: in press.).

Papers in review must be cited as unpublished and should not appear in the Literature Cited section.

Use an initial for the first (given) name and spell out the last name (surname) for other sources of unpublished data or information: (R. Fowler, unpublished data; M.E. Soulé, personal communication).

Software: capitalize the first letter only if the name of the program is a word (e.g., Vortex, ArcGIS). If the name of the program is not a word, use all capital letters (e.g., SAS).

The following are examples of Literature Cited house style:
Campos-Arceiz, A. and R.T. Corlett (2011). Big animals in a shrinking world—studying the ecological role of Asian megafauna as agents of seed dispersal. *Innovation* 10: 50–53.

Campos-Arceiz, A., Larringa, A.R., Weerasinghe, U.R., Takatsuki, S.,

Pastorini, J., Leimgruber, P., Fernando, P. and L. Santamaria (2008). Behavior rather than diet mediates seasonal differences in seed dispersal by Asian elephants. *Ecology* 89: 2684–2691.

MacArthur, R.H. & Wilson, E.O. (1967). *The Theory of Island Biogeography*. Princeton University Press, Princeton, USA.

Sutherland, W.J. (ed.) (1998). *Conservation Science and Action*. Blackwell Science, Oxford, UK.

Beck, B.B., Rapaport, L.G. & Stanley Price, M.R. (1994). Reintroduction of captive-born animals. In *Creative Conservation: Interactive Management of Wild and Captive Animals* (eds P.J.S. Olney, G.M. Mace & A.T.C. Feistner), pp. 265-286. Chapman & Hall, London, UK.

Traeholt, C., Bonthoeun, R., Rawson, B., Samuth, M., Virak, C. and Sok Vuthin (2005). Status review of pileated gibbon, *Hylobates pileatus* and yellow-cheeked crested gibbon, *Nomascus gabriellae*, in Cambodia. *Fauna & Flora International*, Phnom Penh, Cambodia.

Sun H. (2000). Status of the tiger and its conservation in Cambodia. MSc thesis, University of Minnesota, Minneapolis, USA.

IUCN (2010). 2010 IUCN Red List of Threatened Species. [Http://www.redlist.org](http://www.redlist.org) [accessed 1 February 2011].

Biography: This should describe the main research interests of all authors (<150 words total), apart from what is obvious from the subject of the manuscript and the authors' affiliations.

Tables, figures and plates: These should be self-explanatory, each on a separate page and with an appropriate caption. Figures can be submitted in colour as well as in black and white. The Editorial Team may decide to convert coloured figures into black and white should it be necessary due to printing cost and without diluting the message. Plates (black and white only) will only be included in an article if they form part of evidence that is integral to the subject studied (e.g., a photograph of a rare species), if they are of good quality, and if they do not need to be printed in colour.

Appendices: Lengthy tables, and questionnaires are discouraged. In special circumstances these may be made available for viewing online.

Species names: The first time a species is mentioned, its scientific name should follow in parenthesis and in italics: e.g., Asian elephant (*Elephas maximus*). English names should be in lower case throughout except where they incorporate a proper name (e.g., Asian elephant, Cookson's wildebeest, long-billed vulture).

Abbreviations: Full expansion should be given at first mention in the text.

Units of measurement: Use metric units only for measurements of area, mass, height, etc.

CONTENTS

3 GUEST EDITORIAL

6 NEWS AND NOTES

CONTRIBUTING PAPERS

- 10** The attitude, norm and perception of communities towards Sumatran tiger conservation initiatives in Aceh
Yeremiah R. Tjamin, Fachruddin M. Mangunjaya, Imran SL Tobing and Gugah Praharawati
- 16** A rapid bird survey in Bukit Puar, Kerinci Seblat National Park; with notes on different survey method
Muhammad Iqbal, Indra Yustian, Arum Setiawan, Rahmat Pratama Abdullah and Muhammad Isa
- 22** First record of international trafficking of Critically Endangered long-beaked echidnas (*Zaglossus* sp.)
Chris R. Shepherd and Emerson Y. Sy

SHORT COMMUNICATIONS

- 27** Fauna diversity recorded by camera traps in Bukit Tukau, Bukit Belading and Bukit Reban Kambing, Malaysia
Mariani Ramli and Carl Traeholt
- 30** Observations of Glossy Ibis *Plegadis falcinellus* in trade in bird markets in Java, Indonesia
Chris R. Shepherd, John Morgan and Jamie Bouhuys

32 GUIDELINES FOR AUTHORS



9 772338 865002
ISSN: 2338 865x

A rapid bird survey in Bukit Puar, Kerinci Seblat National Park; with notes on different survey methods

By Arum Setiawan

A rapid bird survey in Bukit Puar, Kerinci Seblat National Park; with notes on different survey methods

Muhammad Iqbal¹, Indra Yustian², Arum Setiawan², Rahmat Pratama Abdullah³ and Muhammad Isa⁴

¹KPB-SOS. Jl. Tanjung api-api km 10 Komplek PDK Blok E 1, Palembang 30152, Indonesia.

²Dept. Biology of FMIPA, University of Sriwijaya, Jl. Raya Palembang-Prabumulih km 32, Indralaya, Indonesia

³Community of Conservation, Dept. Biology of FMIPA, University of Sriwijaya, Indralaya, Indonesia

⁴Kerinci Seblat National Park Section V, Jl. Garuda km 7 Kayu Ara, Lubuk Linggau, Sumatera Selatan, Indonesia

Corresponding author: Muhammad Iqbal, Email: kpbsos26@yahoo.com

ABSTRACT

A rapid bird survey was carried out in Bukit Puar, Kerinci Seblat National Park, South Sumatra Province. Three survey methods were applied to assess bird diversity in this area: transect line, mist-netting and riparian survey. A total of 53 species from a total of 53 species from 27 families was recorded during survey. Among three methods applied, transect line is the most effective method to detect more bird species (45 species), followed by mist-netting (14 species) and riparian survey (nine species). There are 10 species protected by Indonesia law and five birds are listed as Near Threatened (NT) by International Union for Conservation of Nature or IUCN. These findings show that Bukit Puar still good habitat for various of birds, and the forest remaining should be protected.

ABSTRAK

Sebuah survei cepat untuk mengetahui keragaman burung telah dilakukan di Bukit Puar, Taman Nasional Kerinci Seblat, Provinsi Sumatera Selatan. Tiga metode survey digunakan dalam survey ini, yaitu: jalur transek (transect line), mist-netting dan survei dengan menyusuri sungai (riparian survey). Tercatat 53 jenis jenis burung dari 27 famil dalam survei ini. Dari ketiga metode yang digunakan, metode jalur transek merupakan metode yang paling efektif untuk mendeteksi lebih banyak keragaman burung dengan 45 jenis, diikuti mist-netting dengan 14 jenis dan survei menyusuri sungai dengan 9 jenis. Tercatat juga 10 jenis merupakan jenis dilindungi oleh Pemerintah Indonesia dan lima jenis masuk dalam kategori terancam punah menurut International Union for Conservation of Nature atau IUCN. Hasil dari survei ini menunjukkan bahwa Bukit Puar masih merupakan habitat yang baik untuk keragaman burung, dan hutan yang masih tersisa disini sebaiknya tetap dilindungi.

Keywords: *Bird, Bukit Puar, Kerinci Seblat National Park, transect line, mist-netting, riparian survey*

INTRODUCTION

Kerinci Seblat National Park (KSNP) with a total area of 1.389.509ha (Kepmenhut No.420/Menhut-II/2004) is a largest National Park in Sumatra, spanning over four provinces: West Sumatra, Jambi, Bengkulu and South Sumatra (Anon, 2016). The area has great habitats diversity that spans over continuous forest from lowlands to montane, including some natural stands of tropical pine (MacKinnon and Philipps, 1993). KSNP has

been listed as an Important Bird Area (IBA), an Endemic Bird Area and World Heritage Site in Indonesia (Anon, 2016; Holmes and Rombang, 2001; Stattersfield, 1998). The park shelters several important populations of Sumatra's charismatic wildlife, including many Sumatran endemic birds (Cochrane and Cubitt, 2004; Myers, 2005).

The avian diversity in KSNP and adjacent areas comprises over 380 species, which include 17 of the 20 Sumatran endemics (Anon, 2002). Most ornithological exploration of KSNP have taken place in the central parts of the park or in areas adjacent to Gunung Kerinci (eg. Holden, 1997; Hurrel, 1989; Lambert and Howes, 1989; Simpson,

Submitted 21st October, 2016. Accepted 26th February, 2017




Figure 1. Location of survey at Bukit Puar, Kerinci Seblat National Park, South Sumatra province. Yellow circle with dotted line is transect line, blue star is mist-net stations, and white arrow show riparian survey tracks.

1995), whereas information about avifauna outside these areas remains little known. In South Sumatra, an area of 250.613ha of KNSP, the avifauna remains unstudied (Anon, 2016).

Bukit Puar is a hilly forest in a part of KSNP that is located in the province of South Sumatra, where the habitat remains relatively intact, yet with no information available on the fauna in Bukit Puar, including its bird diversity. This survey is the first bird diversity survey to be undertaken in the Bukit Puar area of KSNP.

Surveys

From 18-22nd April 2016, three survey were undertaken to rapidly assess the bird diversity in Bukit Puar. We applied standard methods for conducting survey, following protocols for rapid survey of terrestrial birds (eg. Bibby et al., 2000; Herzog et al., 2011). The survey methods used were line transect, mist-netting and survey by canoe along riparian zones. Identification of birds were done with the aid of field guides, mainly of MacKinnon and Phillipps (1993).

METHODS

Study Area

The survey took place around Bukit Puar and its adjacent areas ($02^{\circ} 47'S$, $102^{\circ}23'E$) in KSNP (Fig. 1). The survey took place in approx 500ha of Bukit Puar. The area is under management area of subsection V or SPTN V (SPTN = Seksi Pengelolaan Taman Nasional Wilayah) of KSNP. Administratively, the site is located in Muara Kulam village, South Sumatra province. It consists of secondary dipterocarp forest with little encroachment, apart from low intensity selective illegal logging by local people. The topography is hilly with numerous riparian areas (Fig. 2).



Figure 2. Bukit Puar consists of hilly secondary dipterocarp forest with little encroachment ©Muhammad Iqbal.

Five fixed-radius 100 m transect line surveys were conducted between sunrise to mid-morning. We recorded distance from observer (≤ 50 m or > 50 m, and time of observation of all birds that could be identified either based on vocalisation or direct observation.

Five mist-nets (15 x 3m, 3-4mm mesh) were deployed in the survey area. The nets were stretched between two poles and checked every two hours, or four times a day. Birds caught in the mist-nets were removed and placed in a cage for identification and morphometric measures.

Access to Bukit Puar from Muara Kulam village took place in canoe powered by a 15Hp outboard motor. This time was used to record birds and other wildlife along the river. A total of 6km over a duration of two hours was undertaken for riparian surveys around Bukit Puar and surrounding area.

RESULTS AND DISCUSSION

A total of 53 species from 27 families (taxonomy, scientific name and English name follow Sukmantoro

et al., 2007) was recorded during survey (Table 1). Among the three survey methods used, transect line survey resulted in most recordings (45 species), followed by mist-netting (14 species) and riparian zones survey (nine species). Only one observation of the Greater racket-tailed drongo was recorded across all three survey methods.

From 53 species recorded in Bukit Puar, 10 species are protected by Indonesia law (Noerdjito and Maryanto, 2001), and five species are listed as Near Threatened (NT) on the IUCN red-list (Birdlife International, 2016). Rufous-collared kingfisher is the only species recorded that is listed as both protected and NT.

Whereas the transect line resulted in more species of recorded birds (85% of total), mist-netting is an important tool to detect shy illusive species, such as small birds from the under-storey community (Rahman, 2002; Whitman et al., 1997). In this survey, mist-nets captured four species that were not recorded by the other survey methods: Green broadbill, Hooded pitta, Banded pitta and Chestnut-rumped babbler. The Green broadbill is a good sample of why mist-netting is important too.




Table 1. A list of species recorded during 18-22 April 2016, and methods use to detect the occurrence of birds in Bukit Puar, Kerinci Seblat National Park. P = Protected by Indonesian law, NT = Near threatened (IUCN red-list status), 1 = Transect line, 2 = Mist-netting, 3 = Riparian survey with canoe.

Family	Scientific Name	English name	Methods		
			1	2	3
Accipitridae	<i>Unidentified Accipitridae</i> (P)	Eagle	+		
	<i>Spilornis cheela</i> (P)	Crested Serpent Eagle			+
Columbidae	<i>Treron sp</i>	Green Pigeon	+		
	<i>Ducula aenea</i>	Green Imperial Pigeon	+		
Psittacidae	<i>Loriculus galgulus</i>	Blue-crowned Hanging Parrot	+		
Cuculidae	<i>Cuculus sp</i>	Cuckoo	+		
	<i>Cuculus micropterus</i>	Indian Cuckoo	+		
	<i>Centropus bengalensis</i>	Lesser Coucal	+		+
Strigidae	<i>Ketupa ketupu</i>	Buffy Fish Owl			+
Apodidae	<i>Collocalia sp</i>	Swiftlet	+		
Hemiprocnidae	<i>Hemiproctne comata</i>	Whiskered Treeswift	+		
Alcedinidae	<i>Actenoides concretus</i> (P, NT)	Rufous Collared Kingfisher		+	+
	<i>Halcyon smyrnensis</i> (P)	White-throated Kingfisher	+		+
	<i>Alcedo meninting</i> (P)	Blue-eared Kingfisher	+		+
Bucerotidae	<i>Buceros rhinoceros</i> (P)	Rhinoceros Hornbill	+		
	<i>Anthracoseros sp</i> (P)	Hornbill	+		
Meropidae	<i>Nyctornis amictus</i>	Red-bearded Bee-eater	+		+
Capitonidae	<i>Megalaima rafflesii</i> (NT)	Red-crowned Barbet	+	+	
	<i>Megalaima mystacophanos</i> (NT)	Red-throated Barbet	+	+	
	<i>Megalaima australis</i>	Blue-eared Barbet	+		
Picidae	<i>Picus miniaceus</i>	Banded Woodpecker	+		
	<i>Micropternus brachyurus</i>	Rufous Woodpecker	+		
Eurylaimidae	<i>Corydon sumatranus</i>	Dusky Broadbill	+		
	<i>Eurylaimus ochromolus</i>	Black-and-yellow Broadbill	+		
	<i>Calyptomena viridis</i>	Green Broadbill		+	
	<i>Cymbirhynchus macrorhynchos</i>	Black-and-red Broadbill	+		
Pittidae	<i>Pitta sordida</i> (P)	Hooded Pitta		+	
	<i>Pitta guajana</i> (P)	Banded Pitta		+	
Campephagidae	<i>Hemipus sp</i>	Flycatcher-shrike	+		
Pycnonotidae	<i>Pycnonotus atriceps</i>	Black-headed Bulbul	+		
	<i>Pycnonotus aurigaster</i>	Sooty-headed Bulbul	+		
	<i>Pycnonotus melanicterus</i>	Black-crested Bulbul	+	+	
	<i>Pycnonotus brunneus</i>	Red-eyed Bulbul	+	+	
	<i>Criniger phaeocephalus</i>	Yellow-bellied Bulbul	+	+	
Chloropseidae	<i>Chloropsis cyanopogon</i> (NT)	Lesser Green Leafbird	+		
	<i>Chloropsis sonnerati</i>	Greater Green Leafbird	+		
	<i>Chloropsis cochinchinensis</i>	Blue-winged Leafbird	+		
Irenidae	<i>Irena puella</i>	Asian Fairy Bluebird	+	+	
	<i>Macronus gularis</i>	Striped Tit Babbler	+		

Timaliidae	<i>Stachyris maculata NT</i>	Chestnut-rumped Babbler		+	
	<i>Trichastoma bicolor</i>	Ferruginous Babbler	+	+	
	<i>Trichastoma rostratum</i>	White-chested Babbler	+		
Turdidae	<i>Copsychus saularis</i>	Oriental Magpie Robin	+		
Sylviidae	<i>Orthotomus ruficeps</i>	Ashy Tailorbird	+		
	<i>Prinia familiaris</i>	Bar-winged Prinia	+		
Muscicapidae	<i>Muscicapa dauurica</i>	Asian Brown Flycatcher	+		
Monarchidae	<i>Ficedula sp</i>	Flycatcher			+
	<i>Hypothymis azurea</i>	Black-naped Monarch	+		
Nectariniidae	<i>Arachnothera longirostra P</i>	Little Spiderhunter	+	+	
Dicaeidae	<i>Diaeum sp</i>	Flowerpecker	+		
Dicruridae	<i>Dicrurus aeneus</i>	Bronzed Drongo	+		
	<i>Dicrurus paradiseus</i>	Greater Racquet-tailed Drongo	+	+	+
Corvidae	<i>Corvus enca</i>	Slender-billed Crow	+		
TOTAL			45	14	9

While Green broadbill is a common bird, it is often overlooked due to its behaviour, where it is often found stationary in the middle and high canopy and almost invisible (MacKinnon and Phillipps, 1993). For rapid bird censuses, however, transects line surveys are preferred to get an indication of species richness and abundance, since this method is less costly, less invasive, and less time-consuming (Arizaga et al., 2011).

As tropical landscapes become increasingly deforested, fragmented and human-dominated, riparian zones are becoming disproportionately important in connecting and harbouring populations of tropical forest organisms (Sekercioglu 2009). In South Sumatra, a bird survey was undertaken in Merang peat dome areas and Sembilang National Park (Goenner and Hasudungan, 2001, Iqbal, 2004, Waltert, 2008). Although surveys in riparian zones often result in low number of bird species, it is an important habitat that must be included in general biodiversity assessments. In this survey, we recorded three species Crested serpent eagle, Buffy fish owl and *Ficedula* flycatcher, that were not found elsewhere.

Birds are ideal subjects to assess an area for rapid biodiversity surveys, since they are the best known group of organisms in term of their taxonomy, biology, ecology, biogeography and conservation status (Herzog et al. 2011). Our studies in Bukit Puar suggest that the area is an important habitat for a range of bird species.

ACKNOWLEDGEMENT

We thank GIZ Bioclimate for funding this study. We are very grateful to Dr. Carl Traeholt and Dr. Wilson Novarino for their constructive suggestions and advises. Finally, thanks to our Muara Kulam biodiversity team who have valuable contribution: Doni Setiawan, Catur Prasetyo, Deny Noberio Gusna, Ina Aprilia, Rio Firman and Winda Indriati.

REFERENCES

- Anon. (2002). Management framework for Kerinci Seblat National Park 2002–2006. Kerinci Seblat Integrated Conservation and Development Project, Jambi, Indonesia.
- Anon. (2016). Taman Nasional Kerinci Seblat (TNKS). Seksi Pengelolaan TN. Wilayah V Lubuklinggau, Sumatera Selatan, Indonesia.
- Arizaga, J., Dean, J.I., Vilchels, A., Alonso, D. and A. Mendiburu. (2011). Monitoring communities of small birds: A comparison between mist-netting and counting. *Bird Study* 58: 291–301
- Bibby, C., Jones, M. and S. Marsden. (2000). Teknik-teknik ekspedisi lapangan survei burung. Birdlife International Indonesia Programme, Bogor, Indonesia.

- BirdLife International. (2016). IUCN Red List for birds. Downloaded from <http://www.birdlife.org> on 04/10/2016.
- Cochrane, J. and G. Cubitt. (2004). The national parks and other wild places of Indonesia. New Holland Publisher, London, UK.
- Goenner, C. and F. Hasudungan. (2001). Sembilang Monitoring Report No. 1 Juli/August 2001. Wetlands International-Asia Pacific Indonesia Programme/ Berbak Sembilang Project, Palembang, Indonesia.
- Herzog, S.K., O'Shea, B.J. and T. Pequeno. (2011). Toward a standardized protocol for rapid surveys of terrestrial bird communities. In Larsen, T.H. Core Standardized Methods for Rapid Biological Field Assessment. Conservation International, Arlington, USA.
- Holden, J. (1997). Bird list of Kerinci Seblat National Park, Sumatra and adjoining areas. Fauna and Flora International, Jambi, Indonesia. unpublished
- Holmes, D. and W.M. Rombang. (2001). Daerah penting bagi burung: Sumatera. PKA/Birdlife International-Indonesia Programme, Bogor, Indonesia.
- Hurrell, P. (1989). On the rediscovery of Schneider's Pitta in Sumatra. *Kukila* 4(1-2): 53-56.
- Iqbal, M. (2004). Pemantauan Kawasan Sembilang No. 9, Februari 2004. Laporan Teknis No. 82 Proyek Konservasi Terpadu Lahan Basah Pesisir Berbak Sembilang-Wetland International Indonesia Programme, Palembang, Indonesia.
- Lambert, F. and J.R. Howes. (1989). A recent sighting of Salvadori's Pheasant. *Kukila* 4(1-2): 56-58.
- Mackinnon, J. and K. Phillips. (1993). A field guide to the birds of Borneo, Sumatra, Java and Bali. Oxford University Press, UK.
- Myers, S. (2005). Kerinci-Seblat National Park, Sumatra, Indonesia. *Birding ASIA* 4: 52-57.
- Noerdjito, M. and Maryanto, I. (2001). Jenis-jenis hayati yang dilindungi perundang-undangan Indonesia. Museum Zoologicum Bogoriense/LIPI/The Nature Conservancy/USAID, Cibinong, Indonesia.
- Rahman, M.A. (2002). Using mist-nets on canopy walkways in Malaysia to study canopy avifauna. *The Raffles Bulletin of Zoology* 50(2): 499-506.
- Serkecioglu, C.H. (2009). Tropical ecology: Riparian corridors connect fragmented forest bird populations. *Current Biology* 19(5): 210-213.
- Simpson, B. (1995). Sumatran Cochoa, *Cochoa beccarii*, on Gunung Kerinci, Sumatra. *Oriental Bird Club Bulletin* 21: 50-52.
- Stattersfield, A.J., Crosby, M.J., Long, A.J. and D.C. Wege. 1998. Endemic Bird Areas of the World. Priorities for biodiversity conservation. BirdLife Conservation Series 7. Cambridge, UK.
- Sukmantoro, W., Irham, W., Novarino, W., Hasudungan, F., Kemp, N. and M. Muchtar. (2007). Daftar Burung Indonesia No. 2. The Indonesian Ornithologist's Union/LIPI/OBC Smythies Fund/Gibbon Foundation, Bogor, Indonesia.
- Waltert, M. 2008. Biodiversity assesment in the Merang peat dome area. South Sumatra Forest Management Project, Palembang, Indonesia.
- Whitman, A.A., Hagan, J.M., and N.V.L. Brokaw. (2007). A comparison of two bird survey techniques used in a subtropical forest. *The Condor* 99: 955-965.

A rapid bird survey in Bukit Puar, Kerinci Seblat National Park; with notes on different survey methods

ORIGINALITY REPORT

15%

SIMILARITY INDEX

MATCH ALL SOURCES (ONLY SELECTED SOURCE PRINTED)

★biovalentia.mipa.unsri.ac.id

Internet

10%

EXCLUDE QUOTES ON

EXCLUDE BIBLIOGRAPHY ON

EXCLUDE MATCHES < 1%

**FORMAT PENILAIAN (VALIDASI & PEER REVIEW)
LEMBAR
HASIL PENILAIAN SEJAWAT SEBIDANG ATAU *PEER REVIEW*
KARYA ILMIAH : JURNAL ILMIAH**

Jurnal Artikel Ilmiah : A rapid bird survey in Bukit Puar, Kerinci Seblat National Park; with notes on different survey method

Penulis Artikel Ilmiah : Arum Setiawan

Identitas Jurnal Artikel Ilmiah : a. Nama Jurnal : Journal of Indonesian Natural History
b. Nomor/Volume/Hal : 2/5/16-21
c. Edisi (bulan/tahun) : Desember/2017
d. Penerbit : Jurusan Biologi FMIPA Universitas Andalas
e. Jumlah Halaman : 6

Kategori Publikasi Jurnal Ilmiah : Jurnal Ilmiah Internasional Bereputasi
(beri \checkmark pada kategori yang tepat) Jurnal Ilmiah Internasional
 Jurnal Ilmiah Nasional Terakreditasi S1, S2
 Jurnal Ilmiah Nasional Terakreditasi S3, S4.
 Jurnal Ilmiah Nasional tidak Terakreditasi

I. Hasil Penilaian Validasi :

No.	ASPEK	URAIAN/KOMENTAR PENILAIAN
1.	Indikasi Plagiasi	15 %
2.	Linearitas	Sudah linier dengan bidang keilmuan biologi konservasi

II. Hasil Penilaian *Peer Review* :

Komponen Yang Dinilai	Nilai Maksimal Jurnal Ilmiah (isikan di kolom yang sesuai)					Nilai Akhir Yang Diperoleh
	Internasional Bereputasi (Maks 40)	Internasional (Maks 20)	Nasional Terakreditasi S1, S2 Maks 25	Nasional Terakreditasi S3, S4 Maks 20	Nasional tidak Terakreditasi (maks 10)	
Kelengkapan dan Kesesuaian unsur isi jurnal (10%)					1	1
Ruang lingkup dan kedalaman pembahasan (30%)					3	3
Kecukupan dan Kemutakhiran data/informasi dan metodologi (30%)					3	3
Kelengkapan unsur dan kualitas penerbit (30%)					3	2
Total = (100%)					10	9
Kontribusi Pengusul (Penulis Pertama /Anggota Utama)	Anggota Utama=(0,4x9)/4=0,9					0,9

KOMENTAR/ULASAN *PEER REVIEW*

• Kelengkapan dan Kesesuaian Unsur:	Paper terkait survey burung di Bukit Puar, Taman Nasional Kerinci Seblat. Isi paper sudah memenuhi kaidah-kaidah karya ilmiah, dan sudah sesuai dengan bidang biologi konservasi.
• Ruang Lingkup dan Kedalaman Pembahasan:	Hasil penelitian dibahas cukup komprehensif dengan penyampaian perbandingan dari temuan-temuan penelitian lainnya dan teori terkait. Referensi yang diacu dalam pembahasan sudah cukup update untuk bidang kajian ini.
• Kecukupan & Kemutakhiran Data & Metodologi:	Data-data hasil penelitian cukup baik dan didukung penjelasan, peta lokasi dan gambar yang ditampilkan cukup menarik. Data didapatkan dengan menggunakan metode yang standard.
• Kelengkapan Unsur & Kualitas Penerbit:	Penerbit Jurusan Biologi FMIPA Universitas Andalas berkualitas baik, dan jurnal belum terkreditasi.

Surabaya, 15 Mei 2020
Penilai 1



Prof. Hery Purnobasuki, M.Si., Ph.D.
NIP 196705071991021001
Unit Kerja : Jurusan Biologi FST Unair
Bidang Ilmu : Biologi
Jabatan/Pangkat : Guru Besar/Pembina Utama Madya

FORMAT PENILAIAN (VALIDASI & PEER REVIEW)

LEMBAR

1.56.

HASIL PENILAIAN SEJAWAT SEBIDANG ATAU PEER REVIEW

KARYA ILMIAH : JURNAL ILMIAH

Jurnal Artikel Ilmiah : A rapid bird survey in Bukit Puar, Kerinci Seblat National Park; with notes on different survey method

Penulis Artikel Ilmiah : Arum Setiawan

Identitas Jurnal Artikel Ilmiah : a. Nama Jurnal : Journal of Indonesian Natural History
 b. Nomor/Volume/Hal : 2/5/16-21
 c. Edisi (bulan/tahun) : Desember/2017
 d. Penerbit : Jurusan Biologi FMIPA Universitas Andalas
 e. Jumlah Halaman : 6

Kategori Publikasi Jurnal Ilmiah : Jurnal Ilmiah Internasional Bereputasi
 (beri \surd pada kategori yang tepat) Jurnal Ilmiah Internasional
 Jurnal Ilmiah Nasional Terakreditasi S1, S2
 Jurnal Ilmiah Nasional Terakreditasi S3, S4.
 Jurnal Ilmiah Nasional tidak Terakreditasi

I. Hasil Penilaian Validasi :

No.	ASPEK	URAIAN/KOMENTAR PENILAIAN
1.	Indikasi Plagiasi	15 %
2.	Linearitas	

II. Hasil Penilaian Peer Review :

Komponen Yang Dinilai	Nilai Maksimal Jurnal Ilmiah (isikan di kolom yang sesuai)					Nilai Akhir Yang Diperoleh
	Internasional Bereputasi (Maks 40)	Internasional (Maks 20)	Nasional Terakreditasi S1, S2 Maks 25	Nasional Terakreditasi S3, S4 Maks 20	Nasional tidak Terakreditasi (maks 10)	
Kelengkapan dan Kesesuaian unsur isi jurnal (10%)					1	1
Ruang lingkup dan kedalaman pembahasan (30%)					3	2,5
Kecukupan dan Kemutakhiran data/informasi dan metodologi (30%)					3	3
Kelengkapan unsur dan kualitas penerbit (30%)					3	2,5
Total = (100%)					10	9
Kontribusi Pengusul (Penulis Pertama /Anggota Utama)	Journal of Indonesian Natural History. Vol.5(2): 16-21. Desember 2017 Penulis ke 3 dari 5 penulis. Nilai maksimal 90%. Nilai pengusul: $(0,4 \times 0,9 \times 10)/4 = 0,9$					0,9

KOMENTAR/ULASAN PEER REVIEW

• Kelengkapan dan Kesesuaian Unsur:	Format lengkap memenuhi unsur tulisan ilmiah. Similaritas relatif tinggi walaupun belum melewati batas.
• Ruang Lingkup dan Kedalaman Pembahasan:	Ruang lingkup masih dalam bidang ilmunya. Pembahasan kurang menyeluruh.
• Kecukupan & Kemutakhiran Data & Metodologi:	Data cukup memenuhi. Metode umum dilakukan peneliti.
• Kelengkapan Unsur & Kualitas Penerbit:	Penerbit Jurusan Biologi FMIPA Universitas Andalas memenuhi syarat penerbit.

Yogyakarta, 11 Juli 2020

Penilai 2

tanda tangan 

Prof. Dr. Suwarno Hadisusanto

NIP 195411161983031002

Unit Kerja : Fakultas Biologi UGM

Bidang Ilmu : Biologi

Jabatan/Pangkat : Guru Besar/ Pembina Utama Madya