

Entomopathogenic fungi of soils of freshwater swamps, tidal lowlands, peatlands, and highlands of South Sumatra, Indonesia

AYU SAFITRI¹, SITI HERLINDA^{2,3,*}, ARUM SETIAWAN⁴

¹Department of Environmental Science, Graduate Program, Universitas Sriwijaya. Palembang 30139, South Sumatra, Indonesia

²Department of Plant Pest and Disease, Faculty of Agriculture, Universitas Sriwijaya, Indralaya, Ogan Ilir 30662, South Sumatra, Indonesia.
Tel./fax. +62-711-580663, *email: sitiherlinda@unsri.ac.id

³Research Center for Sub-optimal Lands (PUR-PLSO), Universitas Sriwijaya. Palembang 30139, South Sumatra, Indonesia

⁴Department of Biology, Faculty of Mathematic and Natural Sciences, Universitas Sriwijaya. Indralaya 30662, South Sumatra, Indonesia

Manuscript received: 11 June 2018. Revision accepted: 22 November 2018.

Abstract. Safitri A, Herlinda S, Setiawan A. 2018. *Entomopathogenic fungi of soils of freshwater swamps, tidal lowlands, peatlands, and highlands of South Sumatra, Indonesia. Biodiversitas 19: 2365-2373.* Ecosystems of lowlands and highlands in South Sumatra have specific characteristics of soils and vegetations that can affect the availability of entomopathogenic fungi. This study aimed to explore and identify species and to determine inoculum potentials of the entomopathogenic fungi from soils of freshwater swamps, tidal lowlands, peatlands, and highlands. Baiting of entomopathogenic fungi on soil samples used the larvae of *Tenebrio molitor*. The entomopathogenic fungi species found in this research were *Beauveria bassiana* and *Metarhizium anisopliae*. The number of the fungal isolates were 30 isolates consisting of nine isolates of *B. bassiana* and 21 isolates of *M. anisopliae*. The highest number of isolates was found in the highland ecosystem (11 isolates) and the lowest was found in peatland ecosystem (4 isolates). The highest percentage average of inoculum potentials of the fungi was found in the high land ecosystem (4.04%) and the lowest one was found in freshwater swamps ecosystem (2.11%). Based on the vegetation type, the soil planted with mustard in Talang Patai-Pagaralam (highland ecosystem) had the highest inoculum potentials (9.33%). These fungi will make an important contribution to the biological control for insect pests in lowland to highland ecosystems in Indonesia.

Keywords: *Beauveria bassiana*, lowlands, *Metarhizium anisopliae*, peat soils, wetlands

INTRODUCTION

Many parts of lowlands and highlands of Indonesia are used for agriculture. Lowlands consisting of freshwater swamps, tidal lowlands, and peatlands are the ecosystems having water saturated condition for the whole year (Sudana 2005). The lowland ecosystems for agricultural purposes in Indonesia are distributed in Sumatra, Kalimantan, and Papua Islands covering 11 million ha of tidal lowlands, comprising 9.2 million ha of freshwater swamps, and 14.9 million ha of peatlands, respectively (Mulyani and Sarwani 2013). While the highland ecosystems managed for agriculture with the area of 16.15 million ha are distributed all over Indonesia islands (Center for Agriculture Data and Information System, Secretariat General 2017).

Soils between lowlands and highlands in Indonesia have different characteristics, especially in the moisture, texture, and acidity (pH). The specific characteristics of lowland and highland soils at the four typologies of ecosystems consist of the freshwater swamp, tidal lowlands, peatlands, and highland closely related to specific cultivated vegetations or crops. Paddy usually is cultivated at freshwater swamps (Herlinda et al. 2018; Lakitan et al. 2018a; Lakitan et al. 2018b). There is only a small part of soils in peatlands that can be cultivated with seasonal crops and most peatlands in Indonesia are utilized for forestry and

conservation areas (Suriadikarta and Sutriadi 2007). Highlands area usually planted with various types of seasonal and annual crops. Specific vegetation or crop plants could affect the soil microorganisms associated with plant and crop roots (El-Ghany 2015).

Microorganisms such as fungi are found in freshwater swamps and highlands of South Sumatra and can be used to control insect pest called entomopathogenic fungi such as *Beauveria bassiana* and *Metarhizium anisopliae* (Herlinda et al. 2008; 2010). The entomopathogenic fungi are proven to be an effective agent to control some insect pests (Chinniah et al. 2016), and they are not harmful toward natural enemies of insect pests (Gholamzadeh-Chitgar et al. 2017). However, there is no complete information for fungi of tidal lowlands and peatlands in that region. *B. bassiana* and *M. anisopliae* were found to kill insect pests such as *Crocidolomia pavonana* (Hasyim et al. 2009), *Aphis gossypii* (Herlinda 2010), *Plutella xylostella* (Loc and Chi 2007), *Lygus* spp. (Leland et al. 2005), and *Oryctes rhinoceros* (Moslim et al. 2009). Entomopathogen, such as entomopathogenic fungi can be explored in soils (El-Ghany 2015). Certain entomopathogen species have adaptation capability in certain soils (Bugeme et al. 2008). Entomopathogen already adapted in freshwater swamp soils or highland soils generally has the specific advantage, i.e., it can adapt more effectively in this soil environment (Erler and Ates 2015). Exploration of entomopathogen starting

from lowland to highland ecosystems will produce the high variation of species and genetics that can be utilized in the extended area and specific location. The purposes of this study were to explore and identify species and to determine inoculum potentials of the entomopathogenic fungi of soils of freshwater swamps, tidal lowlands, peatlands, and highlands in South Sumatra.

MATERIALS AND METHODS

Study area

Soil sampling was carried out in December 2017. The study sites were selected in the agricultural center with specific typology ecosystems in some locations of South Sumatra (Figure 1 and Table 1). The sites consisted of freshwater swamps, tidal lowlands, peatlands, and highlands. Those ecosystems chosen for explorations of entomopathogenic fungi aimed to obtain the fungi adapting in lowland and highland soils. The selected freshwater swamp locations were in Gandus and Musi Subdistricts of Palembang City; Rambutan Subdistrict of Banyuasin District, and Pemulutan Subdistrict of Ogan Ilir District. The selected locations of tidal lowlands were Mulya Sari, Telang Sari, Muara Sungsang Subdistricts of Banyuasin District. The selected peatland locations were in Talang Dabok, Sepucuk, and Kedaton Subdistricts of Ogan Komering Ilir District. The selected highland locations were in Lematang, Tanjung Payang, and Pulau Pinang of Lahat District, and Rimau and Talang Patai of Pagaralam City.

Fungal isolation, purification, and identification were conducted from January to March 2018 in the Laboratory of Entomology, Department of Plant Pest and Disease, College of Agriculture, Sriwijaya University, Indonesia. Supporting data were recorded in terms of soil sampling period, village or city names, coordinate points, and vegetation types or crop plants for each exploration locations (Figure 1 and Table 1). The pH of soil samples was measured by using a method of Kartika et al. (2018). In this survey, the application of pesticides in the sampling locations was also monitored.

Procedures

Observation of inoculum potentials of entomopathogenic fungi in the soil

Inoculum potential observation of the entomopathogenic fungi in this research was conducted by exploration of fungi in soils of the freshwater swamps, tidal lowlands, peatlands, and highlands. Entomopathogenic fungi exploration was carried out by modifying the methods of Anwar et al. (2015), i.e., using newly-moulting-third instar of *Tenebrio molitor* (Yellow mealworm beetle) as insect bait (*Tenebrio* bait method) which was infested on fungal inoculum of soil samples.

The soil samples were collected from some crop fields in the sampling locations (Table 1). Soils were taken from each location at the diagonal position. Soil samples of 5000 g were obtained by digging soil at the depth of 5-15 cm in the vicinity of crop roots, and then they were put into the plastic pouch and labeled with the information of period of the sampling and crop plants.




Figure 1. Locations of exploration for entomopathogenic fungi of soils in ecosystems of South Sumatra, Indonesia: point 1 = tidal lowlands (S 02°40.866' E 104°44.298'), point 2 = freshwater swamps (S 03°02.581' E 104°51.231'), point 3 = peatlands (S 03°24.840' E 104°53.362'), and point 4 = highlands (S 03°48.063' E 103°32.072' and S 03°50.174' E 103°31.293')

Table 1. Locations of exploration for entomopathogenic fungi in South Sumatra, Indonesia

Ecosystems	Village or city	Vegetation or crop plants	Height from sea level (m)
Freshwater swamps	Rambutan, Banyuasin	Paddy	15
	Gandus, Palembang	Paddy	12
	Musi Dua, Palembang	Paddy	16.67
	Pemulutan, Ogan Ilir	Paddy	22
Tidal lowlands	Mulya Sari, Banyuasin	Corn, paddy, watermelon	16.33
	Telang Sari, Banyuasin	Corn, coconut, corn + coconut	18.33
	Muara Sungsang, Banyuasin	Coconut, banana, pineapple	15
Peatlands	Talang Dabok, Ogan Komering Ilir	Palm, rubber, pineapple	24
	Sepucuk, Ogan Komering Ilir	Oil palm, rubber, pineapple	23
	Kedaton, Ogan Komering Ilir	Oil palm	19.67
Highlands	Talang Patai, Pagaram	Cabbage	170
	Pulau Pinang, Lahat	Rubber + coffee	161
	Tanjung Payang, Lahat	Rubber	121
	Lematang, Lahat	Paddy	121.3
	Rimau, Pagaram	Tea	1,326

Previously, soil samples were sieved by using 5-mesh sieve to separate them from crop roots. Then, samples were put into plastic tray (size of 32 cm x 25 cm x 5 cm) and yielded 1000 g of finer soil sample. The finer soil samples were subsequently moistened with sterile aquadest until they obtained the soil moisture of 80-90% using the method of Chen et al. (2014). The number of larvae used per sampling location was 150 larvae (5 plastic trays each containing 30 larvae) (Table 1), so the total larvae per ecosystem type was as follows: in the freshwater swamps it was 1,800 (150 larvae x 12 sampling locations), in tidal lowlands it was 1,350 (150 larvae x 9 sampling locations), in peatlands it was 1,350 (150 larvae x 9 sampling locations), and in highlands it was 2,250 (150 larvae x 15 sampling locations).

The larvae were sterilized with 70% alcohol, and put on soil-sample surface in a plastic tray. The body of larvae was sprinkled using soil sample of about 5 mm thick. Subsequently, the plastic trays containing soil samples were covered with black cloth and sprayed with sterile aquadest to maintain humidity of soil samples. The larvae were incubated within soil sample for seven days to provide enough time for entomopathogenic fungi to infect them. Then, the dead larvae infected by the entomopathogenic fungi were recorded daily to determine inoculum potential. The inoculum potentials of entomopathogenic fungi in this research were measured according to the percentage of infected *Tenebrio* bait or hosts (Hofgaard et al. 2016). The inoculum potential (IP) was calculated using the equation below:

$$IP = \frac{ib}{tb} \times 100$$

ib was the number of infected *Tenebrio* bait, and *tb* was the total number of *Tenebrio* baits.

Entomopathogen isolation

The dead *Tenebrio* baits were subsequently isolated and purified by using the methods of Herlinda (2010).

Entomopathogenic fungi infecting and growing on integument of *Tenebrio* bait were isolated and grown on medium of Sabouraud Dextrose Agar (SDA, Merck). The integument surface of larvae infected by the entomopathogenic fungi was previously sterilized using the modified method of Nuraini et al. (2017) with 1% sodium hypochlorite for 3 minutes and subsequently was rinsed three times with sterile aquadest and air dried on sterile filter paper. The larvae were put into a Petri dish containing sterile humid tissue paper and then incubated in order to stimulate the growth of entomopathogenic fungi. Conidia of entomopathogenic fungi emerging from the dead larvae body were taken by using sterile ose needle and moved into a Petri dish containing SDA medium, and incubated for seven days at the constant temperature of 25 °C within the incubator.

Identification of Entomopathogen Fungi

The purified fungi were identified according to macroscopic and microscopic characteristics using the method of Guilherme et al. (2015). Fungi growing on SDA medium with the area of 1 cm² were taken by using ose needle and put into preparations containing SDA medium, incubated for three days and then microscopically observed. Furthermore, fungi were identified by using books of Humber (2005) and El-Ghany (2015).

Data analysis

Data on inoculum potentials based on percentage of *Tenebrio* bait which was infected by the entomopathogenic fungi among the treatments were analyzed descriptively.

RESULTS AND DISCUSSION

Insect characteristics infected by entomopathogenic fungi

The fungi obtained in this study were identified as *B. bassiana* and *M. anisopliae*. There were only 30 isolates found from these two fungal species (Table 2). These 30 isolates were isolated from 223 infected larvae of *Tenebrio*

bait by the entomopathogenic fungi (Table 3-7); however a lot of larvae were failed to be isolated due to contamination mostly by aerial fungi and *Trichoderma* spp. These 30 isolates consisted of nine isolates of *B. bassiana* and 21 isolates of *M. anisopliae*.

Tenebrio bait infected by the entomopathogenic fungi showed macroscopic and microscopic characteristics or symptoms that could be used to confirm the determination of the entomopathogenic fungal species. Sick or dead *Tenebrio* bait infected by *B. bassiana* showed symptoms as follows: insect body was dry and wrinkle, its outer integument was coated by white mycelia similar to silk, rigid, easily broken and no smell (Figure 2). The *Tenebrio* bait attacked by *M. anisopliae* showed symptoms as follows: dry and wrinkle, no smell, brittle and easily broken, but its outer integument was coated by mycelia having greenish white to dark green or dark color (Figure 3).

The pure isolate was obtained from dead *Tenebrio* bait body which was attacked by entomopathogenic fungi with colony characteristics for each species as follows; colony of *B. bassiana* had the white color similar to cotton, but gradually its color changed into yellowish white as fungi become older. Colony of *M. anisopliae* initially had white color similar to color of *B. bassiana*, but the color changed into greenish and dark green or dark as fungi become older (Figure 4). Conidia of *B. bassiana* and *M. anisopliae* with specific characteristics were obtained from each colony species of entomopathogenic fungi. Conidia of *B. bassiana* has a single cell and a round shape, whereas conidia of *M. anisopliae* also has a single cell but with a cylindrical shape (Figure 5). Mycelia of *B. bassiana* and *M. anisopliae* insulate with upright, branches, and layers of conidiophores.

Inoculum potentials of entomopathogenic fungi in the soil of South Sumatra

Inoculum potentials of entomopathogenic fungi in this research were measured according to the percentage of infected *Tenebrio* bait. The inoculum potentials of entomopathogenic fungi derived from the freshwater swamp soil, tidal lowlands, peat soils, and high land each was different among the survey locations (Tables 3-6). The value of inoculum potential in soil of freshwater swamps, tidal lowlands, peatlands and high lands were in the range of 0.67-3.33%, 2.67-7.33%, 0-4%, and 1.33-9.33%, respectively.

From the freshwater swamp soils, we only could obtain five isolates of the entomopathogenic fungi isolated from 35 infected *tenebrio* baits (Table 3). A lot of the infected *Tenebrio* baits were failed to be isolated due to the contamination by other fungi. The inoculum potentials of the entomopathogenic fungi from the tidal lowlands gained were only 10 isolates (Table 4). The peat soils in sepucuk, Ogan Komering Ilir were planted with rubber had higher inoculum potentials (Table 5). The highland soils in Talang Patai, Pagaralam were planted with mustard also had higher inoculum potential (Table 6). The highest percentage of inoculum potentials of the fungi was found in the highland ecosystem and the lowest one was found in

the freshwater swamp ecosystem. However, the highest number of isolates was found in highland ecosystem (11 isolates) and the lowest one found in peatland ecosystem (4 isolates) (Table 7).



Figure 2. *Tenebrio* bait infected by *Beauveria bassiana* (A) and healthy *Tenebrio* (B)



Figure 3. *Tenebrio* bait infected by *Metarhizium anisopliae* (A) and healthy *Tenebrio* (B)



Figure 4. Colony of *Beauveria bassiana* (A) and *Metarhizium anisopliae* (B)




Figure 5. Conidia of *Beauveria bassiana* (A) and *Metarhizium anisopliae* (B) (400x magnification)

Table 2. Species and isolates of entomopathogenic fungi found in South Sumatra, Indonesia

Ecosystems	Species of fungi	Number of isolate	Vegetation or crop plants	Village or city
Freshwater swamps	<i>M. anisopliae</i>	3	Paddy	Rambutan
Freshwater swamps	<i>M. anisopliae</i>	1	Paddy	Pemulutan
Freshwater swamps	<i>B. bassiana</i>	1	Paddy	Rambutan
Tidal lowlands	<i>M. anisopliae</i>	1	Corn + coconut	Telang Sari
Tidal lowlands	<i>M. anisopliae</i>	2	Corn	Telang Sari
Tidal lowlands	<i>M. anisopliae</i>	2	Corn	Telang Sari
Tidal lowlands	<i>M. anisopliae</i>	3	Paddy	Mulya Sari
Tidal lowlands	<i>B. bassiana</i>	1	Corn	Telang Sari
Tidal lowlands	<i>B. bassiana</i>	1	Watermelon	Mulya Sari
Peatlands	<i>B. bassiana</i>	4	Oil palm	Talang Dabok
Highlands	<i>M. anisopliae</i>	1	Rubber + coffee	Pulau Pinang
Highlands	<i>M. anisopliae</i>	4	Cabbage	Talang Patai
Highlands	<i>M. anisopliae</i>	4	Mustard	Talang Patai
Highlands	<i>B. bassiana</i>	1	Rubber + coffee	Pulau Pinang
Highlands	<i>B. bassiana</i>	1	Cabbage	Talang Patai

Table 3. Inoculum potentials of entomopathogenic fungi in the freshwater swamp soils of South Sumatra, Indonesia

Village or city	Vegetation or crop species	GPS (coordinat)	Inoculum potentials (%)			Fungi (total)
			Number of isolate	<i>B. bassiana</i>	<i>M. anisopliae</i>	
Rambutan, Banyuasin	Paddy	S 03°02.581', E 104°51.231'	0	0.00 (0)	0.67 (1)	0.67 (1)
Rambutan, Banyuasin	Paddy	S 03°02.591', E 104°51.217'	2	0.67 (1)	2.00 (3)	2.67 (4)
Rambutan, Banyuasin	Paddy	S 03°02.586', E 104°51.201'	2	0.67 (1)	0.67 (1)	1.33 (2)
Gandus, Palembang	Paddy	S 03°00.401', E 104°42.380'	0	1.33 (2)	2.00 (3)	3.33 (5)
Gandus, Palembang	Paddy	S 03°00.632', E 104°42.532'	0	0.67 (1)	0.67 (1)	1.33 (2)
Gandus, Palembang	Paddy	S 03°00.632', E 104°42.801'	0	1.33 (2)	1.33 (2)	2.67 (4)
Musi Dua, Palembang	Paddy	S 03°02.120', E 104°43.021'	0	0.67 (1)	1.33 (2)	2.00 (3)
Musi Dua, Palembang	Paddy	S 03°02.150', E 104°43.612'	0	1.33 (2)	2.00 (3)	3.33 (5)
Musi Dua, Palembang	Paddy	S 03°02.510', E 104°43.120'	0	1.33 (2)	1.33 (2)	2.67 (4)
Pemulutan, Ogan Ilir	Paddy	S 03°03.148', E 104°46.230'	0	0.67 (1)	0.67 (1)	1.33 (2)
Pemulutan, Ogan Ilir	Paddy	S 03°03.115', E 104°46.218'	0	0.67 (1)	0.67 (1)	1.33 (2)
Pemulutan, Ogan Ilir	Paddy	S 03°03.113', E 104°46.201'	1	1.33 (2)	1.33 (2)	2.67 (4)

Table 4. Inoculum potentials of entomopathogenic fungi in the tidal lowland soils of South Sumatra, Indonesia

Village or city	Vegetation or crop species	GPS (coordinat)	Inoculum potentials (%)			Fungi (total)
			Number of isolate	<i>B. bassiana</i>	<i>M. anisopliae</i>	
Mulya Sari, Banyuasin	Corn	S 02°40.866', E 104°44.298'	2	1.33 (2)	2.67 (4)	4.00 (6)
Mulya Sari, Banyuasin	Paddy	S 02°40.944', E 104°44.621'	3	1.33 (2)	2.00 (3)	3.33 (5)
Mulya Sari, Banyuasin	Watermelon	S 02°40.896', E 104°44.676'	1	1.33 (2)	2.00 (3)	3.33 (5)
Telang Sari, Banyuasin	Corn	S 02°38.842', E 104°45.369'	2	1.33 (2)	2.67 (4)	4.00 (6)
Telang Sari, Banyuasin	Coconut	S 02°38.813', E 104°45.801'	1	0.67 (1)	1.33 (2)	2.00 (3)
Telang Sari, Banyuasin	Corn + coconut	S 02°38.875', E 104°44.495'	1	2.00 (3)	5.33 (8)	7.33 (11)
Muara Sungsang, Banyuasin	Coconut	S 02°21.736', E 104°50.635'	0	1.33 (2)	4.00 (6)	5.33 (8)
Muara Sungsang, Banyuasin	Banana	S 02°21.823', E 104°50.632'	0	0.67 (1)	2.00 (3)	2.67 (4)
Muara Sungsang	Pineapple	S 02°22.542', E 104°50.324'	0	0.67 (1)	2.00 (3)	2.67 (4)

Table 5. Inoculum potentials of entomopathogenic fungi in the peatland soils of South Sumatra, Indonesia

Village or city	Vegetation or crop plants	GPS (coordinat)	Inoculum potentials (%)			
			Number of isolate	<i>B. bassiana</i>	<i>M. anisoplia</i>	Fungi (Total)
Talang Dabok, Ogan Komering Ilir	Oil palm	S 03°23.570', E 104°51.498'	4	3.33 (5)	1.33 (2)	4.67 (7)
Talang Dabok, Ogan Komering Ilir	Rubber	S 03°25.673', E 104°53.258'	0	2.67 (4)	1.33 (2)	4.00 (6)
Talang Dabok, Ogan Komering Ilir	Pineapple	S 03°25.280', E 104°52.940'	0	2.00 (3)	0.67 (1)	2.67 (4)
Sepucuk, Ogan Komering Ilir	Oil palm	S 03°24.840', E 104°53.362'	0	2.00 (3)	1.33 (2)	3.33 (5)
Sepucuk, Ogan Komering Ilir	Rubber	S 03°23.715', E 104°52.275'	0	3.33 (5)	2.00 (3)	5.33 (8)
Sepucuk, Ogan Komering Ilir	Pineapple	S 03°23.535', E 104°51.780'	0	1.33 (2)	1.33 (2)	2.67 (4)
Kedaton, Ogan Komering Ilir	Oil palm	S 03°23.308', E 104°51.487'	0	1.33 (2)	2.00 (3)	3.33 (5)
Kedaton, Ogan Komering Ilir	Oil palm	S 03°23.277', E 104°51.398'	0	1.33 (2)	0.67 (1)	2.00 (3)
Kedaton, Ogan Komering Ilir	Oil palm	S 03°23.204', E 104°51.459'	0	1.33 (2)	0.67 (1)	2.00 (3)

Table 6. Inoculum potentials of entomopathogenic fungi in the highland soils of South Sumatra, Indonesia

Village or city	Vegetation or crop plants	GPS (coordinat)	Inoculum potentials (%)			
			Number of isolate	<i>B. bassiana</i>	<i>M. anisoplia</i>	Fungi (total)
Pulau Pinang, Lahat	Rubber + coffee (A)	S 03°48.819', E 103°32.677'	2	1.33 (2)	1.33 (2)	2.67 (4)
Pulau Pinang, Lahat	Rubber + coffee (B)	S 03°48.852', E 103°32.698'	0	1.33 (2)	2.00 (3)	3.33 (5)
Pulau Pinang, Lahat	Rubber + coffee (C)	S 03°48.883', E 103°32.688'	0	0.67 (1)	0.67 (1)	1.33 (2)
Tanjung Payang, Lahat	Rubber (A)	S 03°48.094', E 103°32.145'	0	0.67 (1)	1.33 (2)	2.00 (3)
Tanjung Payang, Lahat	Rubber (B)	S 03°48.122', E 103°32.162'	0	3.33 (5)	2.67 (4)	6.00 (9)
Tanjung Payang, Lahat	Rubber (C)	S 03°48.177', E 103°32.166'	0	2.00 (3)	2.67 (4)	4.67 (7)
Lematang, Lahat	Paddy	S 03°48.063', E 103°32.072'	0	0.67 (1)	1.33 (2)	2.00 (3)
Lematang, Lahat	Paddy	S 03°48.051', E 103°32.069'	0	3.33 (5)	4.00 (6)	7.33 (11)
Lematang, Lahat	Paddy	S 03°48.020', E 103°32.064'	0	1.33 (2)	1.33 (2)	2.67 (4)
Talang Patai, Pagaram	Cabbage	S 03°50.180', E 103°31.325'	2	4.00 (6)	4.67 (7)	8.67 (13)
Talang Patai, Pagaram	Cabbage	S 03°50.174', E 103°31.313'	3	0.67 (1)	1.33 (2)	2.00 (3)
Talang Patai, Pagaram	Mustard	S 03°50.174', E 103°31.293'	4	4.67 (7)	4.67 (7)	9.33 (14)
Rimau, Pagaram	Tea	S 04°02.161', E 103°10.484'	0	2.00 (3)	2.00 (3)	4.00 (6)
Rimau, Pagaram	Tea	S 04°02.144', E 103°10.487'	0	1.33(2)	1.33(2)	2.67 (4)
Rimau, Pagaram	Tea	S 04°02.136', E 103°10.485'	0	0.67 (1)	1.33(2)	2.00 (4)

Table 7. Inoculum potentials of entomopathogenic fungi in the soil of South Sumatra, Indonesia

Ecosystems	Inoculum potentials (%)			
	Number of isolate	<i>B. bassiana</i>	<i>M. anisoplia</i>	Fungi (total)
Freshwater Swamps	5	0.89 (16)	1.22 (22)	2.11 (38)
Tidal Lowlands	10	1.18 (16)	2.67 (36)	3.85 (52)
Peatlands	4	2.07 (28)	1.25 (17)	3.33 (45)
Highlands	11	1.87 (42)	2.17 (49)	4.04 (91)

Discussion

Entomopathogenic species found in this research were *B. bassiana* and *M. anisopliae*. Macroscopic and microscopic characteristics of both entomopathogenic fungi found in this research matched with the previous studies. Humber (2005) stated that mycelia of *B. bassiana* appeared from exoskeleton of hosts insect, and covered all parts of the exterior surface of host integument, so that host body has white color, reverse pale to yellow colony, hyaline or colorless, single cell, and globular and conidia as well as insulate hypha. *M. anisopliae* causes integument

which has colors of whitish green to dark green because its mycelia cover exoskeleton of hosts insect; it has green to yellow conidia, single cell and cylindrical conidia as well as insulate hypha (Driver et al. 2000; Humber 2005).

Beauveria bassiana and *M. anisopliae* have parasitic and saprophytic phases during the killing process of their host insect (Augustyniuk-Kram and Kram 2012; El-Ghany 2015). Parasitic phase starts viz, the fungal conidia attach to the host insect cuticle, and then the conidia germinate on the host cuticle (El-Ghany 2015). The fungal penetration into the insect cuticle can be performed in producing

specific infection hyphae originating at appressoria of the fungus (Fernandes et al. 2007; El-Ghany 2015). Gürlek et al. (2018) reported that both species could produce germ tubes growing over the surface of the insect cuticle until the tubes contacted weakness area of cuticle where penetration could easily be achieved. After the fungus successfully penetrated, then mycelia distributed into the hemolymph by the formation of blastospores (El-Ghany 2015). Finally, the host insect would die within four days of penetration (Gürlek et al. 2018). Saprophytic phase starts viz, the fungus grew mechanically in the dead insect body, retrieved nutrients from the insect body, and then the fungus produced toxins (El-Ghany 2015).

The success of both fungi in conducting the process of parasitic and saprophytic phases was affected by several external factors such as moisture, pH, temperature, ultraviolet (UV) radiation, and vegetation (El-Ghany 2015). This research showed that highland soil planted with cabbage and mustard in Talang Patai, Pagaram had more inoculum potentials of entomopathogenic fungi than other locations because cabbage and mustard were inhabited by insect pests dominated by Lepidoptera, such as *Plutella xylostella*, *Crociodomia binotalis*, *Spodoptera litura*, and *Chrysodeixis chalcites*; while the larvae of Lepidoptera were the most suitable hosts for entomopathogenic fungi (Godonou et al. 2009; Nunilahwati et al. 2012). This study also found that a lot of infected larvae of the insect pests hung above of the mustard and cabbage canopy. The host insects attacked by entomopathogenic fungi on the cabbage canopy at highlands, South Sumatra were generally dominated by *P. xylostella*, *S. litura*, and *C. chalcites*. The symptoms of sick or infected insect larvae were dry and in the stiff condition, white or greenish white in color and attached on the upper surface of cabbage leaves. The infected larvae which showed the symptoms of dry and stiff condition as mummy and covered with white fungal mycelia was larvae attacked by *B. bassiana*, whereas larvae body covered by fungal mycelia having greenish white or dark green color was symptoms of larvae attacked by *M. anisopliae*. Symptoms of larvae attacked by *B. bassiana* and *M. anisopliae* in this research matched to the symptoms reported by El-Ghany (2015) and Mora et al. (2017).

In this research, more inoculum potentials of the entomopathogenic fungi were found in the highlands and tidal lowlands than that in freshwater swamps and peatlands because it was affected by soil pH and moisture. Zhong et al. (2010) reported that soil pH had a more significant role in determining the existence of fungal propagules within soils than that of soil texture and organic matter. However, Inubushi et al. (2003) had stated that soil moisture is one of the most important controlling factors for biological reactions in the soil. Soil pH in this research was in the range of 4 to 4.5 in freshwater swamps, 4.3 to 5 in tidal lowlands, 3.60 to 4 in peatlands, and 5 to 6.7 in highlands. Kodir and Juwita (2016) stated that the pH value of soil in freshwater swamps in Indonesia are in the ranges of 4 to 4.5, pH 4.17 to 5.35 in tidal lowlands (Marlina et al. 2016), and 3.60 to 3.95 in peatlands (Utami et al. 2009), and 5 to 6 in highlands (Supriadi et al. 2016). The variation of

pH values of soils at each location could affect the adaptation capability of entomopathogenic fungi surviving (Bugeme et al. 2008). The conidial viability of entomopathogenic fungi, such as *B. bassiana* and *M. anisopliae* were affected by pH of in-vitro medium for entomopathogenic fungi. Rizkie et al. (2017) reported that high acidity (pH < 4) of in-vitro medium for fungus growing significantly decrease the conidial viability of *B. bassiana* and *M. anisopliae*. Therefore, inoculum potentials of the entomopathogenic fungi from peatland soil and freshwater swamp soil were lower than that from the soil in the tidal lowlands and highlands.

In addition to soil pH and moisture, soil texture also determines the existence and distribution of fungal propagules. Soil texture has low clay content, and sandy soil texture tends to have the low capability in maintaining the existence of fungal propagules (El-Ghany 2015) as well as the water saturated soil (Garrido-Jurado et al. 2011). Soils from freshwater swamps and peatlands in South Sumatra had lower clay content and in water saturated condition (Marlina et al. 2016; Kartika et al. 2018). The low existence of fungal propagules finding at freshwater swamp and peatland soils was due to both factors. Soils in freshwater swamps are water saturated for more than 6 to 7 months per year usually occurred from November to April (Herlinda et al. 2018) and soil moisture in the peatlands reach 500% (Maftu'ah and Susanti 2009).

Peatlands have soil pore saturated with water all year long resulting in the anaerobic condition of the soil. Peat soil has no clay, sand, and silt content, but it has organic matter (Sudana 2005). The high organic matter finally can decrease the pH of the soil (Utami et al. 2009). Rizkie et al. (2017) confirmed that pH < 4 within medium in-vitro for growing fungi could decrease the ability to live of fungal propagules. In highland ecosystems, the portion of the soil texture among clay, sand, and silt fractions was found in the same composition or balance (Utomo et al. 2013; Supriadi et al. 2016). While tidal lowlands contain silt from sedimentation mixture of river water and seawater with balance clay and sand fractions (Marlina et al. 2016). Higher inoculum potentials in highlands and tidal lowlands in this research was due to balance or higher of clay texture and organic soils than that of freshwater swamps and peatlands.

The balance or higher of clay texture and organic soils are capable of maintaining the existence of fungal propagules (Garrido-Jurado et al. 2011). Zhong et al. (2010) stated that soil in ecosystems which apply composted manure or no synthetic fertilizer had higher propagules content of *B. bassiana* than that of soil in the ecosystem which applies synthetic fertilizers. Also, the application of synthetic pesticides is capable of decreasing the existence of entomopathogenic fungi within the soil (Mietkiewski et al. 2010). Local farmers in freshwater swamp and peatland areas of South Sumatra usually do not apply synthetic pesticides, whereas many local farmers in tidal lowland and highland areas apply synthetic pesticides (Herlinda et al. 2018). Although no synthetic pesticides were applied in the freshwater swamp and peatland areas, fungal propagules or inoculum potentials in these areas were lower than those of tidal lowlands and highlands

areas. In this study, there was no evidence that having no synthetic pesticide application in freshwater swamp and peatland areas can cause the high existence of fungal propagules. However, soil pH and soil texture have more effect on the existence of propagules of *B. bassiana* and *M. anisopliae* in lowland swamp and peatland ecosystems.

Higher inoculum potentials in highlands and tidal lowlands in this research were closely related to specific cultivated vegetations or crops. Most peatlands in Indonesia were utilized for forestry and conservation areas (Suriadikarta and Sutriadi 2007). Paddy generally is cultivated at freshwater swamps (Herlinda et al. 2018; Lakitan et al. 2018a; Lakitan et al. 2018b) as well as in tidal lowlands; however, paddy cultivation at tidal lowlands was more intensive (with two to three planting indices) than that of freshwater swamps (one planting index). Thus, diverse species of vegetation or crop plants could affect abundant and diverse species of the soil microorganisms associated with plant and crop roots (El-Ghany 2015).

This research found two species of entomopathogenic fungi from soils in South Sumatra i.e., *B. bassiana* and *M. anisopliae*. The highest percentage of the inoculum potentials and prevalence of both fungi occurred in the highland ecosystems and the lowest percentage of the inoculum potentials of the fungi was found in the peatland ecosystems. In highland ecosystems, the percentage of the inoculum potentials was affected by the locations and the vegetation or the crop plants. These fungi will make an important contribution to the biological control for insect pests from lowland to highland ecosystems in Indonesia.

ACKNOWLEDGEMENTS

This research was funded by National Strategic Research (PSNI) with budget year of 2018 according to the Director of Research and Community Service, Directorate of Research and Community Service (DRPM), Directorate General for Research and Development, Ministry of Research, Technology, and Higher Education, Number: 093/SP2H/LT/DRPM/IV/2018 chaired by Siti Herlinda. Special thanks to Dr. Suwandi, a microbiologist from Universitas Sriwijaya, Palembang, Indonesia for identification of fungi.

REFERENCES

- Anwar W, Khan SN, Aslam M, Haider MS, Shahid AA, Ali M. 2015. Exploring fungal flora associated with insects of cotton agroecological zones of Punjab, Pakistan. *Pakistan Entomologist* 37: 27-31.
- Augustyniuk-Kram A, Kram KJ. 2012. Entomopathogenic fungi as an important natural regulator of insect outbreaks in forests review. *Forest Ecosystem, More than Just Trees*. Blanco JA. (ed.). In Tech: Croatia
- Bugeme DM, Maniania NK, Knapp M, Boga HI. 2008. Effect of temperature on virulence of *Beauveria bassiana* and *Metarhizium anisopliae* isolates to *Tetranychus evansi*. *Exp Appl Acarol* 46: 275-285.
- Center for Agriculture Data and Information System, Secretariat General. 2017. Statistics of Agricultural Land 2012-2016. In. Hakim M, Wiratno O, and Abdurachman A (eds). Ministry of Agriculture. Jakarta.
- Chen Z, Xu L, Yang F, Ji G, Yang J, Wang J. 2014. Efficacy of *Metarhizium anisopliae* isolate MAX-2 from Shangri-la, China under desiccation stress. *BMC Microbiol* 14: 1-8.
- Driver F, Milner RJ, Trueman JWH. 2000. A taxonomic revision of *Metarhizium* based on a phylogenetic analysis of rDNA sequence data. *Mycol Res* 104: 134-150.
- El-Ghany TMA. 2015. Entomopathogenic Fungi and their Role in Biological Control. Biology Department Faculty of Science Jazan University KSA. Cairo.
- Erlor F, Ates AO. 2015. Potential of two entomopathogenic fungi, *Beauveria bassiana* and *Metarhizium anisopliae* (Coleoptera: Scarabaeidae), as biological control agents against the June beetle. *J Insect Sci* 15: 1-6.
- Fernandes EKK, Rangel DEN, Moraes AM., Bittencourt VREP, Roberts DW. 2007. Variability in tolerance to UV-B radiation among *Beauveria* spp. isolates. *J Invertebr Pathol* 96: 237-243.
- Garrido-Jurado J, Torrent J, Barrón V, Corpas A, Quesada-Moraga E. 2011. Soil properties affect the availability, movement, and virulence of entomopathogenic fungi conidia against puparia of *Ceratitis capitata* (Diptera: Tephritidae). *Biol Control* 58: 277-285.
- Godonou I, James B, Atcha-Ahowé B, Vodouhe S, Kooyman C, Ahanchédé A, Korie S. 2009. Potential of *Beauveria bassiana* and *Metarhizium anisopliae* isolates from Benin to control *Plutella xylostella* L. (Lepidoptera: Plutellidae). *Crop Protection* 28: 220-224.
- Guilherme D, Oliveira P, Pauli G, Moura G, Delalibera I. 2015. A protocol for determination of conidial viability of the fungal entomopathogens *Beauveria bassiana* and *Metarhizium anisopliae* from commercial products. *J Microbiol Methods* 119: 44-52.
- Gürlek S, Sevim A, Sezgin FM, Sevim E. 2018. Isolation and characterization of *Beauveria* and *Metarhizium* spp. from walnut fields and their pathogenicity against the codling moth, *Cydia pomonella* (L.) (Lepidoptera: Tortricidae). *Egyptian J Biol Pest Control* 28: 1-6.
- Hasyim A, Nuraida, Trizelia. 2009. Pathogenicity of entomopathogenic fungi isolates to eggs and larvae of cabbage head caterpillar, *Crociodolomia pavonana* Fabricius. *J Hort* 19: 334-343.
- Herlinda S. 2010. Spore density and viability of entomopathogenic fungal isolates from Indonesia, and their virulence against *Aphis gossypii* Glover (Homoptera: Aphididae). *Tropical Life Sci Res* 21: 11-19.
- Herlinda S, Irsan C, Mayasari R, Septariani S. 2010. Identification and selection of entomopathogenic fungi as biocontrol agents for *Aphis gossypii* from South Sumatra. *Microbiol Indonesia* 4: 137-142.
- Herlinda S, Mulyati SI, Suwandi. 2008. Selection of isolates of entomopathogenic fungi and the bioefficacy of their liquid production against *Leptocorisa oratorius* nymphs. *Microbiol Indonesia* 2: 141-146.
- Herlinda S, Yudha S, Thalib R, Khodijah, Suwandi, Lakitan B, Verawaty M. 2018. Species richness and abundance of spiders inhabiting rice in freshwater swamps and tidal lowlands in South Sumatra, Indonesia. *J ISSAAS* 24: 82-93.
- Hofgaard IS, Seehusen T, Aamot HU, Riley H, Razzaghian J, Le VH, Hjelkrem A-GR, Dill-Macky R, Brodal G. 2016. Inoculum potential of *Fusarium* spp. relates to tillage and straw management in Norwegian fields of spring oats. *Front Microbiol* 7: 1-15.
- Humber RA. 2005. Entomopathogenic Fungal Identification. USDA-ARS Plant Protection Research Unit. Ithaca.
- Inubushi K, Furukawa Y, Hadi A, Purnomo E, Tsuruta H. 2003. Seasonal changes of CO₂, CH₄ and N₂O fluxes in relation to land-use change in tropical peatlands located in coastal area of South Kalimantan. *Chemosphere* 52: 603-608.
- Kartika K, Lakitan B, Wijaya A, Kadir S, Widuri LI, Siaga E. 2018. Effects of particle size and application rate of rice-husk biochar on chemical properties of tropical wetland soil, rice growth and yield. *AJCS* 12: 817-826.
- Kodir KA, Juwita Y. 2016. Morphological characterization and inventory of local wetland rice collected from South Sumatra Province. *Bul Plasma Nutfah* 22: 101-108.
- Lakitan B, Alberto A, Lindiana L, Kartika K, Herlinda S, Kurnianingsih A. 2018. The Benefits of biochar on rice growth and yield in tropical riparian wetland, South Sumatra, Indonesia. *CMUJ Natural Sciences* 17: 111-126.
- Lakitan B, Hadi B, Herlinda S, Siaga E, Widuri LI, Kartika K, Lindiana L, Yunindyawati Y, Meihana M. 2018. Recognizing farmers practices and constraints for intensifying rice production at riparian wetlands in

- Indonesia. NJAS Wageningen J Life Sci 85: 10-20.
- Leland JE, Mcguire MR, Grace JA, Jaronski ST, Ulloa M, Park Y, Plattner RD. 2005. Strain selection of a fungal entomopathogen, *Beauveria bassiana*, for control of plant bugs (*Lygus* spp.) (Heteroptera: Miridae). Biol Control 35: 104-114.
- Loc NT, Chi VTB. 2007. Biocontrol potential of *Metarhizium anisopliae* and *Beauveria bassiana* against diamond back moth, *Plutella xylostella*. Omonrice 93: 86-93.
- Maftu'ah E, Susanti MA. 2009. Earthworms community on several land uses of peat land in Central Kalimantan. Berita Biologi 9:371-378.
- Marlina N, Asmawati, Zairani FY, Midranisiah, Aryani I, Kalasari R. 2016. Biofertilizer utilization in increasing inorganic fertilizer efficiency and rice yield at c-type flooding land of Tanjung Lago tidal lowland. Int J Engg res & Sci & Tech 5: 74-83.
- Mietkiewski RT, Pell JK, Clark SJ. 2010. Influence of pesticide use on the natural occurrence of entomopathogenic fungi in arable soils in the UK: Field and laboratory comparisons. Biocontrol Science and Technology 7: 565-575.
- Mora MAE, Castilho AMC, Fraga ME. 2017. Classification and infection mechanism of entomopathogenic fungi. Arq Inst Biol 84: 1-10.
- Moslim R, Kamarudin N, Wahid MB. 2009. Pathogenicity of granule formulations of *Metarhizium anisopliae* against the larvae of the oil palm rhinoceros beetle, *Oryctes rhinoceros*. J Oil Palm Res 21: 602-612.
- Mulyani A, Sarwani M. 2013. The characteristic and potential of sub-optimal land for agricultural development in Indonesia. Jurnal Sumberdaya Lokal 7: 47-55. [Indonesian]
- Nunilawati H, Herlinda S, Irsan C, Pujiastuti Y. 2012. Exploration, isolation and selection entomopathogenic fungi infectious to *Plutella xylostella* (Lepidoptera: Yponomeutidae) on green mustard (*Brassica chinensis*) crop in South Sumatra. J HPT Tropika 12: 1-11. [Indonesian]
- Nuraini FR, Setyaningsih R, Susilowati ARI. 2017. Screening and characterization of endophytic fungi as antagonistic agents toward *Fusarium oxysporum* on eggplant (*Solanum melongena*). Biodiversitas 18: 1377-1384.
- Rizkie L, Herlinda S, Suwandi, Irsan C, Susilawati, Lakitan B. 2017. Conidial density and viability of *Beauveria bassiana* and *Metarhizium anisopliae* grown on the low-pH in vitro medium. J HPT Tropika 17: 119-127. [Indonesian]
- Sudana W. 2005. The potential and prospect of swamps as a source of agricultural production. Analisis Kebijakan Pertanian 3: 141-151. [Indonesian]
- Supriadi H, Randriani E, Towaha J. 2016. Correlation between altitude, soil chemical properties, and physical quality of arabica coffee beans in highland areas of Garut. J TIDP 3: 45-52.
- Suriadikarta DA, Sutriadi MT. 2007. Potential tidal land for agricultural development. The Jurnal Litbang Pertanian 26: 115-122. [Indonesian]
- Utami SNH, Maas A, Radjaguguk B, Purwanto BH. 2009. Nature on soil physical, chemical and ftir spectrophotometry of hydrophobic peat from Central Kalimantan. J Trop Soils 14: 159-166.
- Utomo M, Banuwa IS, Buchari H, Anggraini Y. 2013. Long-term tillage and nitrogen fertilization effects on soil properties and crop yields. J Trop Soils 18: 131-139.
- Zhong W, Gu T, Wang W. 2010. The effects of mineral fertilizer and organic manure on soil microbial community and diversity. Plant Soil 326: 511-522.



<

Ads by Google




Stop seeing this ad

Why this ad? ⓘ

Biodiversitas

8

H Index









Country	Indonesia -  SJR Ranking of Indonesia
Subject Area and Category	Agricultural and Biological Sciences Animal Science and Zoology Plant Science Biochemistry, Genetics and Molecular Biology Molecular Biology
Publisher	Biology department, Sebelas Maret University Surakarta
Publication type	Journals
ISSN	1412033X, 20854722
Coverage	2014-ongoing
Scope	"Biodiversitas, Journal of Biological Diversity" or Biodiversitas encourages submission of manuscripts dealing with all biodiversit aspects of plants, animals and microbes at the level of gene, species, and ecosystem.
	Homepage How to publish in this journal Contact  Join the conversation about this journal

Publish Research Paper

Call Papers for Scientific Research Journal (Scirj), ISSN: 2201-2796


scirj.org

OPEN



Show this widget in your own website

Just copy the code below and paste within your html code:



F **Fitra Syawal Harahap** 2 weeks ago

I want to submit my manuscript in this journal

reply

SCImago Team



Melanie Ortiz 2 weeks ago

Dear Fitra, thank you very much for your comment, we suggest you to look for author's instructions/submission guidelines in the journal's website. Best Regards, SCImago Team

J **Joko Prasetyo** 2 months ago

I want to submit my manuscript in this journal

reply

SCImago Team



Melanie Ortiz 2 months ago

Dear Joko, thank you very much for your comment, we suggest you to look for author's instructions/submission guidelines in the journal's website. Best Regards, SCImago Team

S **salamiah** 3 months ago

I see the subject area

reply

A **Afik** 7 months ago

Bagus kali jurnal nya

reply



Melanie Ortiz 7 months ago

Dear user, thanks for your participation! Best Regards, SCImago Team

F **Feron** 7 months ago

Wow bagus banget...samangat berkarya untuk Indonesia yang kebih maju

reply



Melanie Ortiz 7 months ago

Dear user, thanks for your participation! Best Regards, SCImago Team



muhamad iksan 2 years ago

i want submit my journal thanks

reply

A **ahmad** 12 months ago

Kindly submit your paper here, <https://smujo.id/biodiv/about/submissions>

Leave a comment

Name

Email

(will not be published)

I'm not a robot

reCAPTCHA
Privacy - Terms

Submit

The users of Scimago Journal & Country Rank have the possibility to dialogue through comments linked to a specific journal. The purpose is to have a forum in which general doubts about the processes of publication in the journal, experiences and other issues derived from the publication of papers are resolved. For topics on particular articles, maintain the dialogue through the usual channels with your editor.

Biological research

Rad Source has the best technology for Cell Research Irradiators around!


radsources.com

OPEN

Developed by:



Powered by:



Follow us on @ScimagoJR

Scimago Lab, Copyright 2007-2020. Data Source: Scopus®

EST MODUS IN REBUS

Horatio (Satire 1.1.106)

[Contact Us](#)
[FAQ](#)
[Useful links](#)
[Home](#)
[Content and Archives](#)
[Aims and Scopes](#)
[Editorial Board](#)
[Guidance for Authors](#)
[Ethical Guidelines](#)
[Charges](#)
[Membership](#)
[Previously Reviewers](#)
[Conference Events](#)

Biodiversitas Vol. 19, No. 6, November 2018

 only search Biodiversitas

Table of Content

[Cover \[PDF\]](#)
[Editorial Board & Guidance for Authors \[PDF\]](#)

1. Short Communication: Second record of *Eumops chiribaya* (Chiroptera, Molossidae) in Peru [\[PDF\]](#)
CESAR EDGARDO MEDINA, DARWIN RONALD DIAZ, BRIAN ANDREW MALAGA, YASMY KARINA MEDINA, EVARISTO LOPEZ
2. Anoa, dwarf buffalo from Sulawesi, Indonesia: Identification based on DNA barcode [\[PDF\]](#)
DWI SENDI PRIYONO, DEDY D. SOLIHIN, ACHMAD FARAJALLAH, DIAH IRAWATI DWI ARINI
3. Genetic diversity and population structure of *Siganus fuscescens* across urban reefs of Seribu Islands, Northern of Jakarta, Indonesia [\[PDF\]](#)
ADITYA BRAMANDITO, BEGINER SUBHAN, TRI PRARTONO, NURLITA PUTRI ANGGRAINI, HEDI INDRA JANUAR, HAWIS H. MADDUPPA
4. Ethnoecology: The traditional landscape of Osing Tribe in Banyuwangi, Indonesia [\[PDF\]](#)
BUDI PRASETYO, TATIK CHIKMAWATI, EKO BAROTO WALUJO, ERVIZAL AMZU
5. Assessing the status of *pinuchu* as indicator of socio-ecological resilience of Chaya Socio-Ecological Production Landscape, Ifugao, Philippines [\[PDF\]](#)
JHOANNA O. SANTIAGO, INOCENCIO E. BUOT JR
6. Spatio-temporal influence of physicochemical parameters on phytoplankton assemblage in coastal brackish lagoon: Gomishan Lagoon, Caspian Sea, Iran [\[PDF\]](#)
MOHAMMAD JABBARI, MORTEZA SALAHI, RASOUL GHORBANI
7. The diversity of endophytic fungi associated with *Piper nigrum* in the tropical areas: A recent study from Kutai Kartanegara, Indonesia [\[PDF\]](#)
SOPIALENA, SUYADI, MUHAMAD SAHIL, JULI NURDIANA
8. Morphometric characteristics of two seagrass species (*Enhalus acoroides* and *Cymodocea rotundata*) in four small islands in North Maluku, Indonesia [\[PDF\]](#)
YUNITA RAMILI, DIETRIECH G. BENGEN, HAWIS H. MADDUPPA, MUJIZAT KAWAROE
9. Ethnobotany of tree ferns in Pasir Menyan Hamlet, Sukamandi Village, Subang, West Java, Indonesia [\[PDF\]](#)
SURYANA, JOHAN ISKANDAR, PARIKESIT, RUHYAT PARTASASMITA
10. Short Communication: Isolation and identification of actinomycetes potential as the antagonist of *Dickeya zeae* pineapple soft rot in Lampung, Indonesia [\[PDF\]](#)
TITIK NUR AENY, JOKO PRASETYO, RADIX SUHARJO, SUSKANDINI R. DIRMAWATI, EFRI, AININ NISWATI
11. Ethnobotany of banana plants (*Musa x paradisiaca*) of Palintang Hamlet, Cipanjalu Village, Bandung, West Java, Indonesia [\[PDF\]](#)
JOHAN ISKANDAR, JOKO KUSMORO, MIRA MUBAROKAH, RUHYAT PARTASASMITA
12. Short Communication: Development of selected PGPR consortium to control *Ralstonia solanaceae* subsp. *indonesiensis* and promote the growth of tomato [\[PDF\]](#)
YULMIRA YANTI, WARNITA, REFLIN, HASMIANDY HAMID
13. Chromosome numbers of some Asteraceae species from Universitas Indonesia Campus, Depok, Indonesia [\[PDF\]](#)
ANDI SALAMAH, RIRIN OKTARINA, EKA AYU AMBARWATI, DWI FAUZIYAH PUTRI, ASTARI DWIRANTI, NOVIAR ANDAYANI
14. The mixed cropping of *Anthocephalus cadamba* and *Glycine max* for rehabilitating sloping lands [\[PDF\]](#)
KARYATI, SRI SARMINAH, KARMINI, GUNAWAN SIMANGUNSONG, JEKSON TAMBA
15. Rapid expansion and biodiversity impacts of the red devil cichlid (*Amphilophus labiatus*, Gunther 1864) in Lake Sentani, Papua, Indonesia [\[PDF\]](#)
HENDERITE L. OHEE, PUGUH SUJARTA, SURIANI BR. SURBAKTI, HOLLY BARCLAY
16. Phytoplankton community structure in the Estuary of Donan River, Cilacap, Central Java, Indonesia [\[PDF\]](#)
HESTI PRATIWI, ARIO DAMAR, SULISTIONO
17. Short Communication: *Cyrtodactylus elok* Dring, 1979 (Sauria, Gekkonidae): A first country record for Thailand [\[PDF\]](#)

YODCHAIY CHUAYNKERN, PECHRKAWIN NURNGSOMSRI, CHANTIP CHUAYNKERN, PRATEEP DUENGKAE, SUNATE KARAPHAN

18. Chromosome numbers of some species of *Pteris* (Pteridaceae) in Java, Indonesia [PDF]
TITIEN NGATINEM PRAPTOSUWIRYO, MUGI MUMPUNI
19. Antifungal activity of soybean rhizosphere actinomycetes producing bioactive compounds against *Fusarium oxysporum* [PDF]
HIMA DEWI MARIASTUTI, SRI LISTIYOWATI, ARIS TRI WAHYUDI
20. A checklist of medically important flies (Order: Diptera) in the central region of Thailand [PDF]
TANAWAT CHAIPHONGPACHARA, SEDTHAPONG LAOJUN, NITIWOOT JONGVISUTTISAN, PATCHARAPRON TUBSAMUT, APICHAYA DASOM
21. Short Communication: Ethnozoology - The perception of Tobelo Dalam tribal community in Saolat Village, East Halmahera, Indonesia towards cuscus (Phalangeridae) [PDF]
ELIZABETH NOVI KUSUMANINGRUM, JATNA SUPRIATNA, ABINAWANTO, ANOM BOWOLAKSONO
22. A checklist of Bombyliidae (Diptera) from Mordovia, Russia and variation of wing shape in *Bombylius* species [PDF]
MARIYA ALEXANDROVNA CHURSINA, ALEXANDER BORISOVICH RUCHIN
23. Molecular characterization of Plant Growth Promoting Rhizobacteria using 16S rRNA sequences in the organic rice field of Sukorejo Village, Central Java, Indonesia [PDF]
SLAMET SANTOSA, SUTARNO, EDI PURWANTO, SURANTO, SAJIDAN
24. Vegetation analysis along the altitudinal gradient of Mt. Ilong, Halcon Range, Mindoro Island, Philippines [PDF]
ELAINE LOREEN C. VILLANUEVA, INOCENCIO E. BUOT, JR.
25. Population genetics and ecology of Sumatran camphor (*Dryobalanops aromatica*) in natural and community-owned forests in Indonesia [PDF]
FAUJIAH NURHASANAH RITONGA, FIFI GUS DWIYANTI, CECEP KUSMANA, ULFAH JUNIARTI SIREGAR, ISKANDAR ZULKARNAEN SIREGAR
26. Planting coffee and take care of forest: A case study on coffee cultivation in the forest carried out among people of Palintang, Highland of Bandung, West Java, Indonesia [PDF]
BUDIAWATI SUPANGKAT ISKANDAR, JOHAN ISKANDAR, RUHYAT PARTASASMITA, RAHMAN LATIF ALFIAN
27. Antimicrobial activities of actinomycete isolates from rhizospheric soils in different mangrove forests of Torosiaje, Gorontalo, Indonesia [PDF]
YULIANA RETNOWATI, SUKARTI MOELJOPAWIRO, TJUT SUGANDAWATY DJOHAN, ENDANG SUTARININGSIH SOETARTO
28. Short Communication: Identification of *Mildew Locus O* (MLO) genes in *Durio zibethinus* genome corresponding with the Powdery Mildew disease [PDF]
RAHMAT AZHARI KEMAL, ERIC BERNARDUS L. SANDJAJA, AUDI PUTRA SANTOSA, JEREMIAS IVAN
29. Short Communication: Trees species diversity and indicator species in Bedugul forest ecosystem, Bali, Indonesia [PDF]
SUTOMO, I DEWA PUTU DARMA, ARIEF PRIYADI, RAJIF IRYADI
30. Ideal planting space for merbau (*Intsia bijuga*) forest plantations in Papua based on distance-dependent competition [PDF]
ELIESER VIKTOR SIRAMI, DJOKO MARSONO, RONGGO SADONO, MUHAMMAD ALI IMRON
31. Knowledge and perception of pelung keepers's toward chicken contests in West Java, Indonesia [PDF]
INDRAWATI Y. ASMARA, DANI GARNIDA, MARINA SULISYATI, SAFITRI TEJANINGSIH, RUHYAT PARTASASMITA
32. Molecular identification, abundance and distribution of the coral-killing sponge *Terpios hoshinota* in Bengkulu and Seribu Islands, Indonesia [PDF]
RISNITA TRI UTAMI, NEVIATY P. ZAMANI, HAWIS H. MADDUPPA
33. Short Communication: Growth and mortality rate of Black Pomfret *Parastromateus niger* (Bloch, 1795) and Silver Pomfret *Pampus argenteus* (Euphrasen, 1788) in Paloh Waters, West Kalimantan, Indonesia [PDF]
ADRIAN DAMORA, DWI ARIYOGAGAUTAMA, RONNY I. WAHJU, HENDRO SUSANTO, JOHN WANG
34. Ichthyofauna checklist (Chordata: Actinopterygii) for indicating water quality in Kampar River catchment, Malaysia [PDF]
CASEY KEAT-CHUAN NG, PETER AUN-CHUAN OOI, WEY-LIM WONG, GIDEON KHOO
35. Bycatch composition of cutlassfish trawlers during fishing season in Bushehr and Hormozgan, Persian Gulf, Iran [PDF]
AHMAD FOULADI SABET, SEYED YOUSEF PAIGHAMBARI, MOJTABA POULADI, HADI RAEISI, REZA ABBASPOUR NADERI
36. Studies on the diversity of macrofungus in Kodaikanal region of Western Ghats, Tamil Nadu, India [PDF]
BOOBALAN THULASINATHAN, MOHAN RASU KULANTHAISAMY, ARUMUGAM NAGARAJAN, SARAVANAN SOORANGKATTAN, JOTHI BASU MUTHURAMALINGAM, JEYAKANTHAN JEYARAMAN4, ALAGARSAMY ARUN
37. Inventory of traditional medicinal plants and their uses from Atinggola, North Gorontalo District, Gorontalo Province, Indonesia [PDF]
NOVRI YOULA KANDOWANGKO, MUKHLISULFATIH LATIEF, RAMPI YUSUF
38. Short Communication: Genetic variability of local corn cultivars from Kisar Island, Maluku, Indonesia based on morphological characters [PDF]
HERMALINA SINAY, RITHA L. KARUWAL
39. Profiles of water quality at Menjangan Besar Island, Karimunjawa, Central Java Province, Indonesia [PDF]
BAMBANG SULARDIONO, CHURUN A'IN, MAX RUDOLF MUSKANANFOLA

40. Geometric morphometrics species discrimination within the Genus *Nemipterus* from Malaysia and its surrounding seas [\[PDF\]](#)
AYESHA IMTIAZ, DARLINA MD NAIM
41. Characterization of phytochemicals and yield components in various okra (*Abelmoschus esculentus*) genotypes [\[PDF\]](#)
MUHARAMA YORA, MUHAMAD SYUKUR, SOBIR
42. Smallholder farmers livestock production on the face of climate change in Bahir Dar, Zuria District, Northwestern Ethiopia [\[PDF\]](#)
BIMREW ASMARE, FENTAHUN MEHERET
43. Lichen diversity in geothermal area of Kamojang, Bandung, West Java, Indonesia and its potential for medicines and dyes [\[PDF\]](#)
JOKO KUSMORO, IIN SUPARTINAH NOER, MUHAMAD FEISAL JATNIKA, RIRIN EKA PERMATASARI, RUHYAT PARTASASMITA
44. Short Communication: Correlation, path analysis, and heritability of phenotypic characters of bread wheat F2 populations [\[PDF\]](#)
YUDIWANTI WAHYU, NURWANITA EKASARI PUTRI, TRIKOESOEMANINGTYAS, SURJONO HADI SUTJAHJO, AMIN NUR
45. Mass vegetative propagation of rare and endangered tree species of Indonesia by shoot cuttings by KOFFCO method and effect of container type on nursery storage of rooted cuttings [\[PDF\]](#)
HENTI HENDALASTUTI RACHMAT, ATOK SUBIAKTO, ARIDA SUSILOWATI
46. Diversity of uropathogenic *Escherichia coli* lytic phage from Cisadane River, West Java, Indonesia based on morphology and protein molecular weight characteristics [\[PDF\]](#)
HAFIZAH ILMU SUFA, SRI BUDIARTI, IMAN RUSMANA
47. Entomopathogenic fungi of soils of freshwater swamps, tidal lowlands, peatlands, and highlands of South Sumatra, Indonesia [\[PDF\]](#)
AYU SAFITRI, SITI HERLINDA, ARUM SETIAWAN
48. Genetic diversity analysis of daluga (*Cyrtosperma merkusii*) using Sequence-Related Amplified Polymorphism in Siau, Sangihe and Talaud Islands, North Sulawesi, Indonesia [\[PDF\]](#)
INA ERLINAWATI, ABINAWANTO, ANDI SALAMAH, RUGAYAH
49. Short Communication: Detection of lectin gene (MLL1 and M35) in mulberry plant (*Morus* spp.) from Bogor, West Java, Indonesia [\[PDF\]](#)
YASINTA RATNA ESTI WULANDARI, YOGIARA, MICHAEL LIZAR
50. Genetic diversity and indirect selection of fine cacao (*Theobroma cacao*) based on bean color [\[PDF\]](#)
LUKITA DEVY, INDAH ANITA-SARI, AGUNG WAHYU SUSILO, ADE WACHJAR, SOBIR

Print on demand [\[PDF\]](#)

BIODIVERSITAS


Journal of Biological Diversity

[Contact Us](#)
[FAQ](#)
[Useful links](#)
[Home](#)
[Content and Archives](#)
[Aims and Scopes](#)
[Editorial Board](#)
[Guidance for Authors](#)
[Ethical Guidelines](#)
[Charges](#)
[Membership](#)
[Previously Reviewers](#)
[Conference Events](#)

Biodiversitas Vol. 21, No. 2, February 2020
ISSN: 1412-033X
E-ISSN: 2085-4722
Editor in Chief: Sutarno
First Published: January 2000

 only search Biodiversitas


NEW UPDATE


21 (1) 2020
20 (12) 2019
20 (11) 2019
20 (10) 2019
20 (9) 2019
20 (8) 2019
20 (7) 2019
20 (6) 2019

ONLINE FIRST PUBLICATION

After the publication of Biodiversitas vol. 15, no. 2, October 2014, furthermore, all manuscripts will be published online as soon as the revision is approved (no time constraints), but they will be printed in the next issue, namely Biodiversitas vol. 16, no. 1, April 2015, and so on.

Examples of how to cite:

- *Prior to the date of printing (before 1 April 2015):*
Tavankar F, Bonyad AE. 2014. Effects of timber harvest on the structural diversity and species composition in hardwood forests. Biodiversitas DOI: 10.13057/biodiv/d160101. (Year is referring to the download time)
- *After the print date (after 1 April 2015):*
Tavankar F, Bonyad AE. 2015. Effects of timber harvest on the structural diversity and species composition in hardwood forests. Biodiversitas 16 (1): 1-9.

NOTIFICATION

Starting on January 1, 2019, Biodiversitas will be issued monthly. New issues will also be found at [Smujo](#), as an online first publication.




DOAJ DIRECTORY OF OPEN ACCESS JOURNALS



[Home](#) | [Archives](#) | [Contact Us](#) | [Affiliate Program](#)



All content on this website, unless otherwise noted, is licensed under a Creative Commons Attribution 4.0 License.



ISSN: 1412-033X
E-ISSN: 2085-4722

BIODIVERSITAS

Journal of Biological Diversity

Volume 19 - Number 6 - November 2018

BIODIVERSITAS

Journal of Biological Diversity
Volume 19 - Number 6 - November 2018

ISSN/E-ISSN:

1412-033X (printed edition), 2085-4722 (electronic)

EDITORIAL BOARD (COMMUNICATING EDITORS):

Abdel Fattah N.A. Rabou (Palestine), Agnieszka B. Najda (Poland), Alan J. Lymbery (Australia), Alireza Ghanadi (Iran), Ankur Patwardhan (India), Bambang H. Saharjo (Indonesia), Daiane H. Nunes (Brazil), Darlina Md. Naim (Malaysia), Ghulam Hassan Dar (India), Faiza Abbasi (India), Hassan Pourbabaie (Iran), I Made Sudiana (Indonesia), Ivan Zambrana-Flores (United Kingdom), Joko R. Witono (Indonesia), Katsuhiko Kondo (Japan), Krishna Raj (India), Livia Wanntorp (Sweden), M. Jayakara Bhandary (India), Mahdi Reyahi-Khoram (Iran), Mahendra K. Rai (India), Mahesh K. Adhikari (Nepal), Maria Panitsa (Greece), Muhammad Akram (Pakistan), Mochamad A. Soendjoto (Indonesia), Mohib Shah (Pakistan), Mohamed M.M. Najim (Srilanka), Morteza Eighani (Iran), Pawan K. Bharti (India), Paul K. Mbugua (Kenya), Rasool B. Tareen (Pakistan), Seweta Srivastava (India), Seyed Aliakbar Hedayati (Iran), Shahabuddin (Indonesia), Shahr Shamsir (Malaysia), Shri Kant Tripathi (India), Stavros Lalas (Greece), Subhash Santra (India), Sugiyarto (Indonesia), T.N. Prakash Kammardi (India)

EDITOR-IN-CHIEF:

Sutarno

EDITORIAL MEMBERS:

English Editors: **Graham Eagleton** (grahameagleton@gmail.com), **Suranto** (surantouns@gmail.com); Technical Editor: **Solichatun** (solichatun_s@yahoo.com), **Artini Pangastuti** (pangastuti_tutut@yahoo.co.id); Distribution & Marketing: **Rita Rakhmawati** (oktia@yahoo.com); Webmaster: **Ari Pitoyo** (aripitoyo@yahoo.com)

MANAGING EDITORS:

Ahmad Dwi Setyawan (unsjournals@gmail.com)

PUBLISHER:

The Society for Indonesian Biodiversity

CO-PUBLISHER:

Department of Biology, Faculty of Mathematics and Natural Sciences, Sebelas Maret University, Surakarta

ADDRESS:

Jl. Ir. Sutami 36A Surakarta 57126. Tel. +62-271-7994097, Tel. & Fax.: +62-271-663375, Email: unsjournals@yahoo.com

ONLINE:

biodiversitas.mipa.uns.ac.id

EXPERTISE AND CORRESPONDING EMAIL OF THE COMMUNICATING EDITORS:

GENETIC DIVERSITY: Agnieszka B. Najda (agnieszka.najda@up.lublin.pl), Alan J. Lymbery (a.lymbery@murdoch.edu.au), Darlina Md. Naim (darlinamdn@usm.my), Mahendra K. Rai (pmkrai@hotmail.com).

SPECIES DIVERSITY: Joko R. Witono (jrwitono@yahoo.com), Katsuhiko Kondo (k3kondo@nodai.ac.jp), Livia Wanntorp (livia.wanntorp@nrm.se), Mahesh K. Adhikari (mkg_adh@wlink.com.np), Maria Panitsa (mpanitsa@upatras.gr), Mohib Shah (mohibshah@awkum.edu.pk), Paul K. Mbugua (paulkmbugua@gmail.com), Rasool B. Tareen (rbtareen@yahoo.com).

ECOSYSTEM DIVERSITY: Abdel Fattah N.A. Rabou (arabou@iugaza.edu), Alireza Ghanadi (aghannadi@yahoo.com), Ankur Patwardhan (ankurpatwardhan@gmail.com), Bambang H. Saharjo (bhsaharjo@gmail.com), Daiane H. Nunes (nunesdaiane@gmail.com), Faiza Abbasi (faeza.abbasi@gmail.com), Ghulam Hassan Dar (profdar99@gmail.com), Hassan Pourbabaie (hassan_pourbabaie@yahoo.com), I Made Sudiana (sudianai@yahoo.com), Ivan Zambrana-Flores (izambrana@gmail.com), Krishna Raj (krishnarajsec@yahoo.co.uk), Mahdi Reyahi-Khoram (phdmrk@gmail.com), Mochamad A. Soendjoto (masoendjoto@gmail.com), Mohamed M.M. Najim (mnajim@kln.ac.lk), Morteza Eighani (morteza_eighani@yahoo.com), Pawan K. Bharti (gurupawanbharti@rediffmail.com), Seweta Srivastava (seweta.21896@lpu.co.in), Seyed Aliakbar Hedayati (Hedayati@gau.ac.ir), Shahabuddin (shahabsaleh@gmail.com), Shahr Shamsir (shahrshamsir@gmail.com), Shri Kant Tripathi (sk_tripathi@rediffmail.com), Stavros Lalas (slalas@teilar.gr), Subhash Santra (scsantra@yahoo.com), Sugiyarto (sugiyarto_ys@yahoo.com), T.N. Prakash Kammardi (prakashtnk@yahoo.com).

ETHNOBIOLOGY: M. Jayakara Bhandary (mbjaikar@gmail.com), Muhammad Akram (makram_0451@hotmail.com).



Society for Indonesia
Biodiversity



Sebelas Maret University
Surakarta

GUIDANCE FOR AUTHORS

Aims and Scope *Biodiversitas*, *Journal of Biological Diversity* or abbreviated as *Biodiversitas* encourages submission of manuscripts dealing with all biodiversity aspects of plants, animals and microbes at the level of the gene, species, and ecosystem as well as ethnobiology.

Article types The journal seeks original full-length research papers, reviews, and short communication. Manuscript of original research should be written in no more than 8,000 words (including tables and picture), or proportional with articles in this publication number. Review articles will be accommodated, while, short communication should be written at least 2,000 words, except for pre-study.

Submission The journal only accepts online submission, through open journal system (<https://smujo.id/biodiv/about/submissions>) or email to the editors at unsjournals@gmail.com. Submitted manuscripts should be the original works of the author(s). The manuscript must be accompanied by a cover letter containing the article title, the first name and last name of all the authors, a paragraph describing the claimed novelty of the findings versus current knowledge. Submission of a manuscript implies that the submitted work has not been published before (except as part of a thesis or report, or abstract); and is not being considered for publication elsewhere. When a manuscript written by a group, all authors should read and approve the final version of the submitted manuscript and its revision; and agree the submission of manuscripts for this journal. All authors should have made substantial contributions to the concept and design of the research, acquisition of the data and its analysis; drafting of the manuscript and correcting of the revision. All authors must be responsible for the quality, accuracy, and ethics of the work.

Ethics Author(s) must obedient to the law and/or ethics in treating the object of research and pay attention to the legality of material sources and intellectual property rights.

Copyright If and when the manuscript is accepted for publication, the author(s) still hold the copyright and retain publishing rights without restrictions. Authors or others are allowed to multiply article as long as not for commercial purposes. For the new invention, authors are suggested to manage its patent before published.

Open access The journal is committed to free-open access that does not charge readers or their institutions for access. Readers are entitled to read, download, copy, distribute, print, search, or link to the full texts of articles, as long as not for commercial purposes. The license type is CC-BY-NC-SA.

Acceptance The only articles written in English (U.S. English) are accepted for publication. Manuscripts will be reviewed by editors and invited reviewers (double blind review) according to their disciplines. Authors will generally be notified of acceptance, rejection, or need for revision within 1 to 2 months of receipt. The manuscript is rejected if the content does not in line with the journal scope, does not meet the standard quality, inappropriate format, complicated grammar, dishonesty (i.e. plagiarism, duplicate publications, fabrication of data, citations manipulation, etc.), or ignoring correspondence in three months. The primary criteria for publication are scientific quality and biodiversity significance. **Uncorrected proofs** will be sent to the corresponding author by email as .doc or .docx files for checking and correcting of typographical errors. To avoid delay in publication, corrected proofs should be returned in 7 days. The accepted papers will be published online in a chronological order at any time, but printed in January, April, July and October.

A charge Starting on January 1, 2017, publishing costs waiver is granted to foreign (non-Indonesian) authors who first publish the manuscript in this journal, especially for graduate students from developing countries. However, other authors are charged USD 250 (IDR 3,500,000).

Reprints The sample journal reprint is only available by special request. Additional copies may be purchased when ordering by sending back the uncorrected proofs by email.

Manuscript preparation Manuscript is typed on A4 (210x297 mm²) paper size, in a single column, single space, 10-point (10 pt) Times New Roman font. The margin text is 3 cm from the top, 2 cm from the bottom, and 1.8 cm from the left and right. Smaller lettering size can be applied in presenting table and figure (9 pt). Word processing program or additional software can be used, however, it must be PC compatible and Microsoft Word based (.doc or .rtf; not .docx). **Scientific names** of species (incl. subspecies, variety, etc.) should be written in italic, except for italic sentence. Scientific name (genera, species, author), and cultivar or strain should be mentioned completely for the first time mentioning it in the body text, especially for taxonomic manuscripts. Name of genera can be shortened after first mentioning, except generating confusion. Name of the author can be eliminated after first mentioning. For example, *Rhizopus oryzae* L. UICC 524, hereinafter can be written as *R. oryzae* UICC 524. Using trivial name should be avoided, otherwise generating confusion. **Biochemical and chemical nomenclature** should follow the order of the IUPAC - IUB. For DNA sequence, it is better used Courier New font. Symbols of standard chemical and abbreviation of chemistry name can be applied for common and clear used, for example, completely written butilic hydroxyl toluene (BHT) to be BHT hereinafter. **Metric measurement** use IS denomination, usage other system should follow the value of equivalent with the denomination of IS first mentioning. Abbreviations set of, like g, mg, mL, etc. do not follow by dot. Minus index (m⁻², L⁻¹, h⁻¹) suggested to be used, except in things like "per-plant" or "per-plot". **Equation of mathematics** does not always can be written down in one column with text, in that case can be written separately. **Number**

one to ten are expressed with words, except if it relates to measurement, while values above them written in number, except in early sentence. The fraction should be expressed in decimal. In the text, it should be used "%" rather than "percent". Avoid expressing ideas with complicated sentence and verbiage, and used efficient and effective sentence.

Title of the article should be written in compact, clear, and informative sentence, preferably not more than 20 words. Name of author(s) should be completely written. **Name and institution** address should also be completely written with street name and number (location), postal code, telephone number, facsimile number, and email address. Manuscript written by a group, author for correspondence along with address is required. First page of the manuscript is used for writing above information.

Abstract should not be more than 200 words. **Keywords** is about five words, covering scientific and local name (if any), research theme, and special methods which used; and sorted from A to Z. All important **abbreviations** must be defined at their first mention. **Running title** is about five words. **Introduction** is about 400-600 words, covering the background and aims of the research. **Materials and Methods** should emphasize on the procedures and data analysis. **Results and Discussion** should be written as a series of connecting sentences, however, for manuscript with long discussion should be divided into subtitles. Thorough discussion represents the causal effect mainly explains for why and how the results of the research were taken place, and do not only re-express the mentioned results in the form of sentences. **Concluding** sentence should be given at the end of the discussion. **Acknowledgments** are expressed in a brief; all sources of institutional, private and corporate financial support for the work must be fully acknowledged, and any potential conflicts of interest are noted.

Figures and Tables of maximum of three pages should be clearly presented. Title of a picture is written down below the picture, while title of a table is written above the table. Colored figures can only be accepted if the information in the manuscript can lose without those images; chart is preferred to use black and white images. Author could consign any picture or photo for the front cover, although it does not print in the manuscript. All images property of others should be mentioned source. **There is no appendix**, all data or data analysis are incorporated into Results and Discussions. For broad data, it can be displayed on the website as a supplement.

References Author-year citations are required. In the text give the authors name followed by the year of publication and arrange from oldest to newest and from A to Z. In citing an article written by two authors, both of them should be mentioned, however, for three and more authors only the first author is mentioned followed by et al., for example: Saharjo and Nurhayati (2006) or (Boonkerd 2003a, b, c; Sugiyarto 2004; El-Bana and Nijs 2005; Balagadde et al. 2008; Webb et al. 2008). Extent citation as shown with word "cit" should be avoided. Reference to unpublished data and personal communication should not appear in the list but should be cited in the text only (e.g., Rifai MA 2007, pers. com. (personal communication); Setyawan AD 2007, unpublished data). In the reference list, the references should be listed in an alphabetical order (better, if only 20 for research papers). Names of journals should be abbreviated. Always use the standard abbreviation of a journal's name according to the **ISSN List of Title Word Abbreviations** (www.issn.org/2-22661-LTWA-online.php). The following examples are for guidance.

Journal:

Saharjo BH, Nurhayati AD. 2006. Domination and composition structure change at hemic peat natural regeneration following burning; a case study in Pelalawan, Riau Province. *Biodiversitas* 7: 154-158.

Book:

Rai MK, Carpinella C. 2006. *Naturally Occurring Bioactive Compounds*. Elsevier, Amsterdam.

Chapter in book:

Webb CO, Cannon CH, Davies SJ. 2008. Ecological organization, biogeography, and the phylogenetic structure of rainforest tree communities. In: Carson W, Schnitzer S (eds) *Tropical Forest Community Ecology*. Wiley-Blackwell, New York.

Abstract:

Assaeed AM. 2007. Seed production and dispersal of *Rhazya stricta*. 50th annual symposium of the International Association for Vegetation Science, Swansea, UK, 23-27 July 2007.

Proceeding:

Alikodra HS. 2000. Biodiversity for development of local autonomous government. In: Setyawan AD, Sutarno (eds.) *Toward Mount Lawu National Park; Proceeding of National Seminary and Workshop on Biodiversity Conservation to Protect and Save Germplasm in Java Island*. Universitas Sebelas Maret, Surakarta, 17-20 July 2000. [Indonesian]

Thesis, Dissertation:

Sugiyarto. 2004. *Soil Macro-invertebrates Diversity and Inter-Cropping Plants Productivity in Agroforestry System based on Sengon*. [Dissertation]. Universitas Brawijaya, Malang. [Indonesian]

Information from internet:

Balagadde FK, Song H, Ozaki J, Collins CH, Barnet M, Arnold FH, Quake SR, You L. 2008. A synthetic *Escherichia coli* predator-prey ecosystem. *Mol Syst Biol* 4: 187. www.molecularsystemsbiology.com

Entomopathogenic fungi of soils of freshwater swamps, tidal lowlands, peatlands, and highlands of South Sumatra, Indonesia

By Arum Setiawan

Entomopathogenic fungi of soils of freshwater swamps, tidal lowlands, peatlands, and highlands of South Sumatra, Indonesia

AYU SAFITRI¹, SITI HERLINDA^{2,3,*}, ARUM SETIAWAN⁴

¹Department of Environmental Science, Graduate Program, Universitas Sriwijaya, Palembang 30139, South Sumatra, Indonesia

²Department of Plant Pest and Disease, Faculty of Agriculture, Universitas Sriwijaya, Indralaya, Ogan Ilir 30662, South Sumatra, Indonesia.
Tel./fax: +62-711-580663, *email: sitherlinda@unsri.ac.id

³Research Center for Sub-optimal Lands (PUR-PLSO), Universitas Sriwijaya, Palembang 30139, South Sumatra, Indonesia

⁴Department of Biology, Faculty of Mathematic and Natural Sciences, Universitas Sriwijaya, Indralaya 30662, South Sumatra, Indonesia

Manuscript received: 11 June 2018. Revision accepted: 22 November 2018.

Abstract. Safitri A, Herlinda S, Setiawan A. 2018. Entomopathogenic fungi of soils of freshwater swamps, tidal lowlands, peatlands, and highlands of South Sumatra, Indonesia. *Biodiversitas* 19: 2365-2373. Ecosystems of lowlands and highlands in South Sumatra have specific characteristics of soils and vegetations that can affect the availability of entomopathogenic fungi. This study aimed to explore and identify species and to determine inoculum potentials of the entomopathogenic fungi from soils of freshwater swamps, tidal lowlands, peatlands, and highlands. Baiting of entomopathogenic fungi on soil samples used the larvae of *Tenebrio molitor*. The entomopathogenic fungi species found in this research were *Beauveria bassiana* and *Metarhizium anisopliae*. The number of the fungal isolates were 30 isolates consisting of nine isolates of *B. bassiana* and 21 isolates of *M. anisopliae*. The highest number of isolates was found in the highland ecosystem (11 isolates) and the lowest was found in peatland ecosystem (4 isolates). The highest percentage average of inoculum potentials of the fungi was found in the high land ecosystem (4.04%) and the lowest one was found in freshwater swamps ecosystem (2.11%). Based on the vegetation type, the soil planted with mustard in Talang Patai-Pagaralam (highland ecosystem) had the highest inoculum potentials (9.33%). These fungi will make an important contribution to the biological control for insect pests in lowland to highland ecosystems in Indonesia.

Keywords: *Beauveria bassiana*, lowlands, *Metarhizium anisopliae*, peat soils, wetlands

INTRODUCTION

Many parts of lowlands and highlands of Indonesia are used for agriculture. Lowlands consisting of freshwater swamps, tidal lowlands, and peatlands are the ecosystems having water saturated condition for the whole year (Sudana 2005). The lowland ecosystems for agricultural purposes in Indonesia are distributed in Sumatra, Kalimantan, and Papua Islands covering 11 million ha of tidal lowlands, comprising 9.2 million ha of freshwater swamps, and 14.9 million ha of peatlands, respectively (Mulyani and Sarwani 2013). While the highland ecosystems managed for agriculture with the area of 16.15 million ha are distributed all over Indonesia islands (Center for Agriculture Data and Information System, Secretariat General 2017).

Soils between lowlands and highlands in Indonesia have different characteristics, especially in the moisture, texture, and acidity (pH). The specific characteristics of lowland and highland soils at the four typologies of ecosystems consist of the freshwater swamp, tidal lowlands, peatlands, and highland closely related to specific cultivated vegetations or crops. Paddy usually is cultivated at freshwater swamps (Herlinda et al. 2018; Lakitan et al. 2018a; Lakitan et al. 2018b). There is only a small part of soils in peatlands that can be cultivated with seasonal crops and most peatlands in Indonesia are utilized for forestry and

conservation areas (Suriadikarta and Sutriadi 2007). Highlands area usually planted with various types of seasonal and annual crops. Specific vegetation or crop plants could affect the soil microorganisms associated with plant and crop roots (El-Ghany 2015).

Microorganisms such as fungi are found in freshwater swamps and highlands of South Sumatra and can be used to control insect pest called entomopathogenic fungi such as *Beauveria bassiana* and *Metarhizium anisopliae* (Herlinda et al. 2008; 2010). The entomopathogenic fungi are proven to be an effective agent to control some insect pests (Chinniah et al. 2016), and they are not harmful toward natural enemies of insect pests (Gholamzadeh-Chitgar et al. 2017). However, there is no complete information for fungi of tidal lowlands and peatlands in that region. *B. bassiana* and *M. anisopliae* were found to kill insect pests such as *Crocidolomia pavonana* (Hasyim et al. 2009), *Aphis gossypii* (Herlinda 2010), *Plutella xylostella* (Loc and Chi 2007), *Lygus* spp. (Leland et al. 2005), and *Oryctes rhinoceros* (Moslim et al. 2009). Entomopathogen, such as entomopathogenic fungi can be explored in soils (El-Ghany 2015). Certain entomopathogen species have adaptation capability in certain soils (Bugeme et al. 2008). Entomopathogen already adapted in freshwater swamp soils or highland soils generally has the specific advantage, i.e., it can adapt more effectively in this soil environment (Erler and Ates 2015). Exploration of entomopathogen starting

from lowland to highland ecosystems will produce the high variation of species and genetics that can be utilized in the extended area and specific location. The purposes of this study were to explore and identify species and to determine inoculum potentials of the entomopathogenic fungi of soils of freshwater swamps, tidal lowlands, peatlands, and highlands in South Sumatra.

MATERIALS AND METHODS

Study area

Soil sampling was carried out in December 2017. The study sites were selected in the agricultural center with specific typology ecosystems in some locations of South Sumatra (Figure 1 and Table 1). The sites consisted of freshwater swamps, tidal lowlands, peatlands, and highlands. Those ecosystems chosen for explorations of entomopathogenic fungi aimed to obtain the fungi adapting in lowland and highland soils. The selected freshwater swamp locations were in Gandus and Musi Subdistricts of Palembang City; Rambutan Subdistrict of Banyuasin District, and Pemulutan Subdistrict of Ogan Ilir District. The selected locations of tidal lowlands were Mulya Sari, Telang Sari, Muara Sungsang Subdistricts of Banyuasin District. The selected peatland locations were in Talang Dabok, Sepueuk, and Kedaton Subdistricts of Ogan Komering Ilir District. The selected highland locations were in Lematang, Tanjung Payang, and Pulau Pinang of Lahat District, and Rimau and Talang Patai of Pagaram City.

Fungal isolation, purification, and identification were conducted from January to March 2018 in the Laboratory of Entomology, Department of Plant Pest and Disease, College of Agriculture, Sriwijaya University, Indonesia. Supporting data were recorded in terms of soil sampling period, village or city names, coordinate points, and vegetation types or crop plants for each exploration locations (Figure 1 and Table 1). The pH of soil samples was measured by using a method of Kartika et al. (2018). In this survey, the application of pesticides in the sampling locations was also monitored.

Procedures

Observation of inoculum potentials of entomopathogenic fungi in the soil

Inoculum potential observation of the entomopathogenic fungi in this research was conducted by exploration of fungi in soils of the freshwater swamps, tidal lowlands, peatlands, and highlands. Entomopathogenic fungi exploration was carried out by modifying the methods of Anwar et al. (2015), i.e., using newly-moulting third instar of *Tenebrio molitor* (Yellow mealworm beetle) as insect bait (*Tenebrio* bait method) which was infested on fungal inoculum of soil samples.

The soil samples were collected from some crop fields in the sampling locations (Table 1). Soils were taken from each location at the diagonal position. Soil samples of 5000 g were obtained by digging soil at the depth of 5-15 cm in the vicinity of crop roots, and then they were put into the plastic pouch and labeled with the information of period of the sampling and crop plants.




Figure 1. Locations of exploration for entomopathogenic fungi of soils in ecosystems of South Sumatra, Indonesia: point 1 = tidal lowlands (S 02°40.866' E 104°44.298'), point 2 = freshwater swamps (S 03°02.581' E 104°51.231'), point 3= peatlands (S 03°24.840' E 104°53.362'), and point 4 = highlands (S 03°48.063' E 103°32.072' and S 03°50.174' E 103°31.293')

Table 1. Locations of exploration for entomopathogenic fungi in South Sumatra, Indonesia

Ecosystems	Village or city	Vegetation or crop plants	Height from sea level (m)
Freshwater swamps	Rambutan, Banyuasin	Paddy	15
	Gandus, Palembang	Paddy	12
	Musi Dua, Palembang	Paddy	16.67
Tidal lowlands	Pemulutan, Ogan Ilir	Paddy	22
	Mulya Sari, Banyuasin	Corn, paddy, watermelon	16.33
	Telang Sari, Banyuasin	Corn, coconut, corn + coconut	18.33
Peatlands	Muara Sungsang, Banyuasin	Coconut, banana, pineapple	15
	Talang Dabok, Ogan Komering Ilir	Palm, rubber, pineapple	24
	Sepucuk, Ogan Komering Ilir	Oil palm, rubber, pineapple	23
Highlands	Kedaton, Ogan Komering Ilir	Oil palm	19.67
	Talang Patai, Pagaram	Cabbage	170
	Pulau Pinang, Lahat	Rubber + coffee	161
	Tanjung Payang, Lahat	Rubber	121
	Lematang, Lahat	Paddy	121.3
	Rimau, Pagaram	Tea	1,326

Previously, soil samples were sieved by using 5-mesh sieve to separate them from crop roots. Then, samples were put into plastic tray (size of 32 cm x 25 cm x 5 cm) and yielded 1000 g of finer soil sample. The finer soil samples were subsequently moistened with sterile aquadest until they obtained the soil moisture of 80-90% using the method of Chen et al. (2014). The number of larvae used per sampling location was 150 larvae (5 plastic trays each containing 30 larvae) (Table 1), so the total larvae per ecosystem type was as follows: in the freshwater swamps it was 1,800 (150 larvae x 12 sampling locations), in tidal lowlands it was 1,350 (150 larvae x 9 sampling locations), in peatlands it was 1,350 (150 larvae x 9 sampling locations), and in highlands it was 2,250 (150 larvae x 15 sampling locations).

The larvae were sterilized with 70% alcohol, and put on soil-sample surface in a plastic tray. The body of larvae was sprinkled using soil sample of about 5 mm thick. Subsequently, the plastic trays containing soil samples were covered with black cloth and sprayed with sterile aquadest to maintain humidity of soil samples. The larvae were incubated within soil sample for seven days to provide enough time for entomopathogenic fungi to infect them. Then, the dead larvae infected by the entomopathogenic fungi were recorded daily to determine inolum potential. The inoculum potentials of entomopathogenic fungi in this research were measured according to the percentage of infected *Tenebrio* bait or hosts (Hofgaard et al. 2016). The inoculum potential (IP) was calculated using the equation below:

$$IP = \frac{ib}{tb} \times 100$$

ib was the number of infected *Tenebrio* bait, and *tb* was the total number of *Tenebrio* baits.

Entomopathogen isolation

The dead *Tenebrio* baits were subsequently isolated and purified by using the methods of Herlinda (2010).

Entomopathogenic fungi infecting and growing on integument of *Tenebrio* bait were isolated and grown on medium of Sabouraud Dextrose Agar (SDA, Merck). The integument surface of larvae infected by the entomopathogenic fungi was previously sterilized using the modified method of Nuraini et al. (2017) with 1% sodium hypochlorite for 3 minutes and subsequently was rinsed three times with sterile aquadest and air dried on sterile filter paper. The larvae were put into a Petri dish containing sterile humid tissue paper and then incubated in order to stimulate the growth of entomopathogenic fungi. Conidia of entomopathogenic fungi emerging from the dead larvae body were taken by using sterile ose needle and moved into a Petri dish containing SDA medium, and incubated for seven days at the constant temperature of 25 °C within the incubator.

Identification of Entomopathogen Fungi

The purified fungi were identified according to macroscopic and microscopic characteristics using the method of Guilherme et al. (2015). Fungi growing on SDA medium with the area of 1 cm² were taken by using ose needle and put into preparations containing SDA medium, incubated for three days and then microscopically observed. Furthermore, fungi were identified by using books of Humber (2005) and El-Ghany (2015).

Data analysis

Data on inoculum potentials based on percentage of *Tenebrio* bait which was infected by the entomopathogenic fungi among the treatments were analyzed descriptively.

RESULTS AND DISCUSSION

Insect characteristics infected by entomopathogenic fungi

The fungi obtained in this study were identified as *B. bassiana* and *M. anisopliae*. There were only 30 isolates found from these two fungal species (Table 2). These 30 isolates were isolated from 223 infected larvae of *Tenebrio*

bait by the entomopathogenic fungi (Table 3-7); however a lot of larvae were failed to be isolated due to contamination mostly by aerial fungi and *Trichoderma* spp. These 30 isolates consisted of nine isolates of *B. bassiana* and 21 isolates of *M. anisopliae*.

Tenebrio bait infected by the entomopathogenic fungi showed macroscopic and microscopic characteristics or symptoms that could be used to confirm the determination of the entomopathogenic fungal species. Sick or dead *Tenebrio* bait infected by *B. bassiana* showed symptoms as follows: insect body was dry and wrinkle, its outer integument was coated by white mycelia similar to silk, rigid, easily broken and no smell (Figure 2). The *Tenebrio* bait attacked by *M. anisopliae* showed symptoms as follows: dry and wrinkle, no smell, brittle and easily broken, but its outer integument was coated by mycelia having greenish white to dark green or dark color (Figure 3).

The pure isolate was obtained from dead *Tenebrio* bait body which was attacked by entomopathogenic fungi with colony characteristics for each species as follows; colony of *B. bassiana* had the white color similar to cotton, but gradually its color changed into yellowish white as fungi become older. Colony of *M. anisopliae* initially had white color similar to color of *B. bassiana*, but the color changed into greenish and dark green or dark as fungi become older (Figure 4). Conidia of *B. bassiana* and *M. anisopliae* with specific characteristics were obtained from each colony species of entomopathogenic fungi. Conidia of *B. bassiana* has a single cell and a round shape, whereas conidia of *M. anisopliae* also has a single cell but with a cylindrical shape (Figure 5). Mycelia of *B. bassiana* and *M. anisopliae* insulate with upright, branches, and layers of conidiophores.

Inoculum potentials of entomopathogenic fungi in the soil of South Sumatra

Inoculum potentials of entomopathogenic fungi in this research were measured according to the percentage of infected *Tenebrio* bait. The inoculum potentials of entomopathogenic fungi derived from the freshwater swamp soil, tidal lowlands, peat soils, and high land each was different among the survey locations (Tables 3-6). The value of inoculum potential in soil of freshwater swamps, tidal lowlands, peatlands and high lands were in the range of 0.67-3.33%, 2.67-7.33%, 0-4%, and 1.33-9.33%, respectively.

From the freshwater swamp soils, we only could obtain five isolates of the entomopathogenic fungi isolated from 35 infected *tenebrio* baits (Table 3). A lot of the infected *Tenebrio* baits were failed to be isolated due to the contamination by other fungi. The inoculum potentials of the entomopathogenic fungi from the tidal lowlands gained were only 10 isolates (Table 4). The peat soils in sepucuk, Ogan Komering Ilir were planted with rubber had higher inoculum potentials (Table 5). The highland soils in Talang Patai, Pagaralam were planted with mustard also had higher inoculum potential (Table 6). The highest percentage of inoculum potentials of the fungi was found in the highland ecosystem and the lowest one was found in

the freshwater swamp ecosystem. However, the highest number of isolates was found in highland ecosystem (11 isolates) and the lowest one found in peatland ecosystem (4 isolates) (Table 7).



Figure 2. *Tenebrio* bait infected by *Beauveria bassiana* (A) and healthy *Tenebrio* (B)



Figure 3. *Tenebrio* bait infected by *Metarhizium anisopliae* (A) and healthy *Tenebrio* (B)




Figure 4. Colony of *Beauveria bassiana* (A) and *Metarhizium anisopliae* (B)




Figure 5. Conidia of *Beauveria bassiana* (A) and *Metarhizium anisopliae* (B) (400x magnification)

Table 2. Species and isolates of entomopathogenic fungi found in South Sumatra, Indonesia

Ecosystems	Species of fungi	Number of isolate	Vegetation or crop plants	Village or city
Freshwater swamps	<i>M. anisopliae</i>	3	Paddy	Rambutan
Freshwater swamps	<i>M. anisopliae</i>	1	Paddy	Pemulutan
Freshwater swamps	<i>B. bassiana</i>	1	Paddy	Rambutan
Tidal lowlands	<i>M. anisopliae</i>	1	Corn + coconut	Telang Sari
Tidal lowlands	<i>M. anisopliae</i>	2	Corn	Telang Sari
Tidal lowlands	<i>M. anisopliae</i>	2	Corn	Mulya Sari
Tidal lowlands	<i>M. anisopliae</i>	3	Paddy	Mulya Sari
Tidal lowlands	<i>B. bassiana</i>	1	Corn	Telang Sari
Tidal lowlands	<i>B. bassiana</i>	1	Watermelon	Mulya Sari
Peatlands	<i>B. bassiana</i>	4	Oil palm	Talang Dabok
Highlands	<i>M. anisopliae</i>	1	Rubber + coffee	Pulau Pinang
Highlands	<i>M. anisopliae</i>	4	Cabbage	Talang Patai
Highlands	<i>M. anisopliae</i>	4	Mustard	Talang Patai
Highlands	<i>B. bassiana</i>	1	Rubber + coffee	Pulau Pinang
Highlands	<i>B. bassiana</i>	1	Cabbage	Talang Patai

Table 3. Inoculum potentials of entomopathogenic fungi in the freshwater swamp soils of South Sumatra, Indonesia

Village or city	Vegetation or crop species	GPS (coordinat)	Number of isolate	Inoculum potentials (%)		
				<i>B. bassiana</i>	<i>M. anisopliae</i>	Fungi (total)
Rambutan, Banyuasin	Paddy	S 03°02.581', E 104°51.231'	0	0.00 (0)	0.67 (1)	0.67 (1)
Rambutan, Banyuasin	Paddy	S 03°02.591', E 104°51.217'	2	0.67 (1)	2.00 (3)	2.67 (4)
Rambutan, Banyuasin	Paddy	S 03°02.586', E 104°51.201'	2	0.67 (1)	0.67 (1)	1.33 (2)
Gandus, Palembang	Paddy	S 03°00.401', E 104°42.380'	0	1.33 (2)	2.00 (3)	3.33 (5)
Gandus, Palembang	Paddy	S 03°00.632', E 104°42.532'	0	0.67 (1)	0.67 (1)	1.33 (2)
Gandus, Palembang	Paddy	S 03°00.632', E 104°42.801'	0	1.33 (2)	1.33 (2)	2.67 (4)
Musi Dua, Palembang	Paddy	S 03°02.120', E 104°43.021'	0	0.67 (1)	1.33 (2)	2.00 (3)
Musi Dua, Palembang	Paddy	S 03°02.150', E 104°43.612'	0	1.33 (2)	2.00 (3)	3.33 (5)
Musi Dua, Palembang	Paddy	S 03°02.510', E 104°43.120'	0	1.33 (2)	1.33 (2)	2.67 (4)
Pemulutan, Ogan Ilir	Paddy	S 03°03.148', E 104°46.230'	0	0.67 (1)	0.67 (1)	1.33 (2)
Pemulutan, Ogan Ilir	Paddy	S 03°03.115', E 104°46.218'	0	0.67 (1)	0.67 (1)	1.33 (2)
Pemulutan, Ogan Ilir	Paddy	S 03°03.113', E 104°46.201'	1	1.33 (2)	1.33 (2)	2.67 (4)

Table 4. Inoculum potentials of entomopathogenic fungi in the tidal lowland soils of South Sumatra, Indonesia

Village or city	Vegetation or crop species	GPS (coordinat)	Number of isolate	Inoculum potentials (%)		
				<i>B. bassiana</i>	<i>M. anisopliae</i>	Fungi (total)
Mulya Sari, Banyuasin	Corn	S 02°40.866', E 104°44.298'	2	1.33 (2)	2.67 (4)	4.00 (6)
Mulya Sari, Banyuasin	Paddy	S 02°40.944', E 104°44.621'	3	1.33 (2)	2.00 (3)	3.33 (5)
Mulya Sari, Banyuasin	Watermelon	S 02°40.896', E 104°44.676'	1	1.33 (2)	2.00 (3)	3.33 (5)
Telang Sari, Banyuasin	Corn	S 02°38.842', E 104°45.369'	2	1.33 (2)	2.67 (4)	4.00 (6)
Telang Sari, Banyuasin	Coconut	S 02°38.813', E 104°45.801'	1	0.67 (1)	1.33 (2)	2.00 (3)
Telang Sari, Banyuasin	Corn + coconut	S 02°38.875', E 104°44.495'	1	2.00 (3)	5.33 (8)	7.33 (11)
Muara Sungsang, Banyuasin	Coconut	S 02°21.736', E 104°50.635'	0	1.33 (2)	4.00 (6)	5.33 (8)
Muara Sungsang, Banyuasin	Banana	S 02°21.823', E 104°50.632'	0	0.67 (1)	2.00 (3)	2.67 (4)
Muara Sungsang	Pineapple	S 02°22.542', E 104°50.324'	0	0.67 (1)	2.00 (3)	2.67 (4)

Table 5. Inoculum potentials of entomopathogenic fungi in the peatland soils of South Sumatra, Indonesia

Village or city	Vegetation or crop plants	GPS (coordinat)	Inoculum potentials (%)			
			Number of isolate	<i>B. bassiana</i>	<i>M. anisoplia</i>	Fungi (Total)
Talang Dabok, Ogan Komering Ilir	Oil palm	S 03°23.570', E 104°51.498'	4	3.33 (5)	1.33 (2)	4.67 (7)
Talang Dabok, Ogan Komering Ilir	Rubber	S 03°25.673', E 104°53.258'	0	2.67 (4)	1.33 (2)	4.00 (6)
Talang Dabok, Ogan Komering Ilir	Pineapple	S 03°25.280', E 104°52.940'	0	2.00 (3)	0.67 (1)	2.67 (4)
Sepucuk, Ogan Komering Ilir	Oil palm	S 03°24.840', E 104°53.362'	0	2.00 (3)	1.33 (2)	3.33 (5)
Sepucuk, Ogan Komering Ilir	Rubber	S 03°23.715', E 104°52.275'	0	3.33 (5)	2.00 (3)	5.33 (8)
Sepucuk, Ogan Komering Ilir	Pineapple	S 03°23.535', E 104°51.780'	0	1.33 (2)	1.33 (2)	2.67 (4)
Kedaton, Ogan Komering Ilir	Oil palm	S 03°23.308', E 104°51.487'	0	1.33 (2)	2.00 (3)	3.33 (5)
Kedaton, Ogan Komering Ilir	Oil palm	S 03°23.277', E 104°51.398'	0	1.33 (2)	0.67 (1)	2.00 (3)
Kedaton, Ogan Komering Ilir	Oil palm	S 03°23.204', E 104°51.459'	0	1.33 (2)	0.67 (1)	2.00 (3)

Table 6. Inoculum potentials of entomopathogenic fungi in the highland soils of South Sumatra, Indonesia

Village or city	Vegetation or crop plants	GPS (coordinat)	Inoculum potentials (%)			
			Number of isolate	<i>B. bassiana</i>	<i>M. anisoplia</i>	Fungi (total)
Pulau Pinang, Lahat	Rubber + coffee (A)	S 03°48.819', E 103°32.677'	2	1.33 (2)	1.33 (2)	2.67 (4)
Pulau Pinang, Lahat	Rubber + coffee (B)	S 03°48.852', E 103°32.698'	0	1.33 (2)	2.00 (3)	3.33 (5)
Pulau Pinang, Lahat	Rubber + coffee (C)	S 03°48.883', E 103°32.688'	0	0.67 (1)	0.67 (1)	1.33 (2)
Tanjung Payang, Lahat	Rubber (A)	S 03°48.094', E 103°32.145'	0	0.67 (1)	1.33 (2)	2.00 (3)
Tanjung Payang, Lahat	Rubber (B)	S 03°48.122', E 103°32.162'	0	3.33 (5)	2.67 (4)	6.00 (9)
Tanjung Payang, Lahat	Rubber (C)	S 03°48.177', E 103°32.166'	0	2.00 (3)	2.67 (4)	4.67 (7)
Lematang, Lahat	Paddy	S 03°48.063', E 103°32.072'	0	0.67 (1)	1.33 (2)	2.00 (3)
Lematang, Lahat	Paddy	S 03°48.051', E 103°32.069'	0	3.33 (5)	4.00 (6)	7.33 (11)
Lematang, Lahat	Paddy	S 03°48.020', E 103°32.064'	0	1.33 (2)	1.33 (2)	2.67 (4)
Talang Patai, Pagaramalam	Cabbage	S 03°50.180', E 103°31.325'	2	4.00 (6)	4.67 (7)	8.67 (13)
Talang Patai, Pagaramalam	Cabbage	S 03°50.174', E 103°31.313'	3	0.67 (1)	1.33 (2)	2.00 (3)
Talang Patai, Pagaramalam	Mustard	S 03°50.174', E 103°31.293'	4	4.67 (7)	4.67 (7)	9.33 (14)
Rimau, Pagaramalam	Tea	S 04°02.161', E 103°10.484'	0	2.00 (3)	2.00 (3)	4.00 (6)
Rimau, Pagaramalam	Tea	S 04°02.144', E 103°10.487'	0	1.33(2)	1.33(2)	2.67 (4)
Rimau, Pagaramalam	Tea	S 04°02.136', E 103°10.485'	0	0.67 (1)	1.33(2)	2.00 (4)

Table 7. Inoculum potentials of entomopathogenic fungi in the soil of South Sumatra, Indonesia

Ecosystems	Inoculum potentials (%)			
	Number of isolate	<i>B. bassiana</i>	<i>M. anisoplia</i>	Fungi (total)
Freshwater Swamps	5	0.89 (16)	1.22 (22)	2.11 (38)
Tidal Lowlands	10	1.18 (16)	2.67 (36)	3.85 (52)
Peatlands	4	2.07 (28)	1.25 (17)	3.33 (45)
Highlands	11	1.87 (42)	2.17 (49)	4.04 (91)

Discussion

Entomopathogenic species found in this research were *B. bassiana* and *M. anisopliae*. Macroscopic and microscopic characteristics of both entomopathogenic fungi found in this research matched with the previous studies. Humber (2005) stated that mycelia of *B. bassiana* appeared from exoskeleton of hosts insect, and covered all parts of the exterior surface of host integument, so that host body has white color, reverse pale to yellow colony, hyaline or colorless, single cell, and globular and conidia as well as insulate hypha. *M. anisopliae* causes integument

which has colors of whitish green to dark green because its mycelia cover exoskeleton of hosts insect; it has green to yellow conidia, single cell and cylindrical conidia as well as insulate hypha (Driver et al. 2000; Humber 2005).

Beauveria bassiana and *M. anisopliae* have parasitic and saprophytic phases during the killing process of their host insect (Augustyniuk-Kram and Kram 2012; El-Ghany 2015). Parasitic phase starts viz, the fungal conidia attach to the host insect cuticle, and then the conidia germinate on the host cuticle (El-Ghany 2015). The fungal penetration into the insect cuticle can be performed in producing

specific infection hyphae originating at appressoria of the fungus (Fernandes et al. 2007; El-Ghany 2015). Gürlek et al. (2018) reported that both species could produce germ tubes growing over the surface of the insect cuticle until the tubes contacted weakness area of cuticle where penetration could easily be achieved. After the fungus successfully penetrated, then mycelia distributed into the hemolymph by the formation of blastospores (El-Ghany 2015). Finally, the host insect would die within four days of penetration (Gürlek et al. 2018). Saprophytic phase starts viz, the fungus grew mechanically in the dead insect body, retrieved nutrients from the insect body, and then the fungus produced toxins (El-Ghany 2015).

The success of both fungi in conducting the process of parasitic and saprophytic phases was affected by several external factors such as moisture, pH, temperature, ultraviolet (UV) radiation, and vegetation (El-Ghany 2015). This research showed that highland soil planted with cabbage and mustard in Talang Patai, Pagaralam had more inoculum potentials of entomopathogenic fungi than other locations because cabbage and mustard were inhabited by insect pests dominated by Lepidoptera, such as *Phutella xylostella*, *Crocidolomia binotalis*, *Spodoptera litura*, and *Chrysodeixis chalcites*; while the larvae of Lepidoptera were the most suitable hosts for entomopathogenic fungi (Godonou et al. 2009; Nunilahwati et al. 2012). This study also found that a lot of infected larvae of the insect pests hung above of the mustard and cabbage canopy. The host insects attacked by entomopathogenic fungi on the cabbage canopy at highlands, South Sumatra were generally dominated by *P. xylostella*, *S. litura*, and *C. chalcites*. The symptoms of sick or infected insect larvae were dry and in the stiff condition, white or greenish white in color and attached on the upper surface of cabbage leaves. The infected larvae which showed the symptoms of dry and stiff condition as mummy and covered with white fungal mycelia was larvae attacked by *B. bassiana*, whereas larvae body covered by fungal mycelia having greenish white or dark green color was symptoms of larvae attacked by *M. anisopliae*. Symptoms of larvae attacked by *B. bassiana* and *M. anisopliae* in this research matched to the symptoms reported by El-Ghany (2015) and Mora et al. (2017).

In this research, more inoculum potentials of the entomopathogenic fungi were found in the highlands and tidal lowlands than that in freshwater swamps and peatlands because it was affected by soil pH and moisture. Zhong et al. (2010) reported that soil pH had a more significant role in determining the existence of fungal propagules within soils than that of soil texture and organic matter. However, Inubushi et al. (2003) had stated that soil moisture is one of the most important controlling factors for biological reactions in the soil. Soil pH in this research was in the range of 4 to 4.5 in freshwater swamps, 4.3 to 5 in tidal lowlands, 3.60 to 4 in peatlands, and 5 to 6.7 in highlands. Kodir and Juwita (2016) stated that the pH value of soil in freshwater swamps in Indonesia are in the ranges of 4 to 4.5, pH 4.17 to 5.35 in tidal lowlands (Marlina et al. 2016), and 3.60 to 3.95 in peatlands (Utami et al. 2009), and 5 to 6 in highlands (Supriadi et al. 2016). The variation of

pH values of soils at each location could affect the adaptation capability of entomopathogenic fungi surviving (Bugeme et al. 2008). The conidial viability of entomopathogenic fungi, such as *B. bassiana* and *M. anisopliae* were affected by pH of in-vitro medium for entomopathogenic fungi. Rizkie et al. (2017) reported that high acidity (pH < 4) of in-vitro medium for fungus growing significantly decrease the conidial viability of *B. bassiana* and *M. anisopliae*. Therefore, inoculum potentials of the entomopathogenic fungi from peatland soil and freshwater swamp soil were lower than that from the soil in the tidal lowlands and highlands.

In addition to soil pH and moisture, soil texture also determines the existence and distribution of fungal propagules. Soil texture has low clay content, and sandy soil texture tends to have the low capability in maintaining the existence of fungal propagules (El-Ghany 2015) as well as the water saturated soil (Garrido-Jurado et al. 2011). Soils from freshwater swamps and peatlands in South Sumatra had lower clay content and in water saturated condition (Marlina et al. 2016; Kartika et al. 2018). The low existence of fungal propagules finding at freshwater swamp and peatland soils was due to both factors. Soils in freshwater swamps are water saturated for more than 6 to 7 months per year usually occurred from November to April (Herlinda et al. 2018) and soil moisture in the peatlands reach 500% (Maftu'ah and Susanti 2009).

Peatlands have soil pore saturated with water all year long resulting in the anaerobic condition of the soil. Peat soil has no clay, sand, and silt content, but it has organic matter (Sudana 2005). The high organic matter finally can decrease the pH of the soil (Utami et al. 2009). Rizkie et al. (2017) confirmed that pH < 4 within medium in-vitro for growing fungi could decrease the ability to live of fungal propagules. In highland ecosystems, the portion of the soil texture among clay, sand, and silt fractions was found in the same composition or balance (Utomo et al. 2013; Supriadi et al. 2016). While tidal lowlands contain silt from sedimentation mixture of river water and seawater with balance clay and sand fractions (Marlina et al. 2016). Higher inoculum potentials in highlands and tidal lowlands in this research was due to balance or higher of clay texture and organic soils than that of freshwater swamps and peatlands.

The balance or higher of clay texture and organic soils are capable of maintaining the existence of fungal propagules (Garrido-Jurado et al. 2011). Zhong et al. (2010) stated that soil in ecosystems which apply composted manure or no synthetic fertilizer had higher propagules content of *B. bassiana* than that of soil in the ecosystem which applies synthetic fertilizers. Also, the application of synthetic pesticides is capable of decreasing the existence of entomopathogenic fungi within the soil (Mietkiewski et al. 2010). Local farmers in freshwater swamp and peatland areas of South Sumatra usually do not apply synthetic pesticides, whereas many local farmers in tidal lowland and highland areas apply synthetic pesticides (Herlinda et al. 2018). Although no synthetic pesticides were applied in the freshwater swamp and peatland areas, fungal propagules or inoculum potentials in these areas were lower than those of tidal lowlands and highlands

areas. In this study, there was no evidence that having no synthetic pesticide application in freshwater swamp and peatland areas can cause the high existence of fungal propagules. However, soil pH and soil texture have more effect on the existence of propagules of *B. bassiana* and *M. anisopliae* in lowland swamp and peatland ecosystems.

Higher inoculum potentials in highlands and tidal lowlands in this research were closely related to specific cultivated vegetations or crops. Most peatlands in Indonesia were utilized for forestry and conservation areas (Suriadikarta and Sutriadi 2007). Paddy generally is cultivated at freshwater swamps (Herlinda et al. 2018; Lakitan et al. 2018a; Lakitan et al. 2018b) as well as in tidal lowlands; however, paddy cultivation at tidal lowlands was more intensive (with two to three planting indices) than that of freshwater swamps (one planting index). Thus, diverse species of vegetation or crop plants could affect abundant and diverse species of the soil microorganisms associated with plant and crop roots (El-Ghany 2015).

This research found two species of entomopathogenic fungi from soils in South Sumatra i.e., *B. bassiana* and *M. anisopliae*. The highest percentage of the inoculum potentials and prevalence of both fungi occurred in the highland ecosystems and the lowest percentage of the inoculum potentials of the fungi was found in the peatland ecosystems. In highland ecosystems, the percentage of the inoculum potentials was affected by the locations and the vegetation or the crop plants. These fungi will make an important contribution to the biological control for insect pests from lowland to highland ecosystems in Indonesia.

ACKNOWLEDGEMENTS

This research was funded by National Strategic Research (PSN) with budget year of 2018 according to the Director of Research and Community Service, Directorate of Research and Community Service (DRPM), Directorate General for Research and Development, Ministry of Research, Technology, and Higher Education, Number: 093/SP2H/LT/DRPM/IV/2018 chaired by Siti Herlinda. Special thanks to Dr. Suwandi, a microbiologist from Universitas Sriwijaya, Palembang, Indonesia for identification of fungi.

REFERENCES

- Anwar W, Khan SN, Aslam M, Haider MS, Shahid AA, Ali M. 2015. Exploring fungal flora associated with insects of cotton agroecological zones of Punjab, Pakistan. *Pakistan Entomologist* 37: 27-31.
- Augustyniuk-Kram A, Kram KJ. 2012. Entomopathogenic fungi as an important natural regulator of insect outbreaks in forests review. *Forest Ecosystem, More than Just Trees*. Blanco JA. (ed.). In Tech: Croatia
- Bugeme DM, Maniania NK, Knapp M, Boga HI. 2008. Effect of temperature on virulence of *Beauveria bassiana* and *Metarhizium anisopliae* isolates to *Tetramesa evansi*. *Exp Appl Acarol* 46: 275-285.
- Center for Agriculture Data and Information System, Secretariat General. 2017. Statistics of Agricultural Land 2012-2016. In: Hakim M, Wiratno O, and Abdurachman A (eds). Ministry of Agriculture. Jakarta.
- Chen Z, Xu L, Yang F, Ji G, Yang J, Wang J. 2014. Efficacy of *Metarhizium anisopliae* isolate MAX-2 from Shangri-la, China under desiccation stress. *BMC Microbiol* 14: 1-8.
- Driver F, Milner RJ, Trueman JWH. 2000. A taxonomic revision of *Metarhizium* based on a phylogenetic analysis of rDNA sequence data. *Mycol Res* 104: 134-150.
- El-Ghany TMA. 2015. Entomopathogenic Fungi and their Role in Biological Control. Biology Department Faculty of Science Jazan University KSA. Cairo.
- Erler F, Ates AO. 2015. Potential of two entomopathogenic fungi, *Beauveria bassiana* and *Metarhizium anisopliae* (Coleoptera: Scarabaeidae), as biological control agents against the June beetle. *J Insect Sci* 15: 1-6.
- Fernandes EKK, Rangel DEN, Moraes AM., Bittencourt VREP, Roberts DW. 2007. Variability in tolerance to UV-B radiation among *Beauveria* spp. isolates. *J Invertebr Pathol* 96: 237-243.
- Garrido-Jurado J, Torrent J, Barrón V, Corpas A, Quesada-Moraga E. 2011. Soil properties affect the availability, movement, and virulence of entomopathogenic fungi conidia against puparia of *Ceratitis capitata* (Diptera: Tephritidae). *Biol Control* 58: 277-285.
- Godonou I, James B, Atcha-Ahowé B, Vodouhe S, Kooyman C, Ahanchédé A, Kone S. 2009. Potential of *Beauveria bassiana* and *Metarhizium anisopliae* isolates from Benin to control *Plutella xylostella* L. (Lepidoptera: Plutellidae). *Crop Protection* 28: 220-224.
- Guilherme D, Oliveira P, Pauli G, Moura G, Delalibera I. 2015. A protocol for determination of conidial viability of the fungal entomopathogens *Beauveria bassiana* and *Metarhizium anisopliae* from commercial products. *J Microbiol Methods* 119: 44-52.
- Gürlek S, Sevim A, Sezgin FM, Sevim E. 2018. Isolation and characterization of *Beauveria* and *Metarhizium* spp. from walnut fields and their pathogenicity against the codling moth, *Cydia pomonella* (L.) (Lepidoptera: Tortricidae). *Egyptian J Biol Pest Control* 28: 1-6.
- Hasyim A, Nuraida, Trizelia. 2009. Pathogenicity of entomopathogenic fungi isolates to eggs and larvae of cabbage head caterpillar, *Crocidolomia pavonana* Fabricius. *J Hort* 19: 334-343.
- Herlinda S. 2010. Spore density and viability of entomopathogenic fungal isolates from Indonesia, and their virulence against *Aphis gossypii* Glover (Homoptera: Aphididae). *Tropical Life Sci Res* 21: 137-142.
- Herlinda S, Irsan C, Mayasari R, Septariani S. 2010. Identification and selection of entomopathogenic fungi as biocontrol agents for *Aphis gossypii* from South Sumatra. *Microbiol Indonesia* 4: 137-142.
- Herlinda S, Mulyati SI, Suwandi. 2008. Selection of isolates of entomopathogenic fungi and the bioefficacy of their liquid production against *Leptocorisa oratorius* nymphs. *Microbiol Indonesia* 2: 141-146.
- Herlinda S, Yudha S, Thalib R, Khodijah, Suwandi, Lakitan B, Verawaty M. 2018. Species richness and abundance of spiders inhabiting rice in freshwater swamps and tidal lowlands in South Sumatra, Indonesia. *J ISSAAS* 24: 82-93.
- Hofgaard IS, Seehusen T, Aamot HU, Riley H, Razzaghian J, Le VH, Hjelkrem A-GR, Dill-Macky R, Brodal G. 2016. Inoculum potential of *Fusarium* spp. relates to tillage and straw management in Norwegian fields of spring oats. *Front Microbiol* 7: 1-15.
- Humber RA. 2005. Entomopathogenic Fungal Identification. USDA-ARS Plant Protection Research Unit. Ithaca.
- Inubushi K, Furukawa Y, Hadi A, Purnomo E, Tsuruta H. 2003. Seasonal changes of CO₂, CH₄, and N₂O fluxes in relation to land-use change in tropical peatlands located in coastal area of South Kalimantan. *Chemosphere* 52: 603-608.
- Kartika K, Lakitan B, Wijaya A, Kadir S, Widuri LI, Siaga E. 2018. Effects of particle size and application rate of rice-husk biochar on chemical properties of tropical wetland soil, rice growth and yield. *AJCS* 12: 817-826.
- Kodir KA, Juwita Y. 2016. Morphological characterization and inventory of local wetland rice collected from South Sumatra Province. *Bul Plasma Nutraf* 22: 101-108.
- Lakitan B, Alberto A, Lindiana L, Kartika K, Herlinda S, Kurnianingsih A. 2018. The Benefits of biochar on rice growth and yield in tropical riparian wetland, South Sumatra, Indonesia. *CMUJ Natural Sciences* 17: 111-126.
- Lakitan B, Hadi B, Herlinda S, Siaga E, Widuri LI, Kartika K, Lindiana L, Yunindyawati Y, Meihana M. 2018. Recognizing farmers practices and constraints for intensifying rice production at riparian wetlands in

- Indonesia. NJAS Wageningen J Life Sci 85: 10-20.
- Leland JE, McGuire MR, Grace JA, Jaronski ST, Ulloa M, Park Y, Plattner RD. 2005. Strain selection of a fungal entomopathogen, *Beauveria bassiana*, for control of plant bugs (*Lygus* spp.) (Heteroptera: Miridae). Biol Control 35: 104-114.
- Loc NT, Chi VTB. 2007. Biocontrol potential of *Metarhizium anisopliae* and *Beauveria bassiana* against diamond back moth, *Plutella xylostella*. Omonrice 93: 86-93.
- Maftu'ah E, Susanti MA. 2009. Earthworms community on several land uses of peat land in Central Kalimantan. Berita Biologi 9:371-378.
- Marlina N, Asmawati, Zairani FY, Midranisiah, Aryani I, Kalasari R. 2016. Biofertilizer utilization in increasing inorganic fertilizer efficiency and rice yield at c-type flooding land of Tanjung Lago tidal lowland. Int J Engg res & Sci & Tech 5: 74-83.
- Mietkiewski RT, Pell JK, Clark SJ. 2010. Influence of pesticide use on the natural occurrence of entomopathogenic fungi in arable soils in the UK: Field and laboratory comparisons. Biocontrol Science and Technology 7: 565-575.
- Mora MAE, Castilho AMC, Fraga ME. 2017. Classification and infection mechanism of entomopathogenic fungi. Arq Inst Biol 84: 1-10.
- Moslim R, Kamarudin N, Wahid MB. 2009. Pathogenicity of granule formulations of *Metarhizium anisopliae* against the larvae of the oil palm rhinoceros beetle, *Oryctes rhinoceros*. J Oil Palm Res 21: 602-612.
- Mulyani A, Sarwani M. 2013. The characteristic and potential of sub-optimal land for agricultural development in Indonesia. Jurnal Sumberdaya Lokal 7: 47-55. [Indonesian]
- Nunilahwati H, Herlinda S, Irsan C, Pujiastuti Y. 2012. Exploration, isolation and selection entomopathogenic fungi infectious to *Plutella xylostella* (Lepidoptera: Yponomeutidae) on green mustard (*Brassica chinensis*) crop in South Sumatra. J HPT Tropika 12: 1-11. [Indonesian]
- Nuraini FR, Setyaningsih R, Susilowati ARI. 2017. Screening and characterization of endophytic fungi as antagonistic agents toward *Fusarium oxysporum* on eggplant (*Solanum melongena*). Biodiversitas 18: 1377-1384.
- Rizkie L, Herlinda S, Suwandi, Irsan C, Susilawati, Lakitan B. 2017. Conidial density and viability of *Beauveria bassiana* and *Metarhizium anisopliae* grown on the low-pH in vitro medium. J HPT Tropika 17: 119-127. [Indonesian]
- Sudana W. 2005. The potential and prospect of swamps as a source of agricultural production. Analisis Kebijakan Pertanian 3: 141-151. [Indonesian]
- Supriadi H, Randriani E, Towaha J. 2016. Correlation between altitude, soil chemical properties, and physical quality of arabica coffee beans in highland areas of Garut J TIDP 3: 45-52.
- Suriadikarta DA, Sutriadi MT. 2007. Potential tidal land for agricultural development. The Jurnal Litbang Pertanian 26: 115-122. [Indonesian]
- Utami SNH, Maas A, Radjaguguk B, Purwanto BH. 2009. Nature on soil physical, chemical and fir spectrophotometry of hydrophobic peat from Central Kalimantan. J Trop Soils 14: 159-166.
- Utome M, Banuwa IS, Buchari H, Anggraini Y. 2013. Long-term tillage and nitrogen fertilization effects on soil properties and crop yields. J Trop Soils 18: 131-139.
- Zhong W, Gu T, Wang W. 2010. The effects of mineral fertilizer and organic manure on soil microbial community and diversity. Plant Soil 326: 511-522.

Entomopathogenic fungi of soils of freshwater swamps, tidal lowlands, peatlands, and highlands of South Sumatra, Indonesia

ORIGINALITY REPORT

7%

SIMILARITY INDEX

MATCH ALL SOURCES (ONLY SELECTED SOURCE PRINTED)

★smujo.id
Internet

7%

EXCLUDE QUOTES ON
EXCLUDE BIBLIOGRAPHY ON

EXCLUDE MATCHES < 1%

FORMAT PENILAIAN (VALIDASI & PEER REVIEW)
LEMBAR
HASIL PENILAIAN SEJAWAT SEBIDANG ATAU PEER REVIEW
KARYA ILMIAH : JURNAL ILMIAH

Jurnal Artikel Ilmiah : Entomopathogenic fungi of soils of freshwater swamps, tidal lowlands, peatlands, and highlands of south sumatra, Indonesia

Penulis Artikel Ilmiah : Arum Setiawan

Identitas Jurnal Artikel Ilmiah : a. Nama Jurnal : Biodiversitas
 b. Nomor/Volume/Hal : 6/19/2365-2373
 c. Edisi (bulan/tahun) : November/2018
 d. Penerbit : Jurusan Biologi FMIPA UNS
 e. Jumlah Halaman : 10

Kategori Publikasi Jurnal Ilmiah : Jurnal Ilmiah Internasional Bereputasi
 (beri \surd pada kategori yang tepat) Jurnal Ilmiah Internasional
 Jurnal Ilmiah Nasional Terakreditasi S1, S2
 Jurnal Ilmiah Nasional Terakreditasi S3, S4
 Jurnal Ilmiah Nasional Tidak Terakreditasi

I. Hasil Penilaian Validasi :

No.	ASPEK	URAIAN/KOMENTAR PENILAIAN
1.	Indikasi Plagiasi	7 %
2.	Linearitas	Isi paper kurang linier dengan bidang biologi konservasi

II. Hasil Penilaian Peer Review :

Komponen Yang Dinilai	Nilai Maksimal Jurnal Ilmiah (isikan di kolom yang sesuai)					Nilai Akhir Yang Diperoleh
	Internasional Bereputasi (Maks 40)	Internasional (Maks 20)	Nasional Terakreditasi S1, S2 Maks 25	Nasional Terakreditasi S3, S4 Maks 20	Nasional tidak Terakreditasi (maks 10)	
Kelengkapan dan Kesesuaian unsur isi jurnal (10%)	4					4
Ruang lingkup dan kedalaman pembahasan (30%)	12					12
Kecukupan dan Kemutakhiran data/informasi dan metodologi (30%)	12					12
Kelengkapan unsur dan kualitas penerbit (30%)	12					12
Total = (100%)	40					40
Kontribusi Pengusul (Penulis Pertama /Anggota Utama)	Anggota Utama $(0,4 \times 40) / 2 = 8$					8
KOMENTAR/ULASAN PEER REVIEW						
• Kelengkapan dan Kesesuaian Unsur:	Paper terkait Entomopathogenic fungi di Sumatera Selatan. Isi paper sudah memenuhi kaidah-kaidah karya ilmiah dan kurang sesuai dengan bidang biologi konservasi					
• Ruang Lingkup dan Kedalaman Pembahasan:	Hasil penelitian dibahas secara komprehensif dengan penyampaian perbandingan dari temuan-temuan penelitian lainnya dan teori terkait. Referensi yang diacu dalam pembahasan sudah cukup update untuk bidang kajian ini.					
• Kecukupan & Kemutakhiran Data & Metodologi:	Data-data hasil penelitian sudah baik dan didukung peta lokasi sampling dan gambar yang ditampilkan menarik. Data didapatkan dengan menggunakan metode yang standard tidak terlalu mutakhir.					
• Kelengkapan Unsur & Kualitas Penerbit:	Penerbit Jurusan Biologi FMIPA UNS berkualitas baik, tidak termasuk predatory publisher, dan jurnal masuk di Q3.					

Surabaya, 15 Mei 2020

Penilai 1

A handwritten signature in black ink, consisting of several loops and a long horizontal stroke at the bottom.

Prof. Hery Purnobasuki, M.Si., Ph.D.

NIP 196705071991021001

Unit Kerja : Jurusan Biologi FST Unair

Bidang Ilmu : Biologi

Jabatan/Pangkat : Guru Besar/ Pembina Utama Madya

FORMAT PENILAIAN (VALIDASI & PEER REVIEW)

LEMBAR

HASIL PENILAIAN SEJAWAT SEBIDANG ATAU PEER REVIEW

KARYA ILMIAH : JURNAL ILMIAH

1.19.

Jurnal Artikel Ilmiah : Entomopathogenic fungi of soils of freshwater swamps, tidal lowlands, peatlands, and highlands of south sumatra, Indonesia

Penulis Artikel Ilmiah : Arum Setiawan

Identitas Jurnal Artikel Ilmiah :

- a. Nama Jurnal : Biodiversitas
- b. Nomor/Volume/Hal : 6/19/2365-2373
- c. Edisi (bulan/tahun) : November/2018
- d. Penerbit : Jurusan Biologi FMIPA UNS
- e. Jumlah Halaman : 10

Kategori Publikasi Jurnal Ilmiah (beri \checkmark pada kategori yang tepat) :

- Jurnal Ilmiah Internasional Bereputasi
- Jurnal Ilmiah Internasional
- Jurnal Ilmiah Nasional Terakreditasi S1, S2
- Jurnal Ilmiah Nasional Terakreditasi S3, S4
- Jurnal Ilmiah Nasional Tidak Terakreditasi.

I. Hasil Penilaian Validasi :

No.	ASPEK	URAIAN/KOMENTAR PENILAIAN
1.	Indikasi Plagiasi	7 %
2.	Linearitas	V


II. Hasil Penilaian Peer Review :

Komponen Yang Dinilai	Nilai Maksimal Jurnal Ilmiah (isikan di kolom yang sesuai)					Nilai Akhir Yang Diperoleh
	Internasional Bereputasi (Maks 40)	Internasional (Maks 20)	Nasional Terakreditasi S1, S2 Maks 25	Nasional Terakreditasi S3, S4 Maks 20	Nasional tidak Terakreditasi (maks 10)	
Kelengkapan dan Kesesuaian unsur isi jurnal (10%)	4					4
Ruang lingkup dan kedalaman pembahasan (30%)	12					12
Kecukupan dan Kemutakhiran data/informasi dan metodologi (30%)	12					12
Kelengkapan unsur dan kualitas penerbit (30%)	12					12
Total = (100%)	40					40
Kontribusi Pengusul (Penulis Pertama /Anggota Utama)	Biodiversitas Vol. 19(6): 2365-2373. November/2018. Terindeks Scopus dengan SJR 0,27 Penulis ke 3 dari 3 penulis. Nilai maksimal 100%. Nilai pengusul: $(0,4 \times 1 \times 40)/2 = 8,0$					8,0

KOMENTAR/ULASAN PEER REVIEW	
• Kelengkapan dan Kesesuaian Unsur:	Format dengan unsur yang lengkap mulai dari abstrak hingga referensi. Acuan relatif baru dan terkait penelitian.
• Ruang Lingkup dan Kedalaman Pembahasan:	Masih dalam bidang ilmu terkait.
• Kecukupan & Kemutakhiran Data & Metodologi:	Data cukup layak karena banyak dan metode tidak ada yang baru.
• Kelengkapan Unsur & Kualitas Penerbit:	Penerbit Jurusan Biologi FMIPA UNS, cukup berkualitas karena kelengkapannya.

Yogyakarta, 11 Juli 2020

Penilai 2

tanda tangan 

Prof. Dr. Suwarno Haditsusanto

NIP 195411161983031002

Unit Kerja : Fakultas Biologi UGM

Bidang Ilmu : Biologi

Jabatan/Pangkat : Guru Besar/ Pembina Utama Madya