

First record of scalloped hammerhead shark *Sphyrna lewini* (Carcharhiniformes: Sphyrnidae) in freshwater habitat

MUHAMMAD IQBAL^{1*}, DONI SETIAWAN², ARUM SETIAWAN² & INDRA YUSTIAN²

¹Biology Program, Faculty of Science, Sriwijaya University, Jalan Padang Selasa 524, Palembang, Sumatera Selatan 30129, Indonesia.

²Department of Biology, Faculty of Science, Sriwijaya University, Jalan Raya Palembang-Prabumulih km 32, Indralaya, Sumatera Selatan 30662, Indonesia.

* Corresponding author [kpbosos26@yahoo.com]

Received 24 March 2020 | Accepted by V. Pešić: 19 April 2020 | Published online .. April 2020.

Abstract

A scalloped hammerhead shark *Sphyraena lewini* (Griffith & Smith, 1834) with c. 40 cm of total length was stranded and photographed on 6 April 2020 in Mahakam river, East Kalimantan province, Indonesia. This unexpected finding is represent a first ever know inland record of *S. lewini* in freshwater habitat.

Key words: *Sphyrna lewini*, record, freshwater, Kalimantan, Indonesia.

Introduction

The hammerhead sharks (family Sphyrnidae) unmistakable sharks have unique head shape with the eyes located at the tips of laterally expanded blades which resemble an hammer (Last & Stevens 1994). These function as a submarine-like bow plane to improve manouvrability and increase sensory capacity by enhancing stereoscopic vision and ability to triangulate sources of scent and electromagnetic signals (Compagno *et al.* 2005). The hammerhead sharks feed mainly at night, good reason for having well-developed senses other than vision (Moyle & Cech 2004).

Family Sphyrnidae is a small group of sharks containing two genera, with eight or ten described species (Ebert *et al.* 2013; Nelson *et al.* 2016; Gallagher & Klimley 2018). The scalloped hammerhead shark *Sphyraena lewini* is one of species of family Sphyrnidae that cosmopolitan in tropical and warm temperate seas (Last & Stevens 1994). The *S. lewini* is a large shark (370-420 cm of total length) with a maximum age of up to 35 years (Ebert *et al.* 2013). However, this species is recently listed as Critically Endangered by IUCN because global population trend estimated median reductions of 76–97%, with the highest probability of >80% reduction over three generation lengths or about 72 years (Rigby *et al.* 2019). As other species of hammerhead sharks, *S. lewini* inhabit marine, primarily continental shelf and occasionally brackish in Atlantic, India and Pacific (Nelson *et al.* 2016; Marie *et al.* 2017); including in Indonesian waters, from Sumatra in the west to the West Papua in the east (Gloerfelt-Tarp & Kailola 1984, Fahmi 2010, Chodriah & Setyadji 2015; Sentosa *et al.* 2016).

A recent inland record of *S. lewini* on the basis photographic evidence was reported in Mahakam river, East Kalimantan, Indonesia. This record presented a first ever known world wide record of *S. lewini* in freshwater habitat.

Materials and Methods

A specimen of *S. lewini* (c. 40 cm of total length) was stranded and photographed on 6 April 2020, at Mahakam river, Muara Baroh village, Muara Pahu subdistrict, Kutai Barat district, East Kalimantan province, Indonesia ($00^{\circ}19'25"S$, $116^{\circ}03'46"E$) (Fig. 1). This area is in inland freshwater habitat located c. 150 km distance from mouth of river. Photos of this shark were shown to us by local people. There is no specific details on measurements, but this small size suggest this species is very early young male juvenile shark, when small claspers are present (Fig. 2). Specimen is unpreserved for a museum collection. It is reported that this specimen requested by a staff of local animal conservation NGO in East Kalimantan, but we are unable to track further for checking specific informations. Identification is therefore based on the features that are visible in the photographs. Proportional morphological characters expressed as percentage of total length based on the photographs shown in Table 1.




Figure 1. Location of known *S. lewini* in Borneo. Yellow circles are *S. lewini* recorded in Bornean waters after Last *et al.* (2010), and red circle is recent inland record from Mahakam river, East Kalimantan province.



Figure 2. The *S. lewini* which stranded in Mahakam river at Muara Baroh village, Muara Pahu subdistrict, Kutai Barat district, East Kalimantan province (Photo: Yuni Sri Wahyuni).



Figure 3. Another side of *S. lewini* which stranded on 6 April 2020 in Mahakam river at Muara Baroh village, Muara Pahu subdistrict, Kutai Barat district, East Kalimantan province (Photo: Yuni Sri Wahyuni).

Table 1. Proportional morphological characters of *S. lewini* expressed as percentage of total length based on the photographs. S1 = specimen from Mahakam river, S2 = other specimens on the basis of Quattro *et al.* (2013).

Characters	S1	S2 (range)
Head length	24	20-24
Pre-first dorsal fin	28	24-28
Pre-second dorsal fin	61	57-61
Prepectoral length	24	19-24
Prepelvic length	47	43-47
Preanal length	59	59
Snout-vent length	48	44-48
Interdorsal space	20	20-25
Second dorsal-caudal space	6	5-6
Pectoral-pelvic space	20	15-20

Results and Discussions

An individual of stranded shark found in Mahakam river has brownish grey dorsally and pale ventrally, head laterally expanded into prominent keels resembling an hammer with maximum width 24-30% of total length; body fusiform and moderately slender; anterior profile of hammer curved anteriorly with median and lateral indentations; first dorsal fin origin about over or slightly behind pectoral fin insertion, first dorsal fin broad and relatively erect, pelvic fin posterior margin nearly straight, no other fin markings (Figs. 2 and 3). These characters are fitted well to the characters of *S. lewini* (Compagno & Niem 1998; Compagno *et al.* 2005; Last *et al.* 2010; Ebert *et al.* 2013). The *S. lewini* is very resemble to smooth hammerhead *S. zygaena*, its closest relative; but *S. lewini* is distinguished from *S. zygaena* by a median indentation (scallop) on anterior margin of head (Fig. 2 and 3) (Gilbert 1967). The Carolina hammerhead *Sphyraña gilberti* in western Atlantic Ocean is a cryptic species of *S. lewini*, and morphologically can only separate with *S. lewini* in the number of precaudal vertebrae (Quattro *et al.* 2013).

The *S. lewini* is a large endangered shark with a world wide distribution, observed in the open ocean but linked ontogenetically to coastal embayments for parturition and juvenile development (Daly-Engel *et al.* 2012). Populations from different locations are generally distinct (Compagno *et al.* 2005; Duncan *et al.* 2006; Ebert *et al.* 2013); and two distinct forms have been identified from molecular analysis of Borneo specimens but, these forms have not yet been clearly distinguished (Last *et al.* 2010). Some aspects of reproductive biology of the *S. lewini* have been studied (eg. Hazin *et al.* 2001; Duncan 2006; Bejarano-Alvarez *et al.* 2010; Salinas-de-León *et al.* 2017; Marie *et al.* 2019), but it is still lacking in Indonesia and Borneo still (Last *et al.* 2010; Wahyudin *et al.* 2019).

The occurrence of up to 150 inland *S. lewini* of in Mahakam river is unexpected. This record is a first ever known report for *S. lewini* in freshwater habitat. *Sphyraña lewini* and other hammerhead sharks are among the most intriguing yet imperiled groups of large sharks globally that understanding of their ecology, diet, life histories and movements was challenged by a lack of studies (Gallagher & Klimley 2018). Nursery populations of *S. lewini* linked by continuous coastline have high connectivity, but that oceanic dispersal by females is rare (Duncan *et al.* 2006). This species is a cosmopolitan and seasonal marine migratory species where they can move up to 600 to 1.700 km away (Diemer *et al.* 2011; Ketchum *et al.* 2012; Hoyos-Padilla *et al.* 2014), but there is no information on movement to freshwater habitat.

Specimen of *S. lewini* found in Mahakam river is a very young individual based on its small size of c. 40 cm of total length. The present of very young *S. lewini* in Mahakam river indicate this river possibly a nursery area for this species. No information available if *S. lewini* occur in freshwater or tolerate low salinities (Compagno & Niem 1998; Compagno 2002; Compagno *et al.* 2005; Last *et al.* 2010; Ebert *et al.* 2013; Gallagher & Klimley 2018). In Indonesia, recent records of young sharks entering freshwater habitat

have been significantly increase (Iqbal *et al.* 2019a, b, c). The invasion to freshwater habitats has been known independently many times in sharks and other elasmobranchs evolution (Lucifora *et al.* 2015). The reasons why affecting the poor penetration of sharks into freshwater habitat are still unclear, however, an important assumption could be the high urea requirement of many proteins in marine sharks (Ballantyne & Robinson 2010). More information and further research are needed to confirm the status of *S. lewini* in freshwater habitat, particularly in Mahakam river and other parts of Bornean waters. Further study in the future should have address if it is an incidental or a regular movements.

Acknowledgements

We are very grateful to Cepi M. Kurniawan and Innal Rahman who draw our attention to the presence of *Sphyrna lewini* in Mahakam river, and particularly to Yuni Sri Wahyuni sharing us her photos. We thank anonymous reviewers who provided invaluable suggestions to make this paper improve.

References

- Ballantyne, J.S. & Robinson, J.W. (2010) Freshwater elasmobranchs: a review of their physiology and biochemistry. *Journal of Comparative Physiology B*, 180(4), 475-93.
- Bejarano-Álvarez, M., Galvan-Magana, F. & Ochoa-Báez, R.I. (2011) Reproductive biology of the Scalloped Hammerhead Shark *Sphyrna lewini* (Chondrichthyes: Sphyrnidae) off south-west Mexico. *Aqua*, 17, 11-22.
- Chodriah, U. & Setyadi, B. (2015) Some biological aspects of Scalloped Hammerhead Sharks (*Sphyrna lewini* Griffith & Smith, 1834) caught from coastal fisheries in the Eastern Indian Ocean. *Indonesian Fisheries Research Journal*, 21(2), 91-97.
- Compagno, L.J.V. (2002) Freshwater and estuarine elasmobranch surveys in the Indo-Pacific Region: threats, distribution and speciation. In: Fowler, S.L., Reed, T.M. & Dipper, F.A. (Eds), *Elasmobranch Biodiversity, conservation and management: Proceedings of the International seminar and workshop, Sabah, Malaysia, July 1997*. IUCN SSC Shark Specialist Group, Switzerland and Cambridge, pp. 185-193.
- Compagno, L.J.V. & Niem, V.H. (1998) Sphyrnidae. Requiem sharks. In: Carpenter K.E. & Niem, V.H. (Eds.), *FAO identification guide for fishery purposes. The living marine resources of the Western Central Pacific) Volume 2. Cephalopods, crustaceans, holothurians and sharks*. FAO, Rome, pp. 1361-1367.
- Compagno, L., Dando, M. & Fowler, S. (2005) *A field guide to the sharks of the world*. Princeton University Press, New Jersey, 368 p.
- Daly-Engel ,T.S., Seraphin, K.D., Holland, K.N., Coffey, J.P., Nance, H.A., Toonen, R.J. & Bowen, B.W. (2012) Global phylogeography with mixed-marker analysis reveals male-mediated dispersal in the Endangered Scalloped Hammerhead Shark (*Sphyrna lewini*). *PLoS ONE*, 7(1), e29986.
- Diemer, K.M., Mann, B.Q. & Hussey, N.E. (2011) Distribution and movement of Scalloped Hammerhead *Sphyrna lewini* and Smooth Hammerhead *Sphyrna zygaena* sharks along the east coast of southern Africa. *African Journal of Marine Science*, 33(2), 229-238.
- Duncan, K.M., Martin, A.P., Bowen, B.W. & De Couet, H.G. (2006) Global phylogeography of the Scalloped Hammerhead Shark (*Sphyrna lewini*). *Molecular Ecology*, 15(8), 2239-2251.
- Ebert, D., Fowler, S. & Compagno, L. (2013) *Sharks of the world, a fully illustrated guide*. Wild Nature Press, Plymouth, 528 p.
- Gallagher, A.J & Klimley, A.P. (2018) The biology and conservation status of the large hammerhead shark complex: the great, scalloped, and smooth hammerheads. *Reviews in Fish Biology and Fisheries*, 28(3), 1-18.
- Gilbert, C.R. (1967) A revision of the Hammerhead Sharks (Family Sphyrnidae). *Proceedings of the United States National Museum*, 119(3539), 1-98.
- Quattro, J.M., Driggers, W.B., Grady, J.M., Ulrich, G.F. & Roberts, M.A. (2013) *Sphyrna gilberti* sp. nov., a new hammerhead shark (Carcharhiniformes, Sphyrnidae) from the western Atlantic Ocean. *Zootaxa*, 3702(2), 159-178.

- Gloerfelt-Tarp, T. & Kailola, P.J. (1984) *Trawled fishes of southern Indonesia and northwestern Australia*. Australian Development Assistance Bureau, Australia, 407 p.
- Fahmi. (2010) Sharks and rays in Indonesia. *Marine Research in Indonesia*, 35(1), 43-54.
- Hazin, F., Fischer, A. & Broadhurst, M. (2001) Aspects of reproductive biology of the Scalloped Hammerhead Shark, *Sphyrna Lewini*, off Northeastern Brazil. *Environmental Biology of Fishes*, 61, 151–159.
- Hoyos-Padilla, E.M., Ketchum, J.T., Klimley, A.P. & Galván-Magana, F. (2014) Ontogenetic migration of a female Scalloped Hammerhead Shark *Sphyrna lewini* in the Gulf of California. *Animal Biotelemetry*, 2(17), 1-9.
- Iqbal, M., Nurnawati, E., Setiawan, A., Dahlan, Z. & Yustian, I. (2019) First photographic inland records of bull shark *Carcharhinus leucas* (Carcharhiniformes: Carcharhinidae) in Sumatran waters, Indonesia. *Ecologica Montenegrina*, 22, 171-176.
- Iqbal, M., Saputra, R.F., Setiawan, A. & Yustian, I. (2019) First photographic inland record of blacktip reef sharks *Carcharhinus melanopterus* (Carcharhiniformes: Carcharhinidae) in Indonesian waters. *Ecologica Montenegrina*, 24, 6-10.
- Iqbal, M., Setiawan, A. & Yustian, I. (2019) First inland record of bull shark *Carcharhinus leucas* (Carcharhiniformes: Carcharhinidae) in Indonesian Borneo. *Ecologica Montenegrina*, 24, 52-57.
- Ketchum, J.T., Hearn, A., Klimley, A.P., Peñaherrera, C., Espinoza, E., Bessudo, S., Soler, G. & Arauz, R. (2014) Inter-island movements of scalloped hammerhead sharks (*Sphyrna lewini*) and seasonal connectivity in a marine protected area of the eastern tropical Pacific. *Marine Biology*, 161(4), 939-951.
- Last, P.R. & Stevens, J.D. (1994) *Sharks and Rays of Australia*. CSIRO, Australia. 513 p.
- Last, P.R., White, W. T., Caira, J. N., Dharmadi. Fahmi., Jensen, K., Lim, A.P.K., Manjaji-Matsumoto, B.M., Naylor, G.J.P., Pogonoski, J.J. Stevens, J.D. & Yearsley, G.K. (2010) *Sharks and rays of Borneo*. CSIRO, Collingwood, 298 p.
- Lucifora, L.O., de Carvalho, M.R., Kyne, P.M. & White, W.T. (2015) Freshwater sharks and rays. *Current Biology*, 25, R971–R973.
- Marie, D.A., Miller, C., Cawich, C., Piovano, S. & Rico, C. (2017) Fisheries-independent surveys identify critical habitats for young Scalloped Hammerhead Sharks (*Sphyrna lewini*) in the Rewa Delta, Fiji. *Scientific Reports*, 7(17273), 1-12.
- Marie, A.D., Herbinger, C., Fullsack, P. & Rico, C. (2019) First reconstruction of kinship in a Scalloped Hammerhead Shark aggregation reveals the mating patterns and breeding sex ratio. *Frontiers in Marine Science* 6:676. doi: 10.3389/fmars.2019.00676.
- Moyle, P.B. & Cech, J.J. (2004) *Fishes, an introduction to ichthyology*. Prentice Hall, Upper Saddle River, 726 p.
- Nelson, J.S., Terry, C., Grande, T. C. & Wilson, M.V.H. (2016) *Fishes of the World*. John Wiley and Sons, Hoboken, 752 pp.
- Quattro, J.M., Driggers, W.B., Grady, J.M., Ulrich, G.F. & Roberts, M.A. (2013) *Sphyrna gilberti* sp. nov., a new hammerhead shark (Carcharhiniformes, Sphyrnidae) from the western Atlantic Ocean. *Zootaxa*, 3702(2), 159-178.
- Rigby, C.L., Dulvy, N.K., Barreto, R., Carlson, J., Fernando, D., Fordham, S., Francis, M.P., Herman, K., Jabado, R.W., Liu, K.M., Marshall, A., Pacourea, N., Romanov, E., Sherley, R.B. & Winker, H. (2019) *Sphyrna lewini*. The IUCN Red List of Threatened Species 2019: e.T39385A2918526. Available from: <https://dx.doi.org/10.2305/IUCN.UK.2019-3.RLTS.T39386A2920499.en>. (08 April 2020).
- Salinas-de-León, P., Hoyos-Padilla, E.M. & Pochet, F. (2017) First observation on the mating behaviour of the endangered Scalloped Hammerhead Shark *Sphyrna lewini* in the Tropical Eastern Pacific. *Environmental Biology of Fishes*, 100, 1603-1608.
- Sentosa, A.S., Dharmadi, D. & Tjahjo, D.W.H. (2016) Parameter populasi hiu martil (*Sphyrna lewini* Griffith & Smith, 1834) di perairan selatan Nusa Tenggara. *Jurnal Penelitian Perikanan Indonesia*, 22(4), 253-262.
- Wahyudin, I., Kamal, M.M., Fahrudin, A. & Boer, M. (2019) Length-weight relationship and reproductive size of silky shark *Carcharhinus falciformis* and scalloped hammerhead shark *Sphyrna lewini* collected in Tanjung Luar fish landing port, East Lombok, Indonesia. *AACL Bioflux*, 12(1), 355-362.

**SJR**

Scimago Journal & Country Rank

Enter Journal Title, ISSN or Publisher Name

[Home](#)[Journal Rankings](#)[Country Rankings](#)[Viz Tools](#)[Help](#)[About Us](#)

Ads by Google

[Stop seeing this ad](#)[Why this ad?](#)

Ecologica Montenegrina

7

H Index

Country	Montenegro - SIR Ranking of Montenegro
Subject Area and Category	Agricultural and Biological Sciences Animal Science and Zoology Ecology, Evolution, Behavior and Systematics Insect Science Plant Science
Publisher	
Publication type	Journals
ISSN	23370173, 23369744
Coverage	2014-2020
Scope	Ecologica Montenegrina (ISSN 2336-9744 (online) ISSN 2337-0173 (print)) is peer-reviewed journal in which scientific articles and reports are quickly published. The papers are in the fields of taxonomy, biogeography and ecology (for example: new taxa for science, taxonomic revision, and/or fundamental ecology and biogeography papers). Open access publishing option is strongly encouraged for authors with research grants and other funds. For those without grants/funds, all accepted manuscripts will be published but access is secured for subscribers only.
Homepage	
How to publish in this journal	
Contact	
Join the conversation about this journal	











Ads by Google

[Stop seeing this ad](#)[Why this ad?](#)

Quartiles







ECOLOGICA MONTENEGRINA



Volume 30
April 2020

Ecologica Montenegrina

HOME ABOUT LOGIN REGISTER SEARCH
 CURRENT ARCHIVES ANNOUNCEMENTS

Home > About the Journal > Editorial Team

Editorial Team

Editor in Chief

Vladimir Pešić, Department of Biology, University of Montenegro, Montenegro

Editorial Board

Badamdorj Bayartogtokh, National University of Mongolia, Mongolia
Holger Braun, Museo de La Plata, Argentina

Dr Patricio De los Rios Escalante, Universidad Católica de Temuco, Chile

Sergey G. Ermilov, Tyumen State University, Tyumen, Russia, Russian Federation

Igor Dovgal, A. Kovalevsky Institute of Marine Biological Research, Russian Federation

Thibault Datry, Irstea, UR MALY, Centre de Lyon-Villeurbanne, France

Zoltán Fehér, Natural History Museum Vienna 3rd Zoology Department, Austria

Dilian Georgiev, University of Plovdiv, Bulgaria

Reinhard Gerecke

Wolfram Graf, University of Natural Resources and Life Sciences, Vienna, Austria

Michał Grabowski, Department of Invertebrate Zoology and Hydrobiology, University of Łódź, Poland

Clemens Grosser, Germany

Borislav Gueorguiev, National Museum of Natural History, Sofia, Bulgaria

Sead Hadžiblahović, Environmental Protection Agency of Montenegro, Montenegro

Dr Laith A. Jawad, Flat Bush, Manukau, Auckland, New Zealand, New Zealand

Dr Roman Yakovlev, Altai State University, Russian Federation

Max Kasparek, Zoology in the Middle East, Germany

Gordan S. Karaman, Montenegrin Academy of Sciences, Montenegro

Jeno Kontschán, Plant Protection Institute, Centre for Agricultural Research, Hungarian Academy of Sciences, Hungary

Andrey Kostianoy, P.P.Shirshov Institute of Oceanology, Russian Academy of Sciences, Moscow, Russia, Russian Federation

Katarina Ljubisavljević, Montenegro

Dr Andrzej Zawal, Department of Invertebrate Zoology & Limnology, University of Szczecin, 71-415 Szczecin, Waska 13, Poland, Poland

Slobodan Evgenije Makarov, University of Belgrade Faculty of Biology, Serbia

Dávid Murányi, Hungarian Natural History Museum Department of Zoology

Baross u. 13, H-1088 Budapest, Hungary, Hungary

Momir M Paunović, Institute for Biological Research "Sinisa Stankovic" University of Belgrade Republic of Serbia Bulevar despota Stefana 142, 11000 Belgrade, Serbia, Serbia

Radmila Petanović, University of Belgrade - Faculty of Agriculture Department of Entomology and Agricultural Zoology Nemanjina 6, P.O. BOX 127 11080 Belgrade - Zemun SERBIA, Serbia

Prof. Alireza Saboori, University of Tehran, Iran, Islamic Republic of

Zoltán Sándor Varga, University of Debrecen, Hungary

Dr Anton V. Volynkin, Altai State University

Paul Selden, University of Kansas, United States

Nelli Grigoryevna Sergeeva, Institute of Marine Biological Researches RAS, Russian Federation

Harry Smit, Naturalis Biodiversity Center, Netherlands

Boris Sket, Univerza v Ljubljani, Slovenia

Vesna Vukasinovic Petic, Department of Technology, University of Montenegro, Podgorica, Montenegro

OPEN JOURNAL SYSTEMS

Journal Help

USER

Username

Password

Remember me

NOTIFICATIONS

- [View](#)
- [Subscribe](#)

LANGUAGE

Select Language

JOURNAL CONTENT

Search

Search Scope

Browse

- [By Issue](#)
- [By Author](#)
- [By Title](#)
- [Other Journals](#)

FONT SIZE

INFORMATION

- [For Readers](#)
- [For Authors](#)
- [For Librarians](#)

KEYWORDS

Amphipoda Black Sea.
Bythinella Carabidae.
China China. Croatia.
Gastropoda Greater Antilles.
Greece Halacaroidea.
Hirudinea Hungary.
Hydrobiidae.
Hydryphantidae.
Macedonia Mollusca.
Montenegro.
Montenegro. New species Republic of.
Adygaea Salicidae.
Serbia. Taxonomy.
Turkey. Ukraine.
biodiversity checklist.
distribution ecology.
faunistics female.
identification key.
jumping spiders.

ISSN: 2336-9744

Ecologica Montenegrina

HOME ABOUT LOGIN REGISTER SEARCH
 CURRENT ARCHIVES ANNOUNCEMENTS

Home > Archives > Vol 30 (2020)

Vol 30 (2020)

Table of Contents

Articles

Contribution to knowledge of the Balkan Lepidoptera (Lepidoptera: Macrolepidoptera).	PDF 1-27
Stoyan Beshkov, Ana Nahirnić	
The study of fidelity measures in the context of using them as a threshold criterion in the allocation of diagnostic species	PDF 28-36
Igor Goncharenko, Alexander Senchylo	
Influence of environmental variability on the body condition of the mangrove horseshoe crab <i>Carcinoscorpius rotundicauda</i> from Banyuasin Estuarine, South Sumatra, Indonesia	PDF 37-46
Nita P. Sari, Noviantrio Gulo, Miftahun Najah, Fauziyah .	
Trophic Connections of Leafroller Moths (Lepidoptera: Tortricidae) and Oaks in Sofia Region, Bulgaria	PDF 47-59
Gergana Zaemdzikova	
Some new <i>Bythinella</i> spp. from southern Greece (Gastropoda: Bythinellidae)	PDF 60-67
Peter Glöer, Hans-Jürgen Hirschfelder	
Diet of the Lesser Spotted Eagle (<i>Clanga pomarina</i>) in Amvrakikos Wetlands National Park, Greece	PDF 68-76
Haralambos Alivizatos, Dimitris Papandropoulos, Stamatis Zogaris	
A new monotypic genus of the subfamily Chilecomadiinae Dyar, 1940 (Lepidoptera: Cossidae) from South America	PDF 77-81
Fernando C. Penco, Roman V. Yakovlev, Artem E. Naydenov	
On the taxonomy of <i>Meganola cretacea</i> (Hampson, 1914) with description of two new species (Lepidoptera, Nolidae, Nolinae).	PDF 82-92
Gyula M. László	
Contribution to the Knowledge of the Balkan Lepidoptera: Nocturnal Lepidoptera of Western Serbia (Lepidoptera).	PDF 93-112
Stoyan Beshkov, Ana Nahirnić	
First record of scalloped hammerhead shark <i>Sphyrna lewini</i> (Carcharhiniformes: Sphyrnidae) in freshwater habitat	PDF 113-118
Muhammad Iqbal, Doni Setiawan, Arum Setiawan, Indra Yustian	
Second record of goblin shark <i>Mitsukurina owstoni</i> (Lamniformes: Mitsukurinidae) in Indonesian waters	PDF 119-123
Muhammad Iqbal, Arum Setiawan, Doni Setiawan, Indra Yustian	
Two new valvatoid genera (Gastropoda, Hydrobiidae) from Morocco	PDF 124-128
Peter Glöer, Youness Mabrouki, Abdelkhaleq Fouzi Taybi	

[OPEN JOURNAL SYSTEMS](#)

[Journal Help](#)

USER

Username
 Password
 Remember me

NOTIFICATIONS

- [View](#)
- [Subscribe](#)

LANGUAGE

Select Language

JOURNAL CONTENT

Search
 Search Scope

Browse

- [By Issue](#)
- [By Author](#)
- [By Title](#)
- [Other Journals](#)

FONT SIZE

INFORMATION

- [For Readers](#)
- [For Authors](#)
- [For Librarians](#)

KEYWORDS

[Amphipoda](#) [Black Sea](#).
[Bythinella](#) [Carabidae](#).
[China](#) [China](#). [Croatia](#).
[Gastropoda](#) [Greater Antilles](#).
[Greece](#) [Halacaroidea](#).
[Hirudinea](#) [Hungary](#).
[Hydrobiidae](#).
[Hydryphantidae](#).
[Macedonia](#) [Mollusca](#).
[Montenegro](#).
[Montenegro](#). [New species](#) [Republic of Adygeya](#).
[Salticidae](#).
[Serbia](#). [Taxonomy](#).
[Turkey](#), [Ukraine](#).
[biodiversity](#), [checklist](#).
[distribution](#), [ecology](#).
[faunistics](#), [female identification](#), [key](#),
[jumping](#), [spiders](#)

First Record of Scalloped hammerhead shark *Sphyrna lewini* (Carcharhiniformes : Sphyrnidae) in freshwater habitat

By Arum Setiawan



Ecologica Montenegrina
ISSN 2339-8744 (online); ISSN 2339-0173 (print)

Ecologica Montenegrina 30: 113-118 (2020)
This journal is available online at: www.biota.org/em
<http://dx.doi.org/10.37828/em.2020.30.10>

Article

First record of scalloped hammerhead shark *Sphyrna lewini* (Carcharhiniformes: Sphyrnidae) in freshwater habitat

MUHAMMAD IQBAL^{1*}, DONI SETIAWAN², ARUM SETIAWAN² & INDRA YUSTIAN²

¹Biology Program, Faculty of Science, Sriwijaya University, Jalan Padang Selasa 524, Palembang, Sumatera Selatan 30129, Indonesia.

²Department of Biology, Faculty of Science, Sriwijaya University, Jalan Raya Palembang-Prabumulih km 32, Indralaya, Sumatera Selatan 30662, Indonesia.

* Corresponding author [kpbosos26@yahoo.com]

Received 24 March 2020 | Accepted by V. Pešić: 19 April 2020 | Published online .. April 2020.

Abstract

A scalloped hammerhead shark *Sphyraena lewini* (Griffith & Smith, 1834) with c. 40 cm of total length was stranded and photographed on 6 April 2020 in Mahakam river, East Kalimantan province, Indonesia. This unexpected finding is represent a first ever known inland record of *S. lewini* in freshwater habitat.

Key words: *Sphyrna lewini*, record, freshwater, Kalimantan, Indonesia.

Introduction

The hammerhead sharks (family Sphyrnidae) unmistakable sharks have unique head shape with the eyes located at the tips of laterally expanded blades which resemble an hammer (Last & Stevens 1994). These function as a submarine-like bow plane to improve manouvrability and increase sensory capacity by enhancing stereoscopic vision and ability to triangulate sources of scent and electromagnetic signals (Compagno *et al.* 2005). The hammerhead sharks feed mainly at night, good reason for having well-developed senses other than vision (Moyle & Cech 2004).

Family Sphyrnidae is a small group of sharks containing two genera, with eight or ten described species (Ebert *et al.* 2013; Nelson *et al.* 2016; Gallagher & Klimley 2018). The scalloped hammerhead shark *Sphyraena lewini* is one of species of family Sphyrnidae that cosmopolitan in tropical and warm temperate seas (Last & Stevens 1994). The *S. lewini* is a large shark (370-420 cm of total length) with a maximum age of up to 35 years (Ebert *et al.* 2013). However, this species is recently listed as Critically Endangered by IUCN because global population trend estimated median reductions of 76–97%, with the highest probability of >80% reduction over three generation lengths or about 72 years (Rigby *et al.* 2019). As other species of hammerhead sharks, *S. lewini* inhabit marine, primarily continental shelf and occasionally brackish in Atlantic, India and Pacific (Nelson *et al.* 2016; Marie *et al.* 2017); including in Indonesian waters, from Sumatra in the west to the West Papua in the east (Gloerfelt-Tarp & Kailola 1984, Fahmi 2010, Chodrijah & Setyadji 2015; Sentosa *et al.* 2016).

A recent inland record of *S. lewini* on the basis photographic evidence was reported in Mahakam river, East Kalimantan, Indonesia. This record presented a first ever known world wide record of *S. lewini* in freshwater habitat.

Materials and Methods

A specimen of *S. lewini* (c. 40 cm of total length) was stranded and photographed on 6 April 2020, at Mahakam river, Muara Baroh village, Muara Pahu subdistrict, Kutai Barat district, East Kalimantan province, Indonesia ($00^{\circ}19'25"S$, $116^{\circ}03'46"E$) (Fig. 1). This area is in inland freshwater habitat located c. 150 km distance from mouth of river. Photos of this shark were shown to us by local people. There is no specific details on measurements, but this small size suggest this species is very early young male juvenile shark, when small claspers are present (Fig. 2). Specimen is unpreserved for a museum collection. It is reported that this specimen requested by a staff of local animal conservation NGO in East Kalimantan, but we are unable to track further for checking specific informations. Identification is therefore based on the features that are visible in the photographs. Proportional morphological characters expressed as percentage of total length based on the photographs shown in Table 1.




Figure 1. Location of known *S. lewini* in Borneo. Yellow circles are *S. lewini* recorded in Bornean waters after Last *et al.* (2010), and red circle is recent inland record from Mahakam river, East Kalimantan province.



Figure 2. The *S. lewini* which stranded in Mahakam river at Muara Baroh village, Muara Pahu subdistrict, Kutai Barat district, East Kalimantan province (Photo: Yuni Sri Wahyuni).



Figure 3. Another side of *S. lewini* which stranded on 6 April 2020 in Mahakam river at Muara Baroh village, Muara Pahu subdistrict, Kutai Barat district, East Kalimantan province (Photo: Yuni Sri Wahyuni).

Table 1. Proportional morphological characters of *S. lewini* expressed as percentage of total length based on the photographs. S1 = specimen from Mahakam river, S2 = other specimens on the basis of Quattro *et al.* (2013).

Characters	S1	S2 (range)
Head length	24	20-24
Pre-first dorsal fin	28	24-28
Pre-second dorsal fin	61	57-61
Prepectoral length	24	19-24
Prepelvic length	47	43-47
Preanal length	59	59
Snout-vent length	48	44-48
Interdorsal space	20	20-25
Second dorsal-caudal space	6	5-6
Pectoral-pelvic space	20	15-20

Results and Discussions

An individual of stranded shark found in Mahakam river has brownish grey dorsally and pale ventrally, head laterally expanded into prominent keels resembling an hammer with maximum width 24-30% of total length; body fusiform and moderately slender; anterior profile of hammer curved anteriorly with median and lateral indentations; first dorsal fin origin about over or slightly behind pectoral fin insertion, first dorsal fin broad and relatively erect, pelvic fin posterior margin nearly straight, no other fin markings (Figs. 2 and 3). These characters are fitted well to the characters of *S. lewini* (Compagno & Niem 1998; Compagno *et al.* 2005; Last *et al.* 2010; Ebert *et al.* 2013). The *S. lewini* is very resemble to smooth hammerhead *S. zygaena*, its closest relative; but *S. lewini* is distinguished from *S. zygaena* by a median indentation (scallop) on anterior margin of head (Fig. 2 and 3) (Gilbert 1967). The Carolina hammerhead *Sphyraena gilberti* in western Atlantic Ocean is a cryptic species of *S. lewini*, and morphologically can only separate with *S. lewini* in the number of precaudal vertebrae (Quattro *et al.* 2013).

The *S. lewini* is a large endangered shark with a world wide distribution, observed in the open ocean but linked ontogenetically to coastal embayments for parturition and juvenile development (Daly-Engel *et al.* 2012). Populations from different locations are generally distinct (Compagno *et al.* 2005; Duncan *et al.* 2006; Ebert *et al.* 2013); and two distinct forms have been identified from molecular analysis of Borneo specimens but, these forms have not yet been clearly distinguished (Last *et al.* 2010). Some aspects of reproductive biology of the *S. lewini* have been studied (eg. Hazin *et al.* 2001; Duncan 2006; Bejarano-Alvarez *et al.* 2010; Salinas-de-León *et al.* 2017; Marie *et al.* 2019), but it is still lacking in Indonesia and Borneo still (Last *et al.* 2010; Wahyudin *et al.* 2019).

The occurrence of up to 150 inland *S. lewini* of in Mahakam river is unexpected. This record is a first ever known report for *S. lewini* in freshwater habitat. *Sphyraena lewini* and other hammerhead sharks are among the most intriguing yet imperiled groups of large sharks globally that understanding of their ecology, diet, life histories and movements was challenged by a lack of studies (Gallagher & Klimley 2018). Nursery populations of *S. lewini* linked by continuous coastline have high connectivity, but that oceanic dispersal by females is rare (Duncan *et al.* 2006). This species is a cosmopolitan and seasonal marine migratory species where they can move up to 600 to 1.700 km away (Diemer *et al.* 2011; Ketchum *et al.* 2012; Hoyos-Padilla *et al.* 2014), but there is no information on movement to freshwater habitat.

Specimen of *S. lewini* found in Mahakam river is a very young individual based on its small size of c. 40 cm of total length. The present of very young *S. lewini* in Mahakam river indicate this river possibly a nursery area for this species. No information available if *S. lewini* occur in freshwater or tolerate low salinities (Compagno & Niem 1998; Compagno 2002; Compagno *et al.* 2005; Last *et al.* 2010; Ebert *et al.* 2013; Gallagher & Klimley 2018). In Indonesia, recent records of young sharks entering freshwater habitat

have been significantly increase (Iqbal *et al.* 2019a, b, c). The invasion to freshwater habitats has been known independently many times in sharks and other elasmobranchs evolution (Lucifora *et al.* 2015). The reasons why affecting the poor penetration of sharks into freshwater habitat are still unclear, however, an important assumption could be the high urea requirement of many proteins in marine sharks (Ballantyne & Robinson 2010). More information and further research are needed to confirm the status of *S. lewini* in freshwater habitat, particularly in Mahakam river and other parts of Bornean waters. Further study in the future should have address if it is an incidental or a regular movements.

Acknowledgements

We are very grateful to Cepi M. Kurniawan and Innal Rahman who draw our attention to the presence of *Sphyrna lewini* in Mahakam river, and particularly to Yuni Sri Wahyuni sharing us her photos. We thank anonymous reviewers who provided invaluable suggestions to make this paper improve.

References

- Ballantyne, J.S. & Robinson, J.W. (2010) Freshwater elasmobranchs: a review of their physiology and biochemistry. *Journal of Comparative Physiology B*, 180(4), 475-93.
- Bejarano-Alvarez, M., Galvan-Magana, F. & Ochoa-Báez, R.I. (2011) Reproductive biology of the Scalloped Hammerhead Shark *Sphyrna lewini* (Chondrichthyes: Sphyrnidae) off south-west Mexico. *Aqua*, 17, 11-22.
- Chodrijah, U. & Setyadiji, B. (2015) Some biological aspects of Scalloped Hammerhead Sharks (*Sphyrna lewini* Griffith & Smith, 1834) caught from coastal fisheries in the Eastern Indian Ocean. *Indonesian Fisheries Research Journal*, 21(2), 91-97.
- Compagno, L.J.V. (2002) Freshwater and estuarine elasmobranch surveys in the Indo-Pacific Region: threats, distribution and speciation. In: Fowler, S.L., Reed, T.M. & Dipper, F.A. (Eds), *Elasmobranch Biodiversity, conservation and management: Proceedings of the International seminar and workshop, Sabah, Malaysia, July 1997*. IUCN SSC Shark Specialist Group, Switzerland and Cambridge, pp. 185-193.
- Compagno, L.J.V. & Niem, V.H. (1998) Sphyrnidae. Requiem sharks. In: Carpenter K.E. & Niem, V.H. (Eds.), *FAO identification guide for fishery purposes. The living marine resources of the Western Central Pacific Volume 2. Cephalopods, crustaceans, holothurians and sharks*. FAO, Rome, pp. 1361-1367.
- Compagno, L., Dando, M. & Fowler, S. (2005) *A field guide to the sharks of the world*. Princeton University Press, New Jersey, 368 p.
- Daly-Engel, T.S., Seraphin, K.D., Holland, K.N., Coffey, J.P., Nance, H.A., Toonen, R.J. & Bowen, B.W. (2012) Global phylogeography with mixed-marker analysis reveals male-mediated dispersal in the Endangered Scalloped Hammerhead Shark (*Sphyrna lewini*). *PLoS ONE*, 7(1), e29986.
- Diemer, K.M., Mann, B.Q. & Hussey, N.E. (2011) Distribution and movement of Scalloped Hammerhead *Sphyrna lewini* and Smooth Hammerhead *Sphyrna zygaena* sharks along the east coast of southern Africa. *African Journal of Marine Science*, 33(2), 229-238.
- Duncan, K.M., Martin, A.P., Bowen, B.W. & De Couet, H.G. (2006) Global phylogeography of the Scalloped Hammerhead Shark (*Sphyrna lewini*). *Molecular Ecology*, 15(8), 2239-2251.
- Ebert, D., Fowler, S. & Compagno, L. (2013) *Sharks of the world, a fully illustrated guide*. Wild Nature Press, Plymouth, 528 p.
- Gallagher, A.J. & Klimley, A.P. (2018) The biology and conservation status of the large hammerhead shark complex: the great, scalloped, and smooth hammerheads. *Reviews in Fish Biology and Fisheries*, 28(3), 1-18.
- Gilbert, C.R. (1967) A revision of the Hammerhead Sharks (Family Sphyrnidae). *Proceedings of the United States National Museum*, 119(3539), 1-98.
- Quattro, J.M., Driggers, W.B., Grady, J.M., Ulrich, G.F. & Roberts, M.A. (2013) *Sphyrna gilberti* sp. nov., a new hammerhead shark (Carcharhiniformes, Sphyrnidae) from the western Atlantic Ocean. *Zootaxa*, 3702(2), 159-178.

- Gloerfelt-Tarp, T. & Kailola, P.J. (1984) *Trawled fishes of southern Indonesia and northwestern Australia*. Australian Development Assistance Bureau, Australia, 407 p.
- Fahmi. (2010) Sharks and rays in Indonesia. *Marine Research in Indonesia*, 35(1), 43-54.
- Hazin, F., Fischer, A. & Broadhurst, M. (2001) Aspects of reproductive biology of the Scalloped Hammerhead Shark, *Sphyrna Lewini*, off Northeastern Brazil. *Environmental Biology of Fishes*, 61, 151–159.
- Hoyos-Padilla, E.M., Ketchum, J.T., Klimley, A.P. & Galván-Magana, F. (2014) Ontogenetic migration of a female Scalloped Hammerhead Shark *Sphyrna lewini* in the Gulf of California. *Animal Biotelemetry*, 2(17), 1-9.
- Iqbal, M., Nurnawati, E., Setiawan, A., Dahlan, Z. & Yustian, I. (2019) First photographic inland records of bull shark *Carcharhinus leucas* (Carcharhiniformes: Carcharhinidae) in Sumatran waters, Indonesia. *Ecologica Montenegrina*, 22, 171-176.
- Iqbal, M., Saputra, R.F., Setiawan, A. & Yustian, I. (2019) First photographic inland record of blacktip reef sharks *Carcharhinus melanopterus* (Carcharhiniformes: Carcharhinidae) in Indonesian waters. *Ecologica Montenegrina*, 24, 6-10.
- Iqbal, M., Setiawan, A. & Yustian, I. (2019) First inland record of bull shark *Carcharhinus leucas* (Carcharhiniformes: Carcharhinidae) in Indonesian Borneo. *Ecologica Montenegrina*, 24, 52-57.
- Ketchum, J.T., Hearn, A., Klimley, A.P., Peñaherrera, C., Espinoza, E., Bessudo, S., Soler, G. & Arauz, R. (2014) Inter-island movements of scalloped hammerhead sharks (*Sphyrna lewini*) and seasonal connectivity in a marine protected area of the eastern tropical Pacific. *Marine Biology*, 161(4), 939-951.
- Last, P.R. & Stevens, J.D. (1994) *Sharks and Rays of Australia*. CSIRO, Australia. 513 p.
- Last, P.R., White, W. T., Caira, J. N., Dharmadi, Fahmi., Jensen, K., Lim, A.P.K., Manjaji-Matsumoto, B.M., Naylor, G.J.P., Pogonoski, J.J. Stevens, J.D. & Yearsley, G.K. (2010) *Sharks and rays of Borneo*. CSIRO, Collingwood, 298 p.
- Lucifora, L.O., de Carvalho, M.R., Kyne, P.M. & White, W.T. (2015) Freshwater sharks and rays. *Current Biology*, 25, R971–R973.
- Marie, D.A., Miller, C., Cawich, C., Piovano, S. & Rico, C. (2017) Fisheries-independent surveys identify critical habitats for young Scalloped Hammerhead Sharks (*Sphyrna lewini*) in the Rewa Delta, Fiji. *Scientific Reports*, 7(17273), 1-12.
- Marie, A.D., Herbinger, C., Fullsack, P. & Rico, C. (2019) First reconstruction of kinship in a Scalloped Hammerhead Shark aggregation reveals the mating patterns and breeding sex ratio. *Frontiers in Marine Science* 6:676. doi: 10.3389/fmars.2019.00676.
- Moyle, P.B. & Cech, J.J. (2004) *Fishes, an introduction to ichthyology*. Prentice Hall, Upper Saddle River, 726 p.
- Nelson, J.S., Terry, C., Grande, T. C. & Wilson, M.V.H. (2016) *Fishes of the World*. John Wiley and Sons, Hoboken, 752 pp.
- Quattro, J.M., Driggers, W.B., Grady, J.M., Ulrich, G.F. & Roberts, M.A. (2013) *Sphyrna gilberti* sp. nov., a new hammerhead shark (Carcharhiniformes, Sphymidae) from the western Atlantic Ocean. *Zootaxa*, 3702(2), 159-178.
- Rigby, C.L., Dulvy, N.K., Barreto, R., Carlson, J., Fernando, D., Fordham, S., Francis, M.P., Herman, K., Jabado, R.W., Liu, K.M., Marshall, A., Pacourea, N., Romanov, E., Sherley, R.B. & Winker, H. (2019) *Sphyrna lewini*. The IUCN Red List of Threatened Species 2019: e.T39385A2918526. Available from: <https://dx.doi.org/10.2305/IUCN.UK.2019-3.RLTS.T39386A2920499.en>. (08 April 2020).
- Salinas-de-León, P., Hoyos-Padilla, E.M. & Pochet, F. (2017) First observation on the mating behaviour of the endangered Scalloped Hammerhead Shark *Sphyrna lewini* in the Tropical Eastern Pacific. *Environmental Biology of Fishes*, 100, 1603-1608.
- Sentosa, A.S., Dharmadi, D. & Tjahjo, D.W.H. (2016) Parameter populasi hiu martil (*Sphyrna lewini* Griffith & Smith, 1834) di perairan selatan Nusa Tenggara. *Jurnal Penelitian Perikanan Indonesia*, 22(4), 253-262.
- Wahyudin, I., Kamal, M.M., Fahrudin, A. & Boer, M. (2019) Length-weight relationship and reproductive size of silky shark *Carcharhinus falciformis* and scalloped hammerhead shark *Sphyrna lewini* collected in Tanjung Luar fish landing port, East Lombok, Indonesia. *AACL Bioflux*, 12(1), 355-362.

First Record of Scalloped hammerhead shark *Sphyrna lewini* (Carchariniformes : Sphyrnidae) in freshwater habitat

ORIGINALITY REPORT

11 %

SIMILARITY INDEX

MATCH ALL SOURCES (ONLY SELECTED SOURCE PRINTED)

★www.rebibio.net 3%
Internet

EXCLUDE QUOTES ON
EXCLUDE ON
BIBLIOGRAPHY

EXCLUDE MATCHES < 1%

FORMAT PENILAIAN (VALIDASI & PEER REVIEW)
LEMBAR
HASIL PENILAIAN SEJAWAT SEBIDANG ATAU *PEER REVIEW*
KARYA ILMIAH : JURNAL ILMIAH

Jurnal Artikel Ilmiah : First record of scalloped hammerhead shark *Sphyraena lewini* (Carcharhiniformes: Sphyrnidae) in freshwater habitat

Penulis Artikel Ilmiah : Arum Setiawan

Identitas Jurnal Artikel Ilmiah : a. Nama Jurnal : Ecologica Montenegrina

b. Nomor/Volume/Hal : 1/30/113-118

c. Edisi (bulan/tahun) : April 2020

d. Penerbit : Center for Biodiversity of Montenegro

e. Jumlah Halaman 6

Kategori Publikasi Jurnal Ilmiah : Jurnal Ilmiah Internasional Bereputasi

(beri √ pada kategori yang tepat) Jurnal Ilmiah Internasional

Jurnal Ilmiah Nasional Terakreditasi S1, S2

Jurnal Ilmiah Nasional Terakreditasi S3, S4

Jurnal Ilmiah Nasional Tidak Terakreditasi

I. Hasil Penilaian Validasi :

No.	ASPEK	URAIAN/KOMENTAR PENILAIAN
1.	Indikasi Plagiasi	11 %
2.	Linearitas	Sudah linier dengan bidang biologi konservasi

II. Hasil Penilaian Peer Review :

Komponen Yang Dinilai	Nilai Maksimal Jurnal Ilmiah (isikan di kolom yang sesuai)					Nilai Akhir Yang Diperoleh
	Internasional Bereputasi (Maks 40)	Internasional (Maks 20)	Nasional Terakreditasi S1, S2 Maks 25	Nasional Terakreditasi S3, S4 Maks 20	Nasional tidak Terakreditasi (maks 10)	
Kelengkapan dan Kesesuaian unsur isi jurnal (10%)	4					4
Ruang lingkup dan kedalaman pembahasan (30%)	12					12
Kecukupan dan Kemutahiran data/informasi dan metodologi (30%)	12					12
Kelengkapan unsur dan kualitas penerbit (30%)	12					12
Total = (100%)	40					40
Kontribusi Pengusul (Penulis Pertama /Anggota Utama)	Anggota Utama $(0,4 \times 40) / 3 = 5,33$					5,33
KOMENTAR/ULASAN PEER REVIEW						
• Kelengkapan dan Kesesuaian Unsur:	Paper terkait deskripsi ikan shark <i>Sphyraena lewini</i> di perairan air tawar. Isi paper sudah memenuhi kaidah-kaidah karya ilmiah dan sudah sesuai dengan bidang biologi konservasi					
• Ruang Lingkup dan Kedalaman Pembahasan:	Hasil penelitian dibahas secara komprehensif dengan penyampaian pembanding dari temuan-temuan penelitian lainnya dan teori terkait. Referensi yang diajukan dalam pembahasan sudah cukup update untuk bidang kajian ini.					
• Kecukupan & Kemutakhir Data & Metodologi:	Data-data hasil penelitian sudah baik dan didukung peta lokasi sampling dan gambar yang ditampilkan menarik. Data didapatkan dengan menggunakan metode yang sudah standard.					
• Kelengkapan Unsur & Kualitas Penerbit:	Penerbit Center for Biodiversity of Montenegro berkualitas baik, tidak termasuk predatory publisher, dan jurnal masuk di Q2.					

Surabaya, 13 Mei 2020
Penilai 1



Prof. Hery Purnobasuki, M.Si., Ph.D.
NIP 196705071991021001
Unit Kerja : Jurusan Biologi FST Unair
Bidang Ilmu : Biologi
Jabatan/Pangkat : Guru Besar/ Pembina Utama Madya

FORMAT PENILAIAN (VALIDASI & PEER REVIEW)**LEMBAR****HASIL PENILAIAN SEJAWAT SEBIDANG ATAU PEER REVIEW****KARYA ILMIAH : JURNAL ILMIAH**

Jurnal Artikel Ilmiah

: First record of scalloped hammerhead shark *Sphyrna lewini*
 (Carcharhiniformes: Sphyrnidae) in freshwater habitat

Penulis Artikel Ilmiah

: Arum Setiawan

Identitas Jurnal Artikel Ilmiah

a. Nama Jurnal : Ecologica Montenegrina
 b. Nomor/Volume/Hal : 1/30/113-118
 c. Edisi (bulan/tahun) : April 2020
 d. Penerbit : Center for Biodiversity of Montenegro
 e. Jumlah Halaman : 6

Kategori Publikasi Jurnal Ilmiah
 (beri ✓ pada kategori yang tepat)

- : Jurnal Ilmiah Internasional Bereputasi
 Jurnal Ilmiah Internasional
 Jurnal Ilmiah Nasional Terakreditasi S1, S2
 Jurnal Ilmiah Nasional Terakreditasi S3, S4
 Jurnal Ilmiah Nasional Tidak Terakreditasi

I. Hasil Penilaian Validasi :

No.	ASPEK	URAIAN/KOMENTAR PENILAIAN
1.	Indikasi Plagiasi	11 %
2.	Linearitas	V

II. Hasil Penilaian Peer Review :

Komponen Yang Dinilai	Nilai Maksimal Jurnal Ilmiah (isikan di kolom yang sesuai)					Nilai Akhir Yang Diperoleh
	Internasional Bereputasi (Maks 40)	Internasional (Maks 20)	Nasional Terakreditasi S1, S2 Maks 25	Nasional Terakreditasi S3, S4 Maks 20	Nasional tidak Terakreditasi (maks 10)	
Kelengkapan dan Kesesuaian unsur isi jurnal (10%)	4					3
Ruang lingkup dan kedalaman pembahasan (30%)	12					11
Kecukupan dan Kemutahiran data/informasi dan metodologi (30%)	12					11
Kelengkapan unsur dan kualitas penerbit (30%)	12					12
Total = (100%)	40					37
Kontribusi Pengusul (Penulis Pertama /Anggota Utama)	Ecologica Montenegrina 30(1): 113-118. Index Factor: 0,79. Penulis ke 3 dari 4. Nilai maksimal %. Nilai pengusul: $(0,4 \times 0,925 \times 40)/3 = 4,9$					4,9

KOMENTAR/ULASAN PEER REVIEW

• Kelengkapan dan Kesesuaian Unsur:	Format lengkap.
• Ruang Lingkup dan Kedalaman Pembahasan:	Masih dalam ruang lingkup dan pembahasan terbatas.
• Kecukupan & Kemutahiran Data & Metodologi:	Data cukup memadai. Metode tidak ada yang baru, sudah terbiasa dilakukan peneliti pada umumnya.
• Kelengkapan Unsur & Kualitas Penerbit:	Cukup lengkap sebagai penerbit. Penerbit berkualitas.

Yogyakarta, 16 Juni 2020

Perfilai 2

tanda tangan ...

Prof. Dr. Suwarno Hadisusanto

NIP 195411161983031002

Unit Kerja : Fakultas Biologi UGM

Bidang Ilmu : Biologi/Ekologi

Jabatan/Pangkat : Guru Besar/ Pembina Utama Madya