

Toxicity of entomopathogenic fungal culture filtrate of lowland and highland soil of South Sumatra (Indonesia) against *Spodoptera litura* larvae

MIMMA GUSTIANINGTYAS¹, SITI HERLINDA^{1,2,3,*}, SUWANDI^{1,2,3}, SUPARMAN^{1,2}, HARMAN HAMIDSON^{1,2}, HASBI^{3,4}, ARUM SETIAWAN⁵, MARIESKA VERAWATY⁵, ELFITA⁶, ARSI^{2,3}

¹Program of Crop Sciences, Faculty of Agriculture, Universitas Sriwijaya. Palembang 30139, South Sumatera, Indonesia

²Department of Plant Pests and Diseases, Faculty of Agriculture, Universitas Sriwijaya, Indralaya 30662, South Sumatera, Indonesia.

Tel.: +62-711-580663, Fax.: +62-711-580276, *email: sitherlinda@unsri.ac.id

³Research Center for Sub-optimal Lands (PUR-PLSO), Universitas Sriwijaya. Palembang 30139, South Sumatera, Indonesia

⁴Department of Agricultural Engineering, Faculty of Agriculture, Universitas Sriwijaya. Indralaya, Ogan Ilir 30662, South Sumatera, Indonesia

⁵Department of Biology, Faculty of Mathematics and Natural Sciences, Universitas Sriwijaya. Indralaya, Ogan Ilir 30662, South Sumatera, Indonesia

⁶Department of Chemistry, Faculty of Mathematics and Natural Sciences, Universitas Sriwijaya. Indralaya, Ogan Ilir 30662, South Sumatera, Indonesia

Manuscript received: 6 March 2020. Revision accepted: 7 April 2020.

Abstract. *Gustianingtyas M, Herlinda S, Suwandi, Suparman, Hamidson H, Hasbi, Setiawan A, Verawaty M, Elfita, Arsi. 2020. Toxicity of entomopathogenic fungal culture filtrate of lowland and highland soil of South Sumatra against Spodoptera litura larvae. Biodiversitas 21: 1839-1849.* The use of secondary fungal metabolites for the active ingredient of mycoinsecticide is more effective and more easily integrated with other pest control techniques. This study aimed to measure the toxicity of the culture filtrate of entomopathogenic fungi originating from South Sumatra against the *Spodoptera litura* larvae. *Beauveria bassiana* (25 isolates) and *Metarhizium anisopliae* (20 isolates) of South Sumatra were cultured in liquid media and filtered to produce culture filtrate. The larvae which were sick due to the filtrate showed the symptoms of decreased appetite and were not actively moving, while the dead larvae were characterized by being wrinkled, dry, black integument, and odorless. Mortality caused by *B. bassiana* filtrate was the highest 98% (BJgTs isolate) and not significantly different from the BSwTd2 isolate (94.67%). Yet, the LT₅₀ BSwTd2 isolate was shorter (5.92 days) compared to the LT₅₀ BJgTs isolate (6.35 days). The most toxic *M. anisopliae* filtrate produced the mortality of 96% (MKbTp2 isolate) and 85.33% (MPdB isolate) each of which had LT₅₀ of 7.36 days and 8.09 days, respectively. So, the most toxic culture filtrate was BSwTd2 isolate of *B. bassiana* and MKbTp2 isolate of *M. anisopliae*. The entomopathogenic fungi producing filtrate which are toxic have the potential to be active ingredients of mycoinsecticides.

Keywords: *Beauveria bassiana*, chili, LT₅₀, *Metarhizium anisopliae*, mortality

INTRODUCTION

South Sumatra is generally dominated by lowland lands, namely freshwater swamps and tidal lowlands (Kartika et al. 2018; Karenina et al. 2019), and a small portion of medium and highland. Freshwater swamps can be found, among others in the Districts of Ogan Ilir, Ogan Komering Ilir, Musi Banyuasin, and Palembang City; the widest tide exists in Banyuasin District and the medium and highlands occur, among others, in the Districts of Ogan Komering Ulu Selatan, Lahat, and Pagaram City. The varied topographical conditions between districts/cities characterize the differences including the crops and microorganisms cultivated there.

In the lowlands of South Sumatra are generally cultivated annual crops, such as paddy (Herlinda et al. 2018a; Prabawati et al. 2019), maize (Juhriah et al. 2019), chili (Johari et al. 2016; Nasution and Respatijarti 2019; Sagrim et al. 2017), and palm oil (Darlan et al. 2016). Meanwhile, in the medium and high plains are generally cultivated annual crops such as coffee, tea (Zamhari et al. 2017), vegetables such as potatoes (Maryanto et al. 2018), cabbage, mustard greens, and caisim (Situmorang et al.

2019). Vegetables and seasonal plants of the lowlands and highlands generally have a major pest, namely *Spodoptera litura* (Turnip et al. 2019). The yield loss caused by its larval stage is more than 35% a year (Bueno et al. 2011). This pest has egg, larval, pupal, and adult stages (Vijaya and Rani 2016). *S. litura* is a polyphagous pest that attacks many vegetables in Indonesia, for example, chili (Nagal et al. 2016), cotton fruit (Maqsood et al. 2017), soybean (Fattah et al. 2018). A polyphagous pest is a pest that has a broad range of host species (more than a family) (Nagal et al. 2016). For the control of *S. litura*, it needs natural enemies such as entomopathogenic fungi that can adapt in low to high altitudes.

In the lowlands and highlands, the soil types have different chemical and physical characteristics; acidic pH soils are generally found in the lowlands and neutral or basic pH soils are found in the highlands (Munir and Herman 2019). Soil pH can affect the presence of entomopathogenic fungi and tends to be neutral or high pH soils which more commonly occur in entomopathogenic fungi than low pH soils (Safitri et al. 2018). Besides pH, the texture of sandy soils tends to be low and muddy which is not suitable for fungal propagule life as well as water-

saturated soil where fungi are rarely spotted to survive (Garrido-Jurado et al. 2011). Temperature (Pinnamaneni et al. 2010) and humidity (Oreste et al. 2015; Liu et al. 2017) also affect pathogenicity of entomopathogenic fungi.

Previous study found that more than 30 isolates of entomopathogenic fungi were found in the freshwater swamp, tidal, and highland soil of South Sumatra (Safitri et al. 2018). Apart from the soil, the entomopathogenic fungi were also found from the insect pests of vegetable and food crops of the lowlands to the highlands of South Sumatra (Herlinda et al. 2018b). Various origins of entomopathogenic fungi tend to produce pathogenicity variations (Sumikarsih et al. 2019).

The pathogenicity of entomopathogenic fungi is influenced by the ability of the fungi to produce secondary metabolites (Zibae et al. 2009). The fungi that develop in insect hemolymph or in liquid media are able to produce toxic metabolites, such as extracellular enzymes, proteins, and toxins (Bandani 2005). In vitro media, toxic metabolites can be produced by the fungi which are grown in liquid media and produce culture filtrate (Soesanto et al. 2019). The use of secondary entomopathogenic fungal metabolites for the active ingredient of mycoinsecticide is more effective and more easily integrated with other pest control techniques (Zibae et al. 2009). Entomopathogenic

fungal culture filtrate found in low to high altitude of South Sumatra potentially gets the most toxic isolates that can be utilized in the development of mycoinsecticide technology. For this reason, this study aimed to measure the toxicity of the culture filtrate isolates of entomopathogenic fungi from low to high land of South Sumatra, Indonesia against *Spodoptera litura* larvae.

MATERIALS AND METHODS

The experiments were carried out at the Entomology Laboratory, Department of Pests and Plant Diseases, Faculty of Agriculture, Universitas Sriwijaya from May to November 2018. The room temperatures and relative humidity during the experiment in the laboratory were 29.78°C and 82.72%, respectively. The entomopathogenic fungi isolates used in this study were explored by Herlinda et al. (2018) and Safitri et al. (2018), (Table 1 and 2) spread from the lowlands to the highlands of South Sumatra. The isolates were grouped into two: 25 isolates of *Beauveria bassiana* (Figure 1) and 20 isolates of *Metarhizium anisopliae* (Figure 2). All isolates were identified by Dr. Suwandi (a mycologist from Universitas Sriwijaya).




Figure 1. *Beauveria bassiana* isolates cultured in SDA media: BJgTs (A), BSsmMs (B), BSwTd1 (C), BSwTd2 (D), BSwTd3 (E), BSwTd4 (F), BPdR (G), BKbTp (H), BKKPp2 (I), Ts1d3 (J), BTmPc (K), BTmTr (L), (M) Ts1d2 (M), BTmTs (N), BlePd2 (O), BTmKt (P), BPCmS (Q), BMkMs (R), BtmGa (S), BTmSr (T), BPcPd2 (U), BtmSo (V), BtmMa (W), BtmPe (X), Blepd (Y)



Figure 2. *Metarhizium* isolates cultured in SDA media: MPdB (A), MPdR1 (B), MPdR2 (C), MpdPe (D), MJgMs1 (E), MJgMs2 (F), MPdMs1 (G), MPdMs2 (H), MPdMs3 (I), MjgKeTs (J), MPdMs4 (K), MJgTs2 (L), MKbTp1 (M), MSwTp1 (N), MSwTp2 (O), MSwTp3 (P), MSwTp4 (Q), MKKpp1 (R), MKbTp2 (S), MagPd (T)

Table 1. *Beauveria bassiana* isolates from South Sumatra used in this research

Isolate code	Isolate soil origin	Geographic origin
BJgTs	Tidal lowlands, maize	Telang Sari
BSmMs	Tidal lowlands, watermelon	Mulya Sari
BSwTd1	Peatlands, oil palm	Talang Dabok
BSwTd2	Peatlands, oil palm	Talang Dabok
BSwTd3	Peatlands, oil palm	Talang Dabok
BSwTd4	Peatlands, oil palm	Talang Dabok
BPdR	Freshwater swamps, paddy	Rambutan
BKbTp	Highlands, cabbage	Talang Patai
BKKpp2	Highlands, rubber, and coffee	Pulau Pinang
Ts1d3	Peatlands	Talang Dabok
BTmPc	Freshwater swamps	Indralaya
BTmTr	Freshwater swamps	Telang Rejo
Ts1d2	Peatlands	Talang Dabok
BTmTs	Highlands	Mulia Sari
BlePd2	<i>Lipaphis erysimi</i>	Pagardin
BTmkt	Freshwater swamps	Kenten
BPCmS	<i>Pseudoplusia chalcites</i>	Muara Siban
BMkMs	Highlands	Muara Siban
BtmGa	Freshwater swamps	Gandus
BTmSr	Tidal lowlands	Srikaton
BPcPd2	<i>Chrysodeixis chalcites</i>	Pagardin
BtmSo	Freshwater swamps	Soak
BtmMa	Freshwater swamps	Mariana
BtmPe	Freshwater swamps	Pemulutan
BlepD	<i>Lipaphis erysimi</i>	Pagardin

Table 2. *Metarhizium anisopliae* isolates from South Sumatra used in this research

Isolate code	Isolate soil origin	Geographic origin
MPdB	Freshwater swamps, paddy	Banyuasin
MPdR1	Freshwater swamps, paddy	Rambutan
MPdR2	Freshwater swamps, paddy	Rambutan
MpdPe	Freshwater swamps, paddy	Pemulutan
MJgMs1	Tidal lowlands, maize	Mulya Sari
MJgMs2	Tidal lowlands, maize	Mulya Sari
MPdMs1	Tidal lowlands, paddy	Mulya Sari
MPdMs2	Tidal lowlands, paddy	Mulya Sari
MPdMs3	Tidal lowlands, paddy	Mulya Sari
MjgKeTs	Tidal lowlands, maize, and oil palm	Telang Sari
MPdMs4	Tidal lowlands, paddy	Mulya Sari
MJgTs2	Tidal lowlands, maize	Telang Sari
MKbTp1	Highlands, cabbage	Talang Patai
MSwTp1	Highlands, mustard	Talang Patai
MSwTp2	Highlands, mustard	Talang Patai
MSwTp3	Highlands, mustard	Talang Patai
MSwTp4	Highlands, mustard	Talang Patai
MKKpp1	Highlands, mustard, rubber, coffee	Pulau Pinang
MKbTp2	Highlands, cabbage	Talang Patai
MagPd	Highlands, <i>Aphis gossypii</i>	Pagardin

Test insect preparation

S. litura larvae were collected from the chili planting area in the experimental field of the Faculty of Agriculture, Sriwijaya University and farmers' vegetable fields which were not applied by synthetic insecticides. Then, the larvae were brought to the laboratory and maintained in a plastic cylinder whose upper part was covered with gauze (Ø 25 cm, height 30 cm) having been washed thoroughly using detergent. Next, water spinach plants were put into it for larvae feed.

Every day the larvae feed was replaced with fresh new feed. The larvae entering the pupae phase were transferred into a plastic cylinder (Ø 10 cm, height 15 cm) with the top open and the bottom of the cylinder sprinkled with sifted soil and sterilized in an oven for 1 hour at 100°C. The thickness of the soil inserted into the cylinder was 3 cm. The plastic cylinder containing the pupae was put into a gauze cage (30x30x30 cm³) which had been chopped with chilies for laying eggs. Adults arising from the pupae were fed with honey smeared on cotton and hung over a cage. The hatched eggs were transferred into a plastic cylinder and fed with water spinach which was replaced every day. The larvae used for toxicity testing were the second instar of the third offspring or afterward.

Culture filtrate production

All isolates before being filtered were first made fit by modifying the method of Herlinda (2010). The media for making fit used Sabouraud Dextrose Agar (SDA, Merck) as much as 16.2 g and added 250 mL aquadest enriched with 1.25 g *Tenebrio molitor* larvae flour (5 g). After that, the ingredients were mixed and stirred evenly and then put in an Erlenmeyer (size 250 mL), sterilized in an autoclave for 120 minutes at a pressure of 1 atm. The media was poured into the Petri dish as much as 10 mL (Ø 9 cm) in aseptic laminar flow air conditions. Then, the entomopathogenic fungi isolates were inoculated in the SDA media and incubated for 14 days.

Isolates of entomopathogenic fungi derived from SDA media of aged 14 days were then transferred and grown into liquid media (broth media), PDB (*Potato Dextrose Broth*) prepared as follows: the PDB media composition consisting of 20 g dextrose monohydrate, 200 g potatoes, and 1000 mL aquadest. Before the potatoes were extracted, they were cut into cubes with a size of ± 2x2x2 cm³ and boiled using 1000 mL aquadest for 20 minutes. This PDB medium was sterilized in the autoclave and after it got cold, the entomopathogenic fungus isolates were inoculated into it. This liquid culture was incubated for 6 weeks.

After the fungus liquid culture (fungal broth) was 6 weeks old, the fungus filtration is carried out. The filtration was conducted in two stages, namely filtering using filter paper and syringe filter. In the first stage, 100 mL of 6-week-old fungal broth was filtered using Whatman filter paper no. 42 and coated with a thickness of 1 cm cotton. The culture filtrate obtained from the first stage of filtering was then sucked up to 10 mL using a hypodermic needle (spike). Then, the needle was removed and the base of the needle was fitted with a syringe filter (0.45 µm-25 mm). The 10 mL spike was pressed so that it released culture

filtrate from the syringe filter. The 100 mL of fungal broth produced ± 70 mL of culture filtrate. The culture filtrate was taken as much as 1 mL to be dripped on a piece of chili leaf which was used as a toxicity test. This culture filtrate is often referred to as raw secondary metabolites (Soesanto et al. 2019). To ensure the culture filtrate did not contain propagules (hyphae, mycelia) and conidia, it was grown on agar media (SDA) before it was applied. If it does not grow fungus on the SDA media, the culture filtrate can be applied.

Insecticidal activity test of entomopathogenic fungus culture filtrate

The chili leaves that had been dropped by the culture filtrate were first drained; before 25 *S. litura* larvae were fasted for 2 hours, and then put into them. The second instar larvae were left to eat leaves that had been dropped with culture filtrate for 6 hours. After 6 hours, the larvae were transferred into a plastic cylinder topped with gauze (Ø 25 cm, height 30 cm) containing 15-30 clean chili leaves which were not dripped with culture filtrate. Every day the chili leaves were replaced with the new ones. Every day the dead larvae were recorded up to 12 days after the application.

Data analysis

The differences in mortality data and lethal time that killed 50% of test insects (LT₅₀) were analyzed using analysis of variance (ANOVA), after that further tests were carried out using the 5% Tukey's Honestly Significant Difference (HSD). All data were calculated using software of SAS University Edition 2.7 9.4 M5.

RESULTS AND DISCUSSION

Insecticidal activities of entomopathogenic fungi

The culture filtrate derived from twenty-five *B. bassiana* isolates tested for the insecticidal activity showed that the most toxic *B. bassiana* isolates were those coded BJgTs (98%) and were not significantly different, among others with BSwTd2 (94.67%) and BTmTs isolates (96%). All *B. bassiana* isolates showed the ability to kill the test larvae (Table 3). The BJgTs, BSwTd2, and BTmTs isolates were the most toxic because they produced the highest mortality and also their ability to kill the shortest as evidenced by the short LT₅₀ (6.35, 5.92, and 7.78 days) isolates. The most toxic isolates were shown with the highest mortality and their shortest LT₅₀. Consequently, for *B. bassiana*, the most toxic isolate was BSwTd2.

The culture filtrate of the twenty *M. anisopliae* isolates which were tested for their insecticidal activity showed that the most toxic isolates were that coded MKbTp2 (96%) and MPdB (85.33%) and the mortality of both isolates was significantly different from the other isolates (Table 4). The BSwTd2 isolate was the most toxic because they produced the highest mortality as well as the ability to kill the shortest as evidenced by the shortest LT₅₀ (7.36 days), and then followed by LT₅₀ isolate MPdB (8.09 days). The most

toxic *M. anisopliae* isolate was proven by the highest mortality and the shortest LT₅₀ which was MKbTp2 isolate.

The color of the fungal broth (Figure 3) and culture filtrate (Figure 4) of *B. bassiana* varied among the isolates. All fungal broth isolates were generally transplanted brown, but there were among those isolates dark brown in color, for example, isolates BSwTd2, BSwTd3, BPdR, TS1d2. There was a tendency for the darker color of the fungal broth and culture filtrate to cause higher mortality. The color of fungal broth (Figure 5) and culture filtrate (Figure 6) of *M. anisopliae* differed from the colors of fungal broth and culture of *B. bassiana*. The *M. anisopliae* fungal broth was generally dark greenish-brown, but the culture filtrate tended to have a more varied color, from greenish dark brown to light brown. The same tendency was also produced in *M. anisopliae* isolates, i.e., the color of the older culture filtrate resulting in higher mortality. The isolates having a darker colored culture filtrate were MPdB, MPdMs1, MPdMs2, MPdMs3, MKbTp1, MKbTp2.

Effect of entomopathogenic fungal culture filtrate on test larvae

Spodoptera litura larvae fed with chili leaves applied with *B. bassiana* or *M. anisopliae* culture filtrate exhibited the same behavior, i.e., they increasingly ate less, while the controls not given culture filtrate continued to eat greedily. The chili leaves given to the larvae which were not given culture filtrate generally only left a leaf bone, whereas in the larvae given culture filtrate there were still leaves remaining (Figures 7 and 8). Thus, the culture filtrate could reduce the appetite of *S. litura* larvae.

The larvae *S. litura* has given culture filtrate besides being lazier to eat showed lazier and lazier to move. After 2 and 3 days being given the culture filtrate, the larvae began to change the color of integument which was previously bright green to become dull and the larvae were discovered to be dead. Four days after being given the culture filtrate, the larvae increasingly began to shrink and die. The dead larvae were wrinkled, dry, black, and odorless. The larvae not given the culture filtrate had larger body sizes and the integument colors were greener, and brighter (Figures 9 and 10).

Table 3. Mortality of larval *Spodoptera litura* after being infested with *Beauveria bassiana* culture filtrates

Isolate codes	Mortality ± SE*(%)	LT ₅₀ (days)± SE
BJgTs	98.00 ± 1.41 ^b	6.35 ± 0.49
BSmMs	50.00 ± 32.53 ^{ab}	11.57 ± 4.08
BSwTd1	49.33 ± 15.36 ^{ab}	12.03 ± 3.52
BSwTd2	94.67 ± 4.35 ^{ab}	5.92 ± 0.46
BSwTd3	73.33 ± 18.51 ^{ab}	8.86 ± 1.78
BSwTd4	26.00 ± 1.41 ^{ab}	16.07 ± 0.97
BPdR	86.67 ± 3.93 ^{ab}	8.17 ± 0.61
BKbTp	62.67 ± 10.39 ^{ab}	10.71 ± 0.91
BKKPp2	58.00 ± 7.07 ^{ab}	10.68 ± 0.34
TS1d3	10.67 ± 2.88 ^a	17.45 ± 0.76
BTmPc	46.67 ± 17.01 ^{ab}	14.32 ± 3.72
BTmTr	29.33 ± 9.68 ^{ab}	14.31 ± 1.79
TS1d2	88.00 ± 8.22 ^{ab}	7.87 ± 1.11
BTmTs	96.00 ± 0.00 ^{ab}	7.78 ± 0.09
BLePd2	70.00 ± 21.21 ^{ab}	9.59 ± 1.43
BTmKt	80.00 ± 14.73 ^{ab}	8.80 ± 1.41
BPcMs	81.33 ± 9.49 ^{ab}	7.87 ± 1.32
BMkMs	72.00 ± 17.99 ^{ab}	10.82 ± 2.46
BTmGa	66.67 ± 22.34 ^{ab}	12.27 ± 3.60
BTmSr	21.33 ± 9.30 ^{ab}	16.80 ± 2.67
BPcPd2	18.67 ± 7.85 ^{ab}	9.43 ± 3.86
BTmSo	80.00 ± 14.14 ^{ab}	7.55 ± 1.35
BTmMa	52.00 ± 19.69 ^{ab}	11.09 ± 1.62
BTmPe	70.00 ± 18.38 ^{ab}	9.30 ± 1.94
BLePd	94.00 ± 4.24 ^{ab}	9.30 ± 0.45
ANOVA F-value	2.09 [*]	1.20 ^{ns}
P value (0.05)	0.02	0.30
Tukey's HSD test	67.09	-

Note: ns = not significantly different; * = significantly different; values within a column (the data of each isolate) followed by the same letters were not significantly different at P < 0.05 according to Tukey's HSD test. Original data were transformed using Arcsin transformation prior to statistical analysis

Table 4. Mortality of larval *Spodoptera litura* after being infested with *Metarhizium anisopliae* culture filtrates

Isolate codes	Mortality ± SE*(%)	LT ₅₀ (days)± SE
MPdB	85.33 ± 14.67 ^b	8.09 ± 1.88 ^a
MPdR1	57.33 ± 19.37 ^{ab}	11.87 ± 2.56 ^a
MPdR2	65.33 ± 13.33 ^{ab}	10.43 ± 0.51 ^a
MPdPe	42.67 ± 9.61 ^{ab}	12.67 ± 0.89 ^a
MJgMs1	53.33 ± 23.25 ^{ab}	13.67 ± 4.52 ^a
MJgMs2	25.33 ± 3.53 ^{ab}	14.50 ± 0.56 ^a
MPdMs1	73.33 ± 10.41 ^{ab}	9.90 ± 0.97 ^a
MPdMs2	84.00 ± 9.24 ^{ab}	8.70 ± 1.26 ^a
MPdMs3	68.00 ± 16.17 ^{ab}	10.08 ± 0.86 ^a
MJgKeTs	42.67 ± 10.67 ^{ab}	12.68 ± 1.05 ^a
MPdMs4	16.00 ± 10.07 ^{ab}	53.70 ± 23.11 ^b
MJgTs2	46.67 ± 13.92 ^{ab}	14.29 ± 2.23 ^a
MKbTp1	84.00 ± 10.07 ^{ab}	8.69 ± 0.95 ^a
MSwTp1	56.00 ± 12.22 ^{ab}	14.64 ± 3.24 ^a
MSwTp2	74.67 ± 13.92 ^{ab}	10.62 ± 1.39 ^a
MSwTp3	61.33 ± 16.38 ^{ab}	10.91 ± 1.99 ^a
MSwTp4	61.33 ± 23.13 ^{ab}	14.60 ± 5.54 ^a
MKKPp1	44.00 ± 10.07 ^{ab}	12.33 ± 0.90 ^a
MKbTp2	96.00 ± 4.00 ^b	7.36 ± 0.62 ^a
MagPd	33.33 ± 3.53 ^{ab}	12.98 ± 0.23 ^a
ANOVA F-value	2.39 [*]	50.13 [*]
P value (0.05)	0.01	0.00
Tukey's HSD test	53.35	2.65

Note: ns = not significantly different; * = significantly different; values within a column (the data of each isolate) followed by the same letters were not significantly different at P < 0.05 according to Tukey's HSD test. Original data were transformed using Arcsin transformation prior to statistical analysis




Figure 3. *Beauveria bassiana* isolates cultured in PDB: BJgTs (A), BSmMs (B), BSwTd1 (C), BSwTd2 (D), BSwTd3 (E), BSwTd4 (F), BPdR (G), BKbTp (H), BKKPp2 (I), Ts1d3 (J), BTmPc (K), BTmTr (L), (M) Ts1d2 (M), BTmTs (N), BlePd2 (O), BTmKt (P), BPCmS (Q), BMkMs (R), BtmGa (S), BTmSr (T), BPcPd2 (U), BtmSo (V), BtmMa (W), BtmPe (X), Blepd (Y)

Discussion

The *B. bassiana* culture filtrate, particularly the isolates coded BSwTd2, were the most toxic compared to other isolates with LT_{50} of only 5.91 days. The BSwTd2 isolate was from peat soils overgrown with oil palm plants. For *M. anisopliae*, the most toxic MKbTp2 isolate was from the cabbage in highland. The larvae began to die after 3x24 hours after the treatment either due to *B. bassiana* or *M. anisopliae*. The larvae death resulted from the 6 weeks of incubation of entomopathogenic fungi causing the toxic culture filtrate. According to Pinnamaneni et al. (2010), the culture of filtrates of the toxic *B. bassiana* was caused by the incubation in liquid culture media in which the fungus produced chitinolytic exochitinase enzymes so that when they entered the body the insect was able to degrade the cuticle. In addition, the fungal broth during the incubation could produce protease enzymes (Qazi 2008).

The time of death of the larvae by this culture filtrate was faster than that of by conidia. El Husseini (2019) stated that the death of *Spodoptera* larvae by *M. anisopliae* conidia began to occur at the fourth day (4x24 hours) or the fifth day after the treatment (post-treatment), whereas in this study the *M. anisopliae* culture filtrate began to kill *S. litura* at the third day after the treatment. The lethal time difference between these was due to the mode of action of fungal conidia being different from the culture filtrate. To kill the host insect, the fungus conidia got contact first with it and then produced toxins to kill the host, while the culture filtrate directly killed the host insect. El-Ghany (2015) states that entomopathogenic fungus conidia kill its host insect in two stages: parasitic and saprophytic phases. The parasitic phase began with the fungal conidia attaching to the host insect cuticle (Augustyniuk-Kram and Kram 2012). Then, if the humidity was high, the conidia started to germinate on the host cuticle (El-Ghany 2015). The

conidia germinating by forming germ tubes continued to grow sticking out to find a soft integument position to facilitate entry into the cuticle (Fernandes et al. 2007). The infection occurs when the germ tubes are able to penetrate insect cuticles and the ability to infect them is a determining factor for the fungus virulence (Altre and Vandenberg 2001). After the germ, tubes penetrated the cuticle and reached the hemocoel, and then they produced specific infection hyphae originating at appressoria (El-Ghany 2015). Furthermore, the hyphae spread to the hemolymph and developed to produce blastospores, and the blastospores produced fungal cytotoxic, for example, destruxins by *M. anisopliae* which killed the host insects

(Mancillas-Paredes et al. 2019), but the death of these host insects was not caused only by the toxin but also due to the mechanical damage by the penetration of fungi into the body of insects (El-Ghany 2015).

After the host insect died, it entered into the saprophytic phase which was influenced by favorable environmental conditions (Peña-Peña et al. 2015). In the body of the dead insect, the fungus formed mycelia and hyphae which continued to grow covering the body of the host insect, and then the hyphae formed conidiogenous cells and the conidia were produced by utilizing the nutrients/fluids of the host insect and finally, the infection process was complete (El-Ghany 2015).



Figure 4. *Beauveria bassiana* culture filtrate: BJgTs (A), BSmmMs (B), BSwTd1 (C), BSwTd2 (D), BSwTd3 (E), BSpdR (F), BSpdR (G), BKbTp (H), BKKPp2 (I), Ts1d3 (J), BTmPc (K), BTmTr (L), (M) Ts1d2 (M), BTmTs (N), BlePd2 (O), BTmKt (P), BPCmS (Q), BMkMs (R), BtmGa (S), BTmSr (T), BPCPd2 (U), BtmSo (V), BtmMa (W), BtmPe (X), Blepd (Y)

In contrast to the mode of action of fungal conidia, a fungal culture filtrate directly kills the host insect due to the toxins produced by fungal broth so that the time needed to kill the host insect is shorter. The entomopathogenic fungus cultured in liquid media can produce conidia in the form of blastospores (Mascarin et al. 2015) which can produce toxins (Mascarin and Jaronski 2016). In this study, the result of calculating conidia density after being incubated for 6 weeks was generally more than 1×10^9 conidia mL^{-1} . Fungal conidia and propagules were separated from the filtrate through two stages of filtration to produce toxic fungal culture filtrate. The *B. bassiana* and *M. anisopliae* culture filtrates were able to kill up to 94.67% and 96% of the larvae of *S. litura*, respectively. The short time of death by this culture filtrate resulted from the death of the direct host insect by consuming feed (chili leaves) moistened with the culture filtrate. Soesanto et al. (2019) stated that entomopathogenic fungal culture filtrate contains secondary metabolites. *B. bassiana* culture filtrate

containing the secondary metabolites can weaken the host insect's immune system (Zibae et al. 2011). The secondary metabolites produce toxins, for example, destruxins and efrapeptins (Zibae et al. 2009). *B. bassiana* also produces a protease enzyme that can kill host insects by dissolving their body proteins (Mancillas-Paredes et al. 2019).

The color of fungal broth and culture of *B. bassiana* and *M. anisopliae* varied between isolates. The darker color of fungal broths and culture filtrates tend to cause higher mortality. In line with the results obtained by Ayudya et al. (2019), the darker culture filtrates tended to be more toxic than the light-colored ones. The darker culture filtrates show higher production of secondary metabolites (Luo et al. 2017) and the activity of extracellular enzymes (Khachatourians et al. 2007) and the enzymes produced such as proteases are able to dissolve integument insects that result in death (Mancillas-Paredes et al. 2019).




Figure 5. *Metarhizium anisopliae* isolates cultured in PDB: MPdB (A), MPdR1 (B), MPdR2 (C), MpdPe (D), MJgMs1 (E), MJgMs2 (F), MPdMs1 (G), MPdMs2 (H), MPdMs3 (I), MjgKeTs (J), MPdMs4 (K), MJgTs2 (L), MKbTp1 (M), MSwTp1 (N), MSwTp2 (O), MSwTp3 (P), MSwTp4 (Q), MKKPp1 (R), MKbTp2 (S), MagPd (T)




Figure 6. *Metarhizium anisopliae* culture filtrate: MPdB (A), MPdR1 (B), MPdR2 (C), MpdPe (D), MJgMs1 (E), MJgMs2 (F), MPdMs1 (G), MPdMs2 (H), MPdMs3 (I), MjgKeTs (J), MPdMs4 (K), MJgTs2 (L), MKbTp1 (M), MSwTp1 (N), MSwTp2 (O), MSwTp3 (P), MSwTp4 (Q), MKKp1 (R), MKbTp2 (S), MagPd (T)




Figure 7. The leaf damage: control (A) and larval *Spodoptera litura* applied with *Beauveria bassiana* culture filtrate (B)



Figure 8. The leaf damage: control (A) and larval *Spodoptera litura* applied with *Metarhizium anisopliae* culture filtrate (B)



Figure 9. The healthy (A) and dead larvae (B) of *Spodoptera litura* caused by *Beauveria bassiana* culture filtrate




Figure 10. The healthy (A) and dead larvae (B) of *Spodoptera litura* caused by *Metarhizium anisopliae* culture filtrate

The symptoms of sick host insects by culture filtrate in this study differed from the deaths by fungal conidia. The insects getting sick by the culture filtrate were not covered by mycelia or conidia on the integument. However, they got sick because of the fungal conidia infection which was generally covered in mycelia (Sumikarsih et al. 2019). The body shape of the sick *S. litura* larvae due to the culture filtrate was shrunk, wrinkled, dry, and odorless. Similar to the results of the study of Ayudya et al. (2019), the *S. litura* larvae treated with *B. bassiana* culture filtrate decreased their body weight and became dried, odorless, and the integument was not overgrown with fungal mycelia because in the culture filtrate it no longer contained conidia but contained toxic compounds for host insects. Therefore, the mortality of *S. litura* larvae in this study was caused by the culture filtrate containing toxic compounds. Zibae et al. (2011) state that toxic compounds in the culture filtrate are secondary metabolites. Zibae et al. (2009) stated that the secondary metabolites are easier to apply in the field because they can be integrated with other control techniques including synthetic insecticides. Consequently, the culture filtrate which is toxic due to the high content of secondary metabolites has the potential to be further developed for the active ingredient of mycoinsecticide.

The most toxic isolates of the culture filtrate were BSwTd2 of *B. bassiana* and MKbTp2 of *M. anisopliae* resulting in the mortality of above 90%. The two isolates that were found each came from the lowlands and highlands so that in the future there is an opportunity for the isolates to be developed and applied in low and highland ecosystems. Entomopathogenic fungi that produce toxic filtrate cultures are capable of killing their hosts in a short time, so that they have the potential to be active ingredients in the development of mycoinsecticides technology.

ACKNOWLEDGEMENTS

Thank you to the sponsors of the donors of this research, the Individual National Strategic Research (PSNI) program for the 2018 fiscal year based on the Research Contracts of the Directorate of Research and Community Service (DRPM), Directorate General of Research and Development Strengthening, Ministry of Technology Research and Higher Education, Contract number: 093/SP2H/LT/DRPM/IV/2018 chaired by SH.

REFERENCES

- Altre JA, Vandenberg JD. 2001. Factors influencing the infectivity of isolates of *Paecilomyces fumosoroseus* against diamondback moth, *Plutella xylostella*. *J Invertebr Pathol* 78: 31-36. DOI: 10.1006/jipa.2001.5043
- Augustyniuk-Kram A, Kram KJ. 2012. Entomopathogenic fungi as an important natural regulator of insect outbreaks in forests review. *Forest Ecosystem, More than Just Trees*. In: Blanco JA. (ed.). In Tech, Croatia. DOI: 10.5772/30596.
- Ayudya D, Herlinda S, Suwandi S. 2019. Insecticidal activity of culture filtrates from liquid medium of *Beauveria bassiana* isolates from South Sumatra (Indonesia) wetland soil against larvae of *Spodoptera litura*. *Biodiversitas* 20: 2101-2109. DOI: 10.13057/biodiv/d200802
- Bandani AR, Khambay BPS, Faull L, Newton R, Deadman M, Butt TM. 2000. Production of efrapeptins by *Tolypocladium* species and evaluation of their insecticidal and antimicrobial properties. *Mycol Res* 104: 537-544. DOI: 10.1017/S0953756299001859.
- Bueno RC, Bueno AF, Moscardi F, Parra JR, Hoffmanncampo CB. 2011. Lepidopteran larva consumption of soybean foliage: basis for developing multiple-species economic thresholds for pest management decisions. *Pest Manag Sci* 67 (2): 170-174.
- Darlan N, Pradiko I, Winarna, Siregar HH. 2016. Effect of el niño 2015 on oil palm performance in central and Southern Sumatera. *J Tanah dan Iklim* 40: 113-120. DOI: 10.2017/jti.v40n2.2016.113-120. [Indonesian]
- El-Ghany TMA. 2015. Entomopathogenic Fungi and their Role in Biological Control. Biology Department Faculty of Science Jazan University KSA, Cairo. DOI:10.4172/978-1-63278-065-2-66
- El-Husseini MMM. 2019. Efficacy of the entomopathogenic fungus, *Metarhizium anisopliae* (Metsch.), against larvae of the cotton leafworm, *Spodoptera littoralis* (Boisd.) (Lepidoptera: Noctuidae), under laboratory conditions. *Egypt J Biol Pest Control* 29: 4-6. DOI: 10.1186/s41938-019-0156-2
- Fattah A, Syam S, Daud ID, Dewi VS, Rahman A. 2018. The intensity of leaf damage caused by attack of *Spodoptera litura* F and seed yield on some soybean varieties in South Sulawesi Indonesia. *Sci Res J* 6: 55-60.
- Fernandes EKK, Rangel DEN, Moraes AM, Bittencourt VREP, Roberts DW. 2007. Variability in tolerance to UV-B radiation among *Beauveria* spp. isolates. *J Invertebr Pathol* 96: 237-243. DOI: 10.1016/j.jip.2007.05.007
- Garrido-Jurado J, Torrent J, Barrón V, Corpas A, Quesada-Moraga E. 2011. Soil properties affect the availability, movement, and virulence of entomopathogenic fungi conidia against puparia of *Ceratitis capitata* (Diptera: Tephritidae). *Biol Control* 58: 277-285. DOI: 10.1016/j.biocontrol.2011.05.017
- Herlinda S, Yudha S, Thalib R, Khodijah, Suwandi, Lakitan B, Verawaty M. 2018a. Species richness and abundance of spiders inhabiting rice in fresh swamps and tidal lowlands in South Sumatera, Indonesia. *J ISSAAS* 24: 82-93.
- Herlinda S, Rizkie L, Suwandi, Susilawati, Lakitan B, Verawaty M, Hasbi. 2018b. Effects of high temperature and ultraviolet-C irradiance on conidial viability and density of *Beauveria bassiana* and *Metarhizium anisopliae* isolated from soils of lowland ecosystems in Indonesia. *Eurasian J Anal Chem* 13: 209-216.
- Johari A, Herlinda S, Pujiastuti Y, Irsan C, Sartiami, D. 2016. Morphological and genetic variation of *Thrips parvispinus* (Thysanoptera: Thripidae) in chili plantation (*Capsicum annum* L.) in the lowland and highland of Jambi Province, Indonesia. *Am J Biol Sci* 2: 17-21. DOI: 10.11648/j.ajbio.s.2014020601.14
- Juhriah, Azrai M, Tambaru E, Rahayu JE. 2019. Phenotypic characteristics and grouping of hybrid waxy corn *Zea mays* L. *J Ilmu Alam dan Lingkungan* 10: 51-60. DOI: 10.20956/jal.v10i1.6624 [Indonesian]
- Karenina T, Herlinda S, Irsan C, Pujiastuti Y. 2019. Abundance and species diversity of predatory arthropods inhabiting rice of refuge habitats and synthetic insecticide application in freshwater swamps in South Sumatra, Indonesia. *Biodiversitas* 20 (8): 2375-2387. DOI: 10.13057/biodiv/d200836
- Kartika K, Lakitan B, Wijaya A, Kadir S, Widuri L.I, Siaga E. 2018. Effects of particle size and application rate of rice-husk biochar on chemical properties of tropical wetland soil, rice growth and yield. *Aust J Crop Sci* 12: 817-826. DOI: 10.21475/ajcs.18.12.05.PNE1043
- Khachatourians GG, Qazi SS, Reid WR. 2007. Biochemistry and molecular biology of entomopathogenic fungi. In: Brakhage AA, Zipfer PF (eds.). *The Mycota VI: Human and animal relationships*. Springer-Verlag, Berlin.
- Liu SF, Wang GJ, Nong XQ, Liu B, Wang M, Li SL, Cao GC, Zhang ZH. 2017. Entomopathogen *Metarhizium anisopliae* promotes the early development of Peanut Root. *Plant Protect Sci* 53: 101-107. DOI: 10.17221/49/2016-PPS
- Luo Z, Ren H, Mousa JJ, Rangel DEN, Zhang Y, Bruner SD, Keyhani NO. 2017. The PacC transcription factor regulates secondary metabolite production and stress response, but has only minor effects on virulence in the insect pathogenic fungus *Beauveria bassiana*. *Environ Microbiol* 19: 788-802. DOI: 10.1111/1462-2920.13648
- Mancillas-Paredes JM, Hernández-Sánchez H, Jaramillo-Flores ME, García-Gutiérrez C. 2019. Proteases and chitinases induced in

- Beauveria bassiana* during infection by *Zabrotes subfasciatus*. Southwestern Entomol 44: 125-137. DOI: 10.3958/059.044.0114
- Maqsood S, Sabri, MA, Ali A, Abbas M, Aziz A. 2017. Comparative toxicity of some insecticides against army worm, *Spodoptera litura* L. (Lepidoptera: Noctuidae) under laboratory conditions. J Entomol Zool Stud 5: 770-773.
- Maryanto M, Sukiyono K, Priyono B. 2018. Technical efficiency analysis and determinants in potato farming (*Solanum tuberosum* L.) in Pagar Alam City, South Sumatra Province. Agraris 4: 1-8. DOI: 10.18196/agr.4154 [Indonesian]
- Mascarin GM, Jackson MA, Behle RW, Kobori NN, Júnior ÍD. 2016. Improved shelf life of dried *Beauveria bassiana* blastospores using convective drying and active packaging processes. Appl Microbiol Biotechnol 100: 8359-8370. DOI: 10.1007/s00253-016-7597-2
- Mascarin GM, Jackson MA, Kobori NN, Behle RW, Delalibera Júnior Í. 2015. Liquid culture fermentation for rapid production of desiccation-tolerant blastospores of *Beauveria bassiana* and *Isaria fumosorosea* strains. J Invertebr Pathol 127: 11-20. DOI: 10.1016/j.jip.2014.12.001
- Munir J, Herman W. 2019. Various physical and soil chemical properties phenomena support food security in West Sumatera. Ziraa'ah 44: 146-153. DOI: 10.31602/zmip.v44i2.1736. [Indonesia]
- Nagal G, Verma K, Rathore L. 2016. Management of *Spodoptera litura* (Fabricius) through some novel insecticides and biopesticides on bell pepper under polyhouse environment. Adv Life Sci 5: 1081-1084.
- Nasution K, Respatijarti. 2019. Yield trial on eight potential lines F7 generation of chili pepper (*Capsicum Annuum* L.) compact type in Low Land. J Produksi Tanaman 7: 464-473. [Indonesian]
- Oreste M, Baser N, Bubici G, Tarasco E. 2015. Effect of *Beauveria bassiana* strains on the *Ceratitis capitata* - *Psytalia concolor* system. Bull Insectol 68: 265-272.
- Peña-Peña AJ, Santillán-Galicia MT, Hernández-López J, Guzmán-Franco AW. 2015. *Metarhizium pingshaense* applied as a seed treatment induces fungal infection in larvae of the white grub *Anomala cincta*. J Invertebr Pathol 130: 9-12. DOI: 10.1016/j.jip.2015.06.010
- Pinnamaneni R., Kalidas P, Rao KRSS. 2010. Cloning and Expression of Bbchit1 gene of *Beauveria bassiana*. Open Entomol J 4: 30-35. DOI: 10.2174/1874407901004010030
- Prabawati G, Herlinda S, Pujiastuti Y. 2019. The abundance of canopy arthropods in South Sumatra (Indonesia) freshwater swamp main and ratooned rice applied with bioinsecticides and synthetic insecticide. Biodiversitas 20: 2921-2930. DOI: 10.13057/biodiv/d201021
- Qazi SS, Khachatourians GG. 2008. Addition of exogenous carbon and nitrogen sources to aphid exuviae modulates synthesis of proteases and chitinase by germinating conidia of *Beauveria bassiana*. Arch Microbiol 189 (6): 589-596.
- Safitri A, Herlinda S, Setiawan, A. 2018. Entomopathogenic fungi of soils of freshwater swamps, tidal lowlands, peatlands, and highlands of South Sumatra, Indonesia. Biodiversitas 19: 2365-2373. DOI: 10.13057/biodiv/d190647
- Sagrim M, Sumule AI, Iyai DA, Baransano M. 2017. Prime potency of agriculture commodities on highland of Arfak Mountains Regency, Papua Barat. J Ilmu Pertanian Indonesia 22: 141-146. [Indonesian]. DOI: 10.18343/jipi.22.3.141
- Situmorang S, Haryono D, Soelaiman A. 2019. Analysis of business performance and welfare of vegetable farmers in Pagar Alam City South Sumatera Province. Indon J Socio Econ 1: 84-92. [Indonesian]
- Soesanto L, Mugiastuti E, Manan A. 2019. Raw secondary metabolites application of two *Trichoderma harzianum* isolates towards vascular streak dieback on cocoa seedlings. Pelita Perkebunan 35: 22-32. DOI: 10.22302/iccri.jur.pelitaperkebunan.v35i1.346
- Sumikarsih E, Herlinda S, Pujiastuti Y. 2019. Conidial density and viability of *Beauveria bassiana* isolate from java and Sumatra and their virulence against *Nilaparvata lugens* at different temperatures. Agrivita 1: 335-349. DOI: 10.17503/agrivita.v41i2.2105
- Turnip A, Runtuboi DYP, Lantang D. 2018. Effectivity test of *Beauveria bassiana* fungus and application time of *Spodoptera litura* in mustard greens (*Brassica juncea*). J Biologi Papua 10: 26-31. DOI: 10.31957/jbp.131. [Indonesian]
- Vijaya M, Rani PU. 2017. Defensive responses in *Capsicum annum* (L) plants, induced due to the feeding by different larval instars of *Spodoptera litura* (F). Arthropod-Plant Interact 11: 193-202.
- Zamhari A, Sitorus S, Pravitasari A. 2017. Analysis of the competitive commodities and its direction of development plan in Pagar Alam City, South Sumatera Province. Tata Loka 19: 218-229. DOI: 10.14710/tataloka.19.3.218-229. [Indonesian]
- Zibae A, Bandani AR, Talaei-Hassanlouei R, Malagoli D. 2011. Cellular Immune Reactions of the Sunn Pest, *Eurygaster integriceps*, to the Entomopathogenic Fungus, *Beauveria bassiana* and its secondary metabolites. J Insect Sci 11: 1-16. DOI: 10.1673/031.011.13801
- Zibae A, Bandani AR, Tork M. 2009. Effect of the entomopathogenic fungus, *Beauveria bassiana*, and its secondary metabolite on detoxifying enzyme activities and acetylcholinesterase (AChE) of the Sunn pest, *Eurygaster integriceps* (Heteroptera: Scutellaridae). Biocontrol Sci Technol 19: 485-498. DOI: 10.1080/09583150902847127.



[i](#) [x](#)

Call for Papers

Submit your research to the newest title from APS, Function.

academic.oup.com OPEN

Biodiversitas

8









H Index

Country	Indonesia - SIR Ranking of Indonesia
Subject Area and Category	Agricultural and Biological Sciences Animal Science and Zoology Plant Science Biochemistry, Genetics and Molecular Biology Molecular Biology
Publisher	Biology department, Sebelas Maret University Surakarta
Publication type	Journals
ISSN	1412033X, 20854722
Coverage	2014-ongoing
Scope	"Biodiversitas, Journal of Biological Diversity" or Biodiversitas encourages submission of manuscripts dealing with all biodiversit aspects of plants, animals and microbes at the level of gene, species, and ecosystem.
	Homepage How to publish in this journal Contact Join the conversation about this journal

Publish Article in USA Journal

Journal of American Academic Research is an academic Journal in the United States.

american-journals.com OPEN



Show this widget in your own website

Just copy the code below and paste within your html code:



F **Fitra Syawal Harahap** 2 weeks ago

I want to submit my manuscript in this journal

reply

SCImago Team



Melanie Ortiz 2 weeks ago

Dear Fitra, thank you very much for your comment, we suggest you to look for author's instructions/submission guidelines in the journal's website. Best Regards, SCImago Team

J **Joko Prasetyo** 2 months ago

I want to submit my manuscript in this journal

reply

SCImago Team



Melanie Ortiz 2 months ago

Dear Joko, thank you very much for your comment, we suggest you to look for author's instructions/submission guidelines in the journal's website. Best Regards, SCImago Team

S **salamiah** 3 months ago

I see the subject area

reply

A **Afik** 7 months ago

Bagus kali jurnal nya

reply



Melanie Ortiz 7 months ago

Dear user, thanks for your participation! Best Regards, SCImago Team

F **Feron** 7 months ago

Wow bagus banget...samangat berkarya untuk Indonesia yang kebih maju

reply



Melanie Ortiz 7 months ago

Dear user, thanks for your participation! Best Regards, SCImago Team



muhamad iksan 2 years ago

i want submit my journal thanks

reply

A **ahmad** 12 months ago

Kindly submit your paper here, <https://smujo.id/biodiv/about/submissions>

Leave a comment

Name

Email

(will not be published)

I'm not a robot

reCAPTCHA
Privacy - Terms

Submit

The users of Scimago Journal & Country Rank have the possibility to dialogue through comments linked to a specific journal. The purpose is to have a forum in which general doubts about the processes of publication in the journal, experiences and other issues derived from the publication of papers are resolved. For topics on particular articles, maintain the dialogue through the usual channels with your editor.

Call for Papers

Submit your research to the newest title from APS, Function.


academic.oup.com

OPEN

Developed by:



Powered by:



Follow us on @ScimagoJR

Scimago Lab, Copyright 2007-2020. Data Source: Scopus®

EST MODUS IN REBUS

Horatio (Satire 1.1.106)

Front cover: *Acropora copiosa* Nemenzo, 1971
(PHOTO: DLSU.EDU.PH)

Published monthly

PRINTED IN INDONESIA

ISSN: 1412-033X

E-ISSN: 2085-4722



9 771412 033764




9 772085 472768

ISSN: 1412-033X
E-ISSN: 2085-4722

BIODIVERSITAS

Journal of Biological Diversity

Volume 21 - Number 5 - May 2020



BIODIVERSITAS

Journal of Biological Diversity
Volume 21 – Number 5 – May 2020

ISSN/E-ISSN:

1412-033X (printed edition), 2085-4722 (electronic)

EDITORIAL BOARD:

Abdel Fattah N.A. Rabou (Palestine), **Agnieszka B. Najda** (Poland), **Ajay Kumar Gautam** (India), **Alan J. Lymbery** (Australia), **Annisa** (Indonesia), **Bambang H. Saharjo** (Indonesia), **Daiane H. Nunes** (Brazil), **Darlina Md. Naim** (Malaysia), **Ghulam Hassan Dar** (India), **Hassan Pourbabaei** (Iran), **Joko R. Witono** (Indonesia), **Kartika Dewi** (Indonesia), **Katsuhiko Kondo** (Japan), **Kusumadewi Sri Yulita** (Indonesia), **Livia Wanntorp** (Sweden), **M. Jayakara Bhandary** (India), **Mahdi Reyahi-Khoram** (Iran), **Mahendra K. Rai** (India), **Mahesh K. Adhikari** (Nepal), **Maria Panitsa** (Greece), **Mochamad A. Soendjoto** (Indonesia), **Mohib Shah** (Pakistan), **Mohamed M.M. Najim** (Srilanka), **Nurhasanah** (Indonesia), **Praptiwi** (Indonesia), **Rasool B. Tareen** (Pakistan), **Seyed Aliakbar Hedayati** (Iran), **Seyed Mehdi Talebi** (Iran), **Shahabuddin** (Indonesia), **Shahir Shamsir** (Malaysia), **Shri Kant Tripathi** (India), **Subhash C. Santra** (India), **Sugeng Budiharta** (Indonesia), **Sugiyarto** (Indonesia), **Taufiq Purna Nugraha** (Indonesia), **Yosep S. Mau** (Indonesia)

EDITOR-IN-CHIEF:

S u t a r n o

EDITORIAL MEMBERS:

English Editors: **Graham Eagleton** (grahameagleton@gmail.com), **Suranto** (surantouns@gmail.com); Technical Editor: **Solichatun** (solichatun_s@yahoo.com), **Artini Pangastuti** (pangastuti_tutut@yahoo.co.id); Distribution & Marketing: **Rita Rakhmawati** (oktia@yahoo.com); Webmaster: **Ari Pitoyo** (aripitoyo@yahoo.com)

MANAGING EDITORS:

Ahmad Dwi Setyawan (unsjournals@gmail.com)

PUBLISHER:

The Society for Indonesian Biodiversity

CO-PUBLISHER:

Department of Biology, Faculty of Mathematics and Natural Sciences, Sebelas Maret University, Surakarta

ADDRESS:

Jl. Ir. Sutami 36A Surakarta 57126. Tel. +62-271-7994097, Tel. & Fax.: +62-271-663375, email: editors@smujo.id

ONLINE:

biodiversitas.mipa.uns.ac.id; smujo.id/biodiv



Society for Indonesia
Biodiversity



Sebelas Maret University
Surakarta

GUIDANCE FOR AUTHORS

Aims and Scope *Biodiversitas*, *Journal of Biological Diversity* or abbreviated as *Biodiversitas* encourages submission of manuscripts dealing with all biodiversity aspects of plants, animals and microbes at the level of the gene, species, and ecosystem as well as ethnobiology.

Article types The journal seeks original full-length research papers, reviews, and short communication. Manuscript of original research should be written in no more than 8,000 words (including tables and picture), or proportional with articles in this publication number. Review articles will be accommodated, while, short communication should be written at least 2,000 words, except for pre-study.

Submission The journal only accepts online submission, through open journal system (<https://smujo.id/biodiv/about/submissions>) or email to the editors at unsjournals@gmail.com. Submitted manuscripts should be the original works of the author(s). The manuscript must be accompanied by a cover letter containing the article title, the first name and last name of all the authors, a paragraph describing the claimed novelty of the findings versus current knowledge. Submission of a manuscript implies that the submitted work has not been published before (except as part of a thesis or report, or abstract); and is not being considered for publication elsewhere. When a manuscript written by a group, all authors should read and approve the final version of the submitted manuscript and its revision; and agree the submission of manuscripts for this journal. All authors should have made substantial contributions to the concept and design of the research, acquisition of the data and its analysis; drafting of the manuscript and correcting of the revision. All authors must be responsible for the quality, accuracy, and ethics of the work.

Ethics Author(s) must obedient to the law and/or ethics in treating the object of research and pay attention to the legality of material sources and intellectual property rights.

Copyright If and when the manuscript is accepted for publication, the author(s) still hold the copyright and retain publishing rights without restrictions. Authors or others are allowed to multiply article as long as not for commercial purposes. For the new invention, authors are suggested to manage its patent before published.

Open access The journal is committed to free-open access that does not charge readers or their institutions for access. Readers are entitled to read, download, copy, distribute, print, search, or link to the full texts of articles, as long as not for commercial purposes. The license type is CC-BY-NC-SA.

Acceptance The only articles written in English (U.S. English) are accepted for publication. Manuscripts will be reviewed by editors and invited reviewers (double blind review) according to their disciplines. Authors will generally be notified of acceptance, rejection, or need for revision within 1 to 2 months of receipt. The manuscript is rejected if the content does not in line with the journal scope, does not meet the standard quality, inappropriate format, complicated grammar, dishonesty (i.e. plagiarism, duplicate publications, fabrication of data, citations manipulation, etc.), or ignoring correspondence in three months. The primary criteria for publication are scientific quality and biodiversity significance. **Uncorrected proofs** will be sent to the corresponding author by email as *.doc* or *.docx* files for checking and correcting of typographical errors. To avoid delay in publication, corrected proofs should be returned in 7 days. The accepted papers will be published online in a chronological order at any time, but printed in the early of each month (12 times).

A charge Starting on January 1, 2019, publishing costs waiver is granted to authors of graduate students from **Least Developed Countries**, who first publish the manuscript in this journal. However, other authors are charged USD 250 (IDR 3,500,000). Additional charges may be billed for language editing, USD 75-150 (IDR 1,000,000-2,000,000).

Reprints The sample journal reprint is only available by special request. Additional copies may be purchased when ordering by sending back the uncorrected proofs by email.

Manuscript preparation Manuscript is typed on A4 (210x297 mm²) paper size, in a single column, single space, 10-point (10 pt) Times New Roman font. The margin text is 3 cm from the top, 2 cm from the bottom, and 1.8 cm from the left and right. Smaller lettering size can be applied in presenting table and figure (9 pt). Word processing program or additional software can be used, however, it must be PC compatible and Microsoft Word based (*.doc* or *.rtf*; not *.docx*). **Scientific names** of species (incl. subspecies, variety, etc.) should be written in italic, except for italic sentence. Scientific name (genera, species, author), and cultivar or strain should be mentioned completely for the first time mentioning it in the body text, especially for taxonomic manuscripts. Name of genera can be shortened after first mentioning, except generating confusion. Name of the author can be eliminated after first mentioning. For example, *Rhizopus oryzae* L. UICC 524, hereinafter can be written as *R. oryzae* UICC 524. Using trivial name should be avoided, otherwise generating confusion. **Biochemical and chemical nomenclature** should follow the order of the IUPAC - IUB. For DNA sequence, it is better used Courier New font. Symbols of standard chemical and abbreviation of chemistry name can be applied for common and clear used, for example, completely written butilic hydroxyl toluene (BHT) to be BHT hereinafter. **Metric measurement** use IS denomination, usage other system should follow the value of equivalent with the denomination of IS first mentioning. Abbreviations set of, like g, mg, mL, etc. do not follow by dot. Minus index (m⁻², L⁻¹, h⁻¹) suggested to be used, except in things like "per-plant" or "per-plot". **Equation of mathematics** does not always can be written

down in one column with text, in that case can be written separately. **Number** one to ten are expressed with words, except if it relates to measurement, while values above them written in number, except in early sentence. The fraction should be expressed in decimal. In the text, it should be used "%" rather than "percent". Avoid expressing ideas with complicated sentence and verbiage, and used efficient and effective sentence.

Title of the article should be written in compact, clear, and informative sentence, preferably not more than 20 words. Name of author(s) should be completely written. **Name and institution** address should also be completely written with street name and number (location), postal code, telephone number, facsimile number, and email address. Manuscript written by a group, author for correspondence along with address is required. First page of the manuscript is used for writing above information.

Abstract should not be more than 200 words. **Keywords** is about five words, covering scientific and local name (if any), research theme, and special methods which used; and sorted from A to Z. All important **abbreviations** must be defined at their first mention. **Running title** is about five words. **Introduction** is about 400-600 words, covering the background and aims of the research. **Materials and Methods** should emphasize on the procedures and data analysis. **Results and Discussion** should be written as a series of connecting sentences, however, for manuscript with long discussion should be divided into subtitles. Thorough discussion represents the causal effect mainly explains for why and how the results of the research were taken place, and do not only re-express the mentioned results in the form of sentences. **Concluding** sentence should be given at the end of the discussion. **Acknowledgments** are expressed in a brief; all sources of institutional, private and corporate financial support for the work must be fully acknowledged, and any potential conflicts of interest are noted.

Figures and Tables of maximum of three pages should be clearly presented. Title of a picture is written down below the picture, while title of a table is written above the table. Colored figures can only be accepted if the information in the manuscript can lose without those images; chart is preferred to use black and white images. Author could consign any picture or photo for the front cover, although it does not print in the manuscript. All images property of others should be mentioned source. **There is no appendix**, all data or data analysis are incorporated into Results and Discussions. For broad data, it can be displayed on the website as a supplement.

References Author-year citations are required. In the text give the authors name followed by the year of publication and arrange from oldest to newest and from A to Z. In citing an article written by two authors, both of them should be mentioned, however, for three and more authors only the first author is mentioned followed by et al., for example: Saharjo and Nurhayati (2006) or (Boonkerd 2003a, b, c; Sugiyarto 2004; El-Bana and Nijs 2005; Balagadde et al. 2008; Webb et al. 2008). Extent citation as shown with word "*cit*" should be avoided. Reference to unpublished data and personal communication should not appear in the list but should be cited in the text only (e.g., Rifai MA 2007, pers. com. (personal communication); Setyawan AD 2007, unpublished data). In the reference list, the references should be listed in an alphabetical order (better, if only 20 for research papers). Names of journals should be abbreviated. Always use the standard abbreviation of a journal's name according to the **ISSN List of Title Word Abbreviations** (www.issn.org/2-22661-LTWA-online.php). The following examples are for guidance.

Journal:

Saharjo BH, Nurhayati AD. 2006. Domination and composition structure change at hemic peat natural regeneration following burning; a case study in Pelalawan, Riau Province. *Biodiversitas* 7: 154-158.

Book:

Rai MK, Carpinella C. 2006. Naturally Occurring Bioactive Compounds. Elsevier, Amsterdam.

Chapter in book:

Webb CO, Cannon CH, Davies SJ. 2008. Ecological organization, biogeography, and the phylogenetic structure of rainforest tree communities. In: Carson W, Schnitzer S (eds) *Tropical Forest Community Ecology*. Wiley-Blackwell, New York.

Abstract:

Assaed AM. 2007. Seed production and dispersal of *Rhazya stricta*. 50th annual symposium of the International Association for Vegetation Science, Swansea, UK, 23-27 July 2007.

Proceeding:

Alikodra HS. 2000. Biodiversity for development of local autonomous government. In: Setyawan AD, Sutarno (eds.) *Toward Mount Lawu National Park; Proceeding of National Seminary and Workshop on Biodiversity Conservation to Protect and Save Germplasm in Java Island*. Universitas Sebelas Maret, Surakarta, 17-20 July 2000. [Indonesian]

Thesis, Dissertation:

Sugiyarto. 2004. Soil Macro-invertebrates Diversity and Inter-Cropping Plants Productivity in Agroforestry System based on Sengon. [Dissertation]. Universitas Brawijaya, Malang. [Indonesian]

Information from internet:

Balagadde FK, Song H, Ozaki J, Collins CH, Barnet M, Arnold FH, Quake SR, You L. 2008. A synthetic *Escherichia coli* predator-prey ecosystem. *Mol Syst Biol* 4: 187. www.molecularsystemsbiology.com

[Contact Us](#)
[FAQ](#)
[Useful links](#)
[Home](#)
[Content and Archives](#)
[Aims and Scopes](#)
[Editorial Board](#)
[Guidance for Authors](#)
[Ethical Guidelines](#)
[Charges](#)
[Membership](#)
[Previously Reviewers](#)
[Conference Events](#)

Biodiversitas Vol. 21, No. 4, April 2020

 only search Biodiversitas

Table of Content

[Cover \[PDF\]](#)
[Editorial Board & Guidance for Authors \[PDF\]](#)

1. Anatomical examination of the petiole of eupolypods I (Polypodiales)
JEANETTE MARA P. TAN, MARIA CELESTE BANATICLA-HILARIO, PASTOR MALABRIGO, MARJORIE DELOS ANGELES, INOCENCIO E. BUOT, JR. [\[PDF\]](#)
2. The growth of three varieties of black pepper (*Piper nigrum*) under different light intensities related to indigenous hormones role
ISSUKINDARSYAH, ENDANG SULISTYANINGSIH, DIDIK INDRADEWA, EKA TARWACA SUSILA PUTRA [\[PDF\]](#)
3. Short Communication: The composition of undergrowth vegetation in the Gendol Riverbank, Sleman District, Yogyakarta, Indonesia
ATUS SYAHBUDIN, ALNUS MEINATA, WIYONO, RIDLA ARIFRIANA [\[PDF\]](#)
4. Diversity of soil organic carbon and water characteristics under different vegetation types in northern Bengkulu, Indonesia
BANDI HERMAWAN, HERY SUHARTOYO, BAMBANG SULISTYO, BAMBANG GONGGO MURCITRO, WELLY HERMAN [\[PDF\]](#)
5. Distribution of *Gyrinops versteegii* (Gilg.) Domke in varying vegetation structures, soil properties, and microclimates in Manggarai, Flores, East Nusa Tenggara
TITUT YULISTYARINI, ABBAN PUTRI FIQA, SUGENG BUDIHARTA, RIDESTI RINDYASTUTI [\[PDF\]](#)
6. Short Communication: Isolation and characterization of the endophytic bacteria, and their potential as maize diseases control
ENDANG MUGIASTUTI, SUPRAYOGI, NUR PRIHATININGSIH, LOEKAS SOESANTO [\[PDF\]](#)
7. Coral transplantation on a multilevel substrate of Artificial Patch Reefs: effect of fixing methods on the growth rate of two *Acropora* species
MUNASIK, AGUS SABDONO, AZELIA N. ASSYFA, DIAH PERMATA WIJAYANTI, SUGIYANTO, IRWANI IRWANI, RUDHI PRIBADI [\[PDF\]](#)
8. Morphological variation of two common sea grapes (*Caulerpa lentillifera* and *Caulerpa racemosa*) from selected regions in the Philippines
JEREMIAH L. ESTRADA, NONNATUS S. BAUTISTA, MARIBEL L. DIONISIO-SESE [\[PDF\]](#)
9. Effects of supplementation with phosphorus, calcium and manganese during oil palm frond fermentation by *Phanerochaete chrysosporium* on ligninase enzyme activity
RONI PAZLA, NOVIRMAN JAMARUN, FAUZIA AGUSTIN, MARDIATI ZAIN, ARIEF, NESYA OKTIA CAHYANI [\[PDF\]](#)
10. Toxicity of entomopathogenic fungal culture filtrate of lowland and highland soil of South Sumatra (Indonesia) against *Spodoptera litura* larvae
MIMMA GUSTIANINGTYAS, SITI HERLINDA, SUWANDI, SUPARMAN, HARMAN HAMIDSON, HASBI, ARUM SETIAWAN, MARIESKA VERAWATY, ELFITA, ARSI [\[PDF\]](#)
11. Ethnobotany and conservation of indigenous edible fruit plants in South Aceh, Indonesia
ADI BEJO SUWARDI, ZIDNI ILMAN NAVIA, TISNA HARMAWAN, SYAMSUARDI, ERIZAL MUKHTAR [\[PDF\]](#)
12. Ethnobotanical knowledge and conservation practices of indigenous people of Mbeliling Forest Area, Indonesia
MARLINDA MULU, ZEPHISIUS R.E. NTELOK, PETRUS SII, HILDEGARDIS MULU [\[PDF\]](#)
13. Short Communication: Community of phytoplankton in peatland canal, Riau, and wet dune slacks of Parangtritis, Yogyakarta, Indonesia
ANNISA MAWARNI, FIRDA N.N. AZIZAH, HENI W. SARTIKA, SUWARNO HADISUSANTO, DWINDA M. PUTRI, AKBAR REZA [\[PDF\]](#)
14. Assessment on the growth performance of planted *Dryobalanops beccarii* at reforestation sites after implementation of selective girdling
MOHD EFFENDI WASLI, DOUGLAS BUNGAN AMBUN, MEEKIONG KALU, MOGERET SIDI, HAFSAH NAHRAWI, HASHIMAH ELIAS [\[PDF\]](#)
15. Better providers of habitat for Javan slow loris (*Nycticebus javanicus* E. Geoffroy 1812): A species distribution modeling approach in Central Java, Indonesia

- MAHFUT SODIK, SATYAWAN PUDYATMOKO, PUJO SEMEDI HARGO YUWONO, MUHAMMAD TAFRICHAN, MUHAMMAD ALI IMRON [PDF]
16. Identification of growth genes diversity of swamp buffalo using RFLP in Kabaena Island, Bombana District, Southeast Sulawesi, Indonesia
LA ODE NAFIU, MUZUNI, MUHAMMAD AMRULLAH PAGALA, WIDHI KURNIAWAN, SYAM RAHADI [PDF]
 17. Short Communication: Genetic diversity of lemon (*Citrus* spp.) from Ternate Island (Indonesia) based on morphological and molecular characters
ABDULRASYID TOLANGARA, ALOYSIUS DURAN COREBIMA, ABDU MAS'UD, SUNDARI [PDF]
 18. Short Communication: The type and sound diversity of Kukuak Balenggek chicken (*Gallus gallus domesticus*) reared in West Sumatra, Indonesian
FIRDA ARLINA, RUSFIDRA, DICKY ANDRIANO, CECE SUMATRI [PDF]
 19. The potency of endophytic bacteria isolated from *Ficus septica* as phytoremediation promoting agent of Cr (VI) contaminated soil
NITA SHILFIANI ROHMAH, SUHARJONO, YOGA DWI JATMIKO, DIAN SISWANTO, IRFAN MUSTAFA [PDF]
 20. Evaluation of wild plants as lead (Pb) and cadmium (Cd) accumulators for phytoremediation of contaminated rice fields
NURIL HIDAYATI [PDF]
 21. Effectiveness of biological control of *Trichoderma harzianum* on soybean leaf rust disease and the production in West Papua Lowland, Indonesia
EKO AGUS MARTANTO, ADELIN E. TANATI, SAMEN BAAN, HERMAN R. TATA, AGUSTINUS MURDJOKO [PDF]
 22. Fauna diversity, production potential and total economic value of mangrove ecosystems in Mentawir Village, East Kalimantan, Indonesia
ROCHADI KRISTININGRUM, ABUBAKAR M. LAHJIE, MASJAYA, SYAHRIR YUSUF, YOSEP RUSLIM, AMIR MA'RUF [PDF]
 23. Fungal isolates from marine sponge *Chelonaplysilla* sp.: Diversity, antimicrobial and cytotoxic activities
DIAN HANDAYANI, MUH. ADE ARTASASTA, NILDA SAFIRNA, DIANA FITRI AYUNI, TRINA EKAWATI TALLEI, TRIANA HERTIANI [PDF]
 24. Helminth fauna of *Microtus cf. arvalis* (Rodentia, Cricetidae) in Russia and adjacent countries
NADEZHDA Yu. KIRILLOVA, ALEXANDER A. KIRILLOV, ALEXANDER B. RUCHIN, MAXIM V. TRUKHACHEV [PDF]
 25. Assessment of the health status of the Sidi R'Ghies forest, Oum El Bouaghi, north-east Algerian
MALIKA RACHED-KANOUNI, ALIA ZERROUKI, MAROUA LAHMAR, AMINA BELDJAZIA, KARIMA KARA, LABED ABABSA [PDF]
 26. Floristic analysis of semi-arid mountain ecosystems of the Griqualand West centre of plant endemism, Northern Cape, South Africa
NANETTE VAN STADEN, STEFAN JOHN SIEBERT, DIRK PETRUS CILLIERS, DIAN WILSENACH, ARNOLD WALTER FRISBY [PDF]
 27. Spontaneous plant recolonization on reclaimed post-coal mining sites in East Kalimantan, Indonesia: Native versus alien and succession progress
LIA HAPSARI, TRIMANTO, SUGENG BUDIHARTA [PDF]
 28. Growth prediction for rubber tree and intercropped forest trees to facilitate environmental services valuation in South Thailand
NARUN NATTHAROM, SAOWALAK ROONGTAWANREONGSRI, SARA BUMRONGSRI [PDF]
 29. Dispersion of tongka langit banana in Buru and Seram, Maluku Province, Indonesia, based on topographic and climate factors
HALVINA GASELA SAIYA, ADRIANA HIARIEJ, ANNEKE PESIK, ELIZABETH KAYA, MEITTY LOUISE HEHANUSSA, FERAD PUTURUHU [PDF]
 30. Genetic variability of Indonesian *Oryctes rhinoceros nudivirus* (OrNV) as genus of *Alphanudivirus*
SAT RAHAYUWATI, YAYI MUNARA KUSUMAH, SUDHARTO PRAWIROKARTO, DADANG, TEGUH SANTOSO [PDF]
 31. Population ecology size and habitat preference of the ghost orchid *Didymoplexis pallens* in Bogor Botanic Gardens, Indonesia
RIZMOON NURUL ZULKARNAEN, R. VITRI GARVITA, HARY WAWANGNINGRUM, KARTIKA NING TYAS [PDF]
 32. Antioxidant activity screening of seven Indonesian herbal extract
JOHAN SUKWEENADHI, OEKE YUNITA, FINNA SETIAWAN, KARTINI, MAYA THERESA SIAGIAN, ANGGREYNI PRATIWI DANDURU, CHRISTINA AVANTI [PDF]
 33. Short Communication: First record of *Hirschmanniella mucronata* (Nematoda: Pratylenchidae) in Yogyakarta, Indonesia
SIWI INDARTI, ALAN SOFFAN, MUHAMMAD MAULANA FARDANI ANDRASMARA [PDF]
 34. Short Communication: Polymorphism at third exon of the Myostatin gene and its association with growth and carcass traits in Batur sheepa
HASSAN ISHAG HASSAN HAREN [PDF]
 35. Abundance of ants (Hymenoptera: Formicidae) and the functional groups in two different habitats
ANANTO TRIYOGO, BUDIADI, S.M. WIDYASTUTI, SENA ADI SUBRATA, SUWITO SETYO BUDI [PDF]
 36. Nesting behavior of reintroduced Bornean Orangutan in Bukit Batikap Conservation Forest, Central Kalimantan, Indonesia
IKE NURJUITA NAYASILANA, SRI SUCI UTAMI ATMOKO, AHMAT SUYOKO, SUWARNO HADISUSANTO [PDF]
 37. Genetic diversity of blue swimming crab (*Portunus pelagicus* Linn 1758) from Indonesian waters (Sunda and Sahul Shelf, Wallacea region): Phylogenetic approach

- ANDI ALIAH HIDAYANI, YUSHINTA FUJAYA, DODY DH.TRIJUNO; NITA RUKMINASARI; ALIMUDDIN ALIMUDDIN [PDF]
38. The Isolation and identification of cellulolytic bacteria at fibric, hemic and sapric peat in Teluk Bakung Peatland, Kubu Raya District, Indonesia
SITI KHOTIMAH, SUHARJONO, TRI ARDYATI, YULIA NURANI [PDF]
 39. Short Communication: Preliminary phylogenetic analysis of bacteria producing laccase isolated from Gunung Pancar, Bogor, Indonesia
WIN WIN MAR, ALI ROHMAN, NUR H. MUWAFIQI, GALIH AYHUSTA LARAS, DYAH AGUSTINA, ONE ASMARANI, NI NYOMAN TRI PUSPANINGSIH [PDF]
 40. Short Communication: Biological control of Fusarium wilt on banana plants using biofertilizers
ARIEF WIDYANTORO, HADIWIYONO, SUBAGIYA [PDF]
 41. The impact of bioinsecticide overdoses of *Beauveria bassiana* on species diversity and abundance of not targeted arthropods in South Sumatra (Indonesia) freshwater swamp paddy
KHOIRUL IKHSANUDIN HANIF, SITI HERLINDA, CHANDRA IRSAN, YULIA PUJIASTUTI, GHANNI PRABAWATI, HASBI, TILI KARENINA [PDF]
 42. Autecology of *Drosera burmanni* in the Wolobobo Botanic Gardens, Ngada District, Flores Island, Indonesia
JOKO RIDHO WITONO, DIDI USMADI, WIHERMANTO, DANANG WAHYU PURNOMO, DINA SAFARINANUGRAHA, YANUAR PAKIDING, YANUAR PAKIDING, NELWAN NETOSES [PDF]
 43. An inventory of useful threatened plant species in Vhembe Biosphere Reserve, Limpopo Province, South Africa
LUAMBO JEFFREY RAMARUMO, ALFRED MAROYI [PDF]
 44. The genetic relationships and Indo-Pacific connectivity of whale sharks (*Rhincodon typus*) with particular reference to mitochondrial COI gene sequences from Cendrawasih Bay, Papua, Indonesia
ABDUL HAMID A. TOHA, MUHAMMAD DAILAMI, SAIFUL ANWAR, JUSWONO B. SETIAWAN, YUSUP JENTEWO, IDA LAPADI, SANNY SUTANTO, RATIH ARYASARI, AMBARIYANTO, FERAWATI RUNTUBOI, HAWIS MADDUPPA [PDF]
 45. Nesting behavior of Bornean immature Orangutan (*Pongo pygmaeus wurmbii*) in Nyaru Menteng Arboretum School, Palangka Raya, Central Kalimantan, Indonesia
FOUAD FAUZI, SUEMARNO, AMINUDIN AFANDHI, AMIN SETYO LEKSONO [PDF]
 46. The effect of drying treatment to metabolite profile and cytotoxic potential of *Rhizophora apiculata* leaves
MADA TRIANDALA SIBERO, ANGGUN PUSPITARINI SISWANTO, RUDHI PRIBADI, AGUS SABDONO, OCKY KARNA RADJASA, AGUS TRIANTO, EVAN HANSEL FREDERICK, ALDI PRATAMA WIJAYA, DWI HARYANTI, DESY WULAN TRININGSIH, SASTRA JENDRA HAYUNINGRAT, YASUHIRO IGARASHI [PDF]
 47. Germination and salinity tolerance of seeds of sixteen Fabaceae species in Thailand for reclamation of salt-affected lands
YONGKRIAT KU-OR, NISA LEKSUNGOEN, DAMRONGVUDHI ONWIMON, PEERAPAT DOOMNIL [PDF]
 48. Short Communication: Antioxidant and antibacterial properties of tree fern *Cyathea* contaminants
AHMAD FAIZAL, INTAN TAUFIK, ANISAH FIRDA RACHMANI, ALDA WYDIA PRIHARTINI AZAR [PDF]
 49. The identification of plant reliefs in the Lalitavistara story of Borobudur temple, Central Java, Indonesia
DESTARIO METUSALA, FAUZIAH, DEWI AYU LESTARI, JANIS DAMAIYANI, SHOFIYATUL MASâ€™UDAH, HARI SETYAWAN [PDF]
 50. Microbiological and physicochemical characteristics of bakasang laor, a traditional fermented fishery product from Maluku, Indonesia
FERYMON MAHULETTE, TRI SANTI KURNIA [PDF]
 51. Distribution survey of Klossâ€™s Gibbons (*Hylobates klossii*) in Mentawai Islands, Indonesia
ARIF SETIAWAN, CHRISTIAN SIMANJUNTAK, ISMAEL SAUMANUK, DAMIANUS TATEBURUK, YOAN DINATA, DARMAWAN LISWANTO, ANJAR RAFIASTANTO [PDF]
 52. Feeding selectivity of *Holothuria atra* in different microhabitat in Panjang Island, Jepara (Java, Indonesia)s
RETNO HARTATI, AMBARIYANTO AMBARIYANTO, WIDIANINGSIH WIDIANINGSIH, MUHAMMAD ZAINURI [PDF]
 53. Short Communication: Application of bacteriocin from *Lactobacillus plantarum* SRCM 1 004 34 strain isolated from okara as a natural preservative in beef sausage
SALAM ARITONANG, ELLY ROZA, AFRIANI SANDRA [PDF]
 54. Orchid exploration in Tanjung Peropa Wildlife Reserves for Kendari Botanic Gardens collection, Indonesia
SRI HARTINI, POPI APRILIANI [PDF]
 55. Diversity of capsaicin content, quantitative, and yield components in chili (*Capsicum annuum*) genotypes and their F1 hybrid
ZULFIKAR DAMARALAM SAHID, MUHAMAD SYUKUR, AWANG MAHARIJAYAI [PDF]
 56. Potential mitochondrial diversity role in the productivity of three lines of Japanese quails
TAMADHUR HUSSEIN , MOHAMMED BAQUR S. AL-SHUHAIB, TAHREER M. AL-THUWAINI [PDF]
 57. Environment carrying capacity and willingness to pay for bird-watching ecotourism in Kerandangan Natural Park, Lombok, Indonesia
I WAYAN SUANA, HILMAN AHYADI, GITO HADIPRAYITNO, SALEH AMIN, LALU ACHMAD TAN TILAR WANGSAJATI SUKMARING KALIH, FRANCISCUS XAVERIUS SUDARYANTO [PDF]
 58. The combination of mare's milk and grape polyphenol extract for treatment of dysbiosis induced by dextran sulfate sodium
SAMAT KOZHAKHMETOV, DMITRIY BABENKO, MADIYAR NURGAZIYEV, ALTYNAY TUYAKOVA, AYAUPLYM NURGOZHINA, NURISLAM MUHANBETGANOV, LAURA CHULENBAYEVA, SHYNGGYS SERGAZY, ALEXANDR GULYAYEV, TIMUR SALIEV, ALMAGUL KUSHUGULOVA [PDF]
 59. Synthesis of lactic acid from sugar palm trunk waste (*Arenga pinnata*): Preliminary hydrolysis and fermentation

studies

WHINY HARDIYATI ERLIANA, TRI WIDJAJA, ALI ALTWAY, LILY PUDJIASTUTI [[PDF](#)]

60. Effectiveness of herbs containing *Curcuma xanthorrhiza*, *Elephantopus scaber* L and *Amaranthus tricolor* L in iron deficiency anemia patients
ULFA FITRIANI, ZURAIIDA ZULKARNAIN, FAJAR NOVIANTO, ENGGAR WIJAYANTI, AGUS TRIYONO [[PDF](#)]
61. Biodecolorization of methyl orange by mixed cultures of brown-rot fungus *Daedalea dickinsii* and bacterium *Pseudomonas aeruginosa*
ADI SETYO PURNOMO, MITHA OCDYANI MAWADDAH [[PDF](#)]
62. Autecology of *Melastoma malabathricum*, an invasive species in the Way Kambas National Park, Indonesias
JANI MASTER, IBNUL QAYIM, DEDE SETIADI, NYOTO SANTOSO [[PDF](#)]
63. Diversity, abundance, activity period, and factors affecting the appearance of wildlife around the corridors between Khao Yai- Thap Lan National Parks, Thailand by camera trapping
RONGLARP SUKMASUANG, KHWANRUTAI CHARASPET, TARAPORN PANGANTA, MANANYA PLA-ARD, NORASET KHIOESREE, JIDAPA THONGBANTHUM [[PDF](#)]
64. Tiger grass (*Thysanolaena maxima*) cultivation in CALSANAG watershed in Romblon, Philippines: dilemmas and prospects for sustainable natural resources management
LEILA LANDICHO, MARIA THERESA NEMESIS P. OCAMPO, ROWENA ESPERANZA D. CABAUG, ROMNICK S. BALITON, EDGARDO ANDALECIO, RAYMUND INOCENCIO, MARCELINA SERVANEZ, RUSSEL SON A. COSICO, MARYANNE G. ABADILLOS, ARNOLD KARL A. CASTILLO [[PDF](#)]
65. Isolation and characterization of *Botryococcus braunii* from a freshwater environment in Tenggara, Kutai Kartanegara, Indonesia
RUDY AGUNG NUGROHO, DIRGARINI JULIA NURLIANTI SUBAGYONO, ENOS TANGKE ARUNG [[PDF](#)]

[Print on demand \[PDF\]](#)

[Home](#) | [Archives](#) | [Contact Us](#) | [Affiliate Program](#)


Copyright © ½ Biodiversitas, 2012. All Rights Reserved

[Contact Us](#)
[FAQ](#)
[Useful links](#)
[Home](#)
[Content and Archives](#)
[Aims and Scopes](#)
[Editorial Board](#)
[Guidance for Authors](#)
[Ethical Guidelines](#)
[Charges](#)
[Membership](#)
[Previously Reviewers](#)
[Conference Events](#)

Biodiversitas Vol. 21, No. 5, May 2020
ISSN: 1412-033X
E-ISSN: 2085-4722
Editor in Chief: Sutarno
First Published: January 2000

 only search Biodiversitas


NEW UPDATE


21 (4) 2020
21 (3) 2020
21 (2) 2020
21 (1) 2020
**20 (12)
2019**
**20 (11)
2019**
**20 (10)
2019**
20 (9) 2019

ONLINE FIRST PUBLICATION

After the publication of Biodiversitas vol. 15, no. 2, October 2014, furthermore, all manuscripts will be published online as soon as the revision is approved (no time constraints), but they will be printed in the next issue, namely Biodiversitas vol. 16, no. 1, April 2015, and so on.

Examples of how to cite:

- *Prior to the date of printing (before 1 April 2015):*
Tavankar F, Bonyad AE. 2014. Effects of timber harvest on the structural diversity and species composition in hardwood forests. Biodiversitas DOI: 10.13057/biodiv/d160101. (Year is referring to the download time)
- *After the print date (after 1 April 2015):*
Tavankar F, Bonyad AE. 2015. Effects of timber harvest on the structural diversity and species composition in hardwood forests. Biodiversitas 16 (1): 1-9.

NOTIFICATION

Starting on January 1, 2019, Biodiversitas will be issued monthly. New issues will also be found at [Smujo](#), as an online first publication.



DOAJ DIRECTORY OF
OPEN ACCESS
JOURNALS



[Home](#) | [Archives](#) | [Contact Us](#) | [Affiliate Program](#)



All content on this website, unless otherwise noted, is licensed under a Creative Commons Attribution 4.0 License.

Toxicity of entomopathogenic fungal culture filtrate of lowland and highland soil of South Sumatra (Indonesia) against *Spodoptera litura* larvae

By Arum Setiawan

Toxicity of entomopathogenic fungal culture filtrate of lowland and highland soil of South Sumatra (Indonesia) against *Spodoptera litura* larvae

MIMMA GUSTIANINGTYAS¹, SITI HERLINDA^{1,2,3,*}, SUWANDI^{1,2,3}, SUPARMAN^{1,2}, HARMAN HAMIDSON^{1,2}, HASBI^{3,4}, ARUM SETIAWAN⁵, MARIESKA VERAWATY⁵, ELFITA⁶, ARSI^{2,3}

¹Program of Crop Sciences, Faculty of Agriculture, Universitas Sriwijaya, Palembang 30139, South Sumatera, Indonesia

²Department of Plant Pests and Diseases, Faculty of Agriculture, Universitas Sriwijaya, Indralaya 30662, South Sumatera, Indonesia.

Tel.: +62-711-580663, Fax.: +62-711-580276, *email: sitiherlinda@unsri.ac.id

³Research Center for Sub-optimal Lands (PUR-PLSO), Universitas Sriwijaya, Palembang 30139, South Sumatera, Indonesia

⁴Department of Agricultural Engineering, Faculty of Agriculture, Universitas Sriwijaya, Indralaya, Ogan Ilir 30662, South Sumatera, Indonesia

⁵Department of Biology, Faculty of Mathematics and Natural Sciences, Universitas Sriwijaya, Indralaya, Ogan Ilir 30662, South Sumatera, Indonesia

⁶Department of Chemistry, Faculty of Mathematics and Natural Sciences, Universitas Sriwijaya, Indralaya, Ogan Ilir 30662, South Sumatera, Indonesia

Manuscript received: 6 March 2020. Revision accepted: 7 April 2020.

Abstract. *Gustianingtyas M, Herlinda S, Suwandi, Suparman, Hamidson H, Hasbi, Setiawan A, Verawaty M, Elfita, Arsi. 2020. Toxicity of entomopathogenic fungal culture filtrate of lowland and highland soil of South Sumatra against Spodoptera litura larvae. Biodiversitas 21: 1839-1849.* The use of secondary fungal metabolites for the active ingredient of mycoinsecticide is more effective and more easily integrated with other pest control techniques. This study aimed to measure the toxicity of the culture filtrate of entomopathogenic fungi originating from South Sumatra against the *Spodoptera litura* larvae. *Beauveria bassiana* (25 isolates) and *Metarhizium anisopliae* (20 isolates) of South Sumatra were cultured in liquid media and filtered to produce culture filtrate. The larvae which were sick due to the filtrate showed the symptoms of decreased appetite and were not actively moving, while the dead larvae were characterized by being wrinkled, dry, black integument, and odorless. Mortality caused by *B. bassiana* filtrate was the highest 98% (BJgTs isolate) and not significantly different from the BSwTd2 isolate (94.67%). Yet, the LT₅₀ BSwTd2 isolate was shorter (5.92 days) compared to the LT₅₀ BJgTs isolate (6.35 days). The most toxic *M. anisopliae* filtrate produced the mortality of 96% (MKbTp2 isolate) and 85.33% (MPdB isolate) each of which had LT₅₀ of 7.36 days and 8.09 days, respectively. So, the most toxic culture filtrate was BSwTd2 isolate of *B. bassiana* and MKbTp2 isolate of *M. anisopliae*. The entomopathogenic fungi producing filtrate which are toxic have the potential to be active ingredients of mycoinsecticides.

Keywords: *Beauveria bassiana*, chili, LT₅₀, *Metarhizium anisopliae*, mortality

INTRODUCTION

South Sumatra is generally dominated by lowland lands, namely freshwater swamps and tidal lowlands (Kartika et al. 2018; Karenina et al. 2019), and a small portion of medium and highland. Freshwater swamps can be found, among others in the Districts of Ogan Ilir, Ogan Komering Ilir, Musi Banyuasin, and Palembang City; the widest tide exists in Banyuasin District and the medium and highlands occur, among others, in the Districts of Ogan Komering Ulu Selatan, Lahat, and Pagaralam City. The varied topographical conditions between districts/cities characterize the differences including the crops and microorganisms cultivated there.

In the lowlands of South Sumatra are generally cultivated annual crops, such as paddy (Herlinda et al. 2018a; Prabawati et al. 2019), maize (Juhriah et al. 2019), chili (Johari et al. 2016; Nasution and Respatijarti 2019; Sagrim et al. 2017), and palm oil (Darlan et al. 2016). Meanwhile, in the medium and high plains are generally cultivated annual crops such as coffee, tea (Zamhari et al. 2017), vegetables such as potatoes (Maryanto et al. 2018), cabbage, mustard greens, and caisim (Situmorang et al.

2019). Vegetables and seasonal plants of the lowlands and highlands generally have a major pest, namely *Spodoptera litura* (Turnip et al. 2019). The yield loss caused by its larval stage is more than 35% a year (Bueno et al. 2011). This pest has egg, larval, pupal, and adult stages (Vijaya and Rani 2016). *S. litura* is a polyphagous pest that attacks many vegetables in Indonesia, for example, chili (Nagal et al. 2016), cotton fruit (Maqsood et al. 2017), soybean (Fattah et al. 2018). A polyphagous pest is a pest that has a broad range of host species (more than a family) (Nagal et al. 2016). For the control of *S. litura*, it needs natural enemies such as entomopathogenic fungi that can adapt in low to high altitudes.

In the lowlands and highlands, the soil types have different chemical and physical characteristics; acidic pH soils are generally found in the lowlands and neutral or basic pH soils are found in the highlands (Munir and Herman 2019). Soil pH can affect the presence of entomopathogenic fungi and tends to be neutral or high pH soils which more commonly occur in entomopathogenic fungi than low pH soils (Safitri et al. 2018). Besides pH, the texture of sandy soils tends to be low and muddy which is not suitable for fungal propagule life as well as water-

saturated soil where fungi are rarely spotted to survive (Garrido-Jurado et al. 2011). Temperature (Pinnamaneni et al. 2010) and humidity (Oreste et al. 2015; Liu et al. 2017) also affect pathogenicity of entomopathogenic fungi.

Previous study found that more than 30 isolates of entomopathogenic fungi were found in the freshwater swamp, tidal, and highland soil of South Sumatra (Safitri et al. 2018). Apart from the soil, the entomopathogenic fungi were also found from the insect pests of vegetable and food crops of the lowlands to the highlands of South Sumatra (Herlinda et al. 2018b). Various origins of entomopathogenic fungi tend to produce pathogenicity variations (Sumikarsih et al. 2019).

The pathogenicity of entomopathogenic fungi is influenced by the ability of the fungi to produce secondary metabolites (Zibae et al. 2009). The fungi that develop in insect hemolymph or in liquid media are able to produce toxic metabolites, such as extracellular enzymes, proteins, and toxins (Bandani 2005). In vitro media, toxic metabolites can be produced by the fungi which are grown in liquid media and produce culture filtrate (Soesanto et al. 2019). The use of secondary entomopathogenic fungal metabolites for the active ingredient of mycoinsecticide is more effective and more easily integrated with other pest control techniques (Zibae et al. 2009). Entomopathogenic

fungal culture filtrate found in low to high altitude of South Sumatra potentially gets the most toxic isolates that can be utilized in the development of mycoinsecticide technology. For this reason, this study aimed to measure the toxicity of the culture filtrate isolates of entomopathogenic fungi from low to high land of South Sumatra, Indonesia against *Spodoptera litura* larvae.

MATERIALS AND METHODS

The experiments were carried out at the Entomology Laboratory, Department of Pests and Plant Diseases, Faculty of Agriculture, Universitas Sriwijaya from May to November 2018. The room temperatures and relative humidity during the experiment in the laboratory were 29.78°C and 82.72%, respectively. The entomopathogenic fungi isolates used in this study were explored by Herlinda et al. (2018) and Safitri et al. (2018), (Table 1 and 2) spread from the lowlands to the highlands of South Sumatra. The isolates were grouped into two: 25 isolates of *Beauveria bassiana* (Figure 1) and 20 isolates of *Metarhizium anisopliae* (Figure 2). All isolates were identified by Dr. Suwandi (a mycologist from Universitas Sriwijaya).




Figure 1. *Beauveria bassiana* isolates cultured in SDA media: BJgTs (A), BSmMs (B), BSwTd1 (C), BSwTd2 (D), BSwTd3 (E), BSwTd4 (F), BPdR (G), BKbTp (H), BKKPp2 (I), Ts1d3 (J), BTmPc (K), BTmTr (L), (M) Ts1d2 (M), BTmTs (N), BlePd2 (O), BTmKt (P), BPCmS (Q), BMkMs (R), BtmGa (S), BTmSr (T), BPcPd2 (U), BtmSo (V), BtmMa (W), BtmPe (X), Blepd (Y)



Figure 2. *Metarhizium* isolates cultured in SDA media: MPdB (A), MPdR1 (B), MPdR2 (C), MpdPe (D), MJgMs1 (E), MJgMs2 (F), MPdMs1 (G), MPdMs2 (H), MPdMs3 (I), MjgKeTs (J), MPdMs4 (K), MJgTs2 (L), MKbTp1 (M), MSwTp1 (N), MSwTp2 (O), MSwTp3 (P), MSwTp4 (Q), MKKp1 (R), MKbTp2 (S), MagPd (T)

Table 1. *Beauveria bassiana* isolates from South Sumatra used in this research

Isolate code	Isolate soil origin	Geographic origin
BJgTs	Tidal lowlands, maize	Telang Sari
BsmMs	Tidal lowlands, watermelon	Mulya Sari
BSwTd1	Peatlands, oil palm	Talang Dabok
BSwTd2	Peatlands, oil palm	Talang Dabok
BSwTd3	Peatlands, oil palm	Talang Dabok
BSwTd4	Peatlands, oil palm	Talang Dabok
BPdR	Freshwater swamps, paddy	Rambutan
BKbTp	Highlands, cabbage	Talang Patai
BKKPp2	Highlands, rubber, and coffee	Pulau Pinang
Ts1d3	Peatlands	Talang Dabok
BTmPc	Freshwater swamps	Indralaya
BTmTr	Freshwater swamps	Telang Rejo
Ts1d2	Peatlands	Talang Dabok
BTmTs	Highlands	Mulia Sari
BlePd2	<i>Lipaphis erysimi</i>	Pagardin
BTmkt	Freshwater swamps	Kenten
BPCmS	<i>Pseudopius chalcites</i>	Muara Siban
BMkMs	Highlands	Muara Siban
BtmGa	Freshwater swamps	Gandus
BTmSr	Tidal lowlands	Srikaton
BPcPd2	<i>Chrysodeixis chalcites</i>	Pagardin
BtmSo	Freshwater swamps	Soak
BtmMa	Freshwater swamps	Mariana
BtmPe	Freshwater swamps	Pemulutan
BlepD	<i>Lipaphis erysimi</i>	Pagardin

Table 2. *Metarhizium anisopliae* isolates from South Sumatra used in this research

Isolate code	Isolate soil origin	Geographic origin
MPdB	Freshwater swamps, paddy	Banyuasin
MPdR1	Freshwater swamps, paddy	Rambutan
MPdR2	Freshwater swamps, paddy	Rambutan
MpdPe	Freshwater swamps, paddy	Pemulutan
MJgMs1	Tidal lowlands, maize	Mulya Sari
MJgMs2	Tidal lowlands, maize	Mulya Sari
MPdMs1	Tidal lowlands, paddy	Mulya Sari
MPdMs2	Tidal lowlands, paddy	Mulya Sari
MPdMs3	Tidal lowlands, paddy	Mulya Sari
MjgKeTs	Tidal lowlands, maize, and oil palm	Telang Sari
MPdMs4	Tidal lowlands, paddy	Mulya Sari
MJgTs2	Tidal lowlands, maize	Telang Sari
MKbTp1	Highlands, cabbage	Talang Patai
MSwTp1	Highlands, mustard	Talang Patai
MSwTp2	Highlands, mustard	Talang Patai
MSwTp3	Highlands, mustard	Talang Patai
MSwTp4	Highlands, mustard	Talang Patai
MKKPp1	Highlands, mustard, rubber, coffee	Pulau Pinang
MKbTp2	Highlands, cabbage	Talang Patai
MagPd	Highlands, <i>Aphis gossypii</i>	Pagardin

Test insect preparation

S. litura larvae were collected from the chili planting area in the experimental field of the Faculty of Agriculture, Sriwijaya University and farmers' vegetable fields which were not applied by synthetic insecticides. Then, the larvae were brought to the laboratory and maintained in a plastic cylinder whose upper part was covered with gauze (Ø 25 cm, height 30 cm) having been washed thoroughly using detergent. Next, water spinach plants were put into it for larvae feed.

Every day the larvae feed was replaced with fresh new feed. The larvae entering the pupae phase were transferred into a plastic cylinder (Ø 10 cm, height 15 cm) with the top open and the bottom of the cylinder sprinkled with sifted soil and sterilized in an oven for 1 hour at 100°C. The thickness of the soil inserted into the cylinder was 3 cm. The plastic cylinder containing the pupae was put into a gauze cage (30x30x30 cm³) which had been chopped with chilies for laying eggs. Adults arising from the pupae were fed with honey smeared on cotton and hung over a cage. The hatched eggs were transferred into a plastic cylinder and fed with water spinach which was replaced every day. The larvae used for toxicity testing were the second instar of the third offspring or afterward.

Culture filtrate production

All isolates before being filtered were first made fit by modifying the method of Herlinda (2010). The media for making fit used Sabouraud Dextrose Agar (SDA, Merck) as much as 16.2 g and added 250 mL aquadest enriched with 1.25 g *Tenebrio molitor* larvae flour (5 g). After that, the ingredients were mixed and stirred evenly and then put in an Erlenmeyer (size 250 mL), sterilized in an autoclave for 120 minutes at a pressure of 1 atm. The media was poured into the Petri dish as much as 10 mL (Ø 9 cm) in aseptic laminar flow air conditions. Then, the entomopathogenic fungi isolates were inoculated in the SDA media and incubated for 14 days.

Isolates of entomopathogenic fungi derived from SDA media of aged 14 days were then transferred and grown into liquid media (broth media), PDB (*Potato Dextrose Broth*) prepared as follows: the PDB media composition consisting of 20 g dextrose monohydrate, 200 g potatoes, and 1000 mL aquadest. Before the potatoes were extracted, they were cut into cubes with a size of $\pm 2 \times 2 \times 2$ cm³ and boiled using 1000 mL aquadest for 20 minutes. This PDB medium was sterilized in the autoclave and after it got cold, the entomopathogenic fungus isolates were inoculated into it. This liquid culture was incubated for 6 weeks.

After the fungus liquid culture (fungal broth) was 6 weeks old, the fungus filtration is carried out. The filtration was conducted in two stages, namely filtering using filter paper and syringe filter. In the first stage, 100 mL of 6-week-old fungal broth was filtered using Whatman filter paper no. 42 and coated with a thickness of 1 cm cotton. The culture filtrate obtained from the first stage of filtering was then sucked up to 10 mL using a hypodermic needle (spike). Then, the needle was removed and the base of the needle was fitted with a syringe filter (0.45 µm-25 mm). The 10 mL spike was pressed so that it released culture

filtrate from the syringe filter. The 100 mL of fungal broth produced ± 70 mL of culture filtrate. The culture filtrate was taken as much as 1 mL to be dripped on a piece of chili leaf which was used as a toxicity test. This culture filtrate is often referred to as raw secondary metabolites (Soesanto et al. 2019). To ensure the culture filtrate did not contain propagules (hyphae, mycelia) and conidia, it was grown on agar media (SDA) before it was applied. If it does not grow fungus on the SDA media, the culture filtrate can be applied.

Insecticidal activity test of entomopathogenic fungus culture filtrate

The chili leaves that had been dropped by the culture filtrate were first drained; before 25 *S. litura* larvae were fasted for 2 hours, and then put into them. The second instar larvae were left to eat leaves that had been dropped with culture filtrate for 6 hours. After 6 hours, the larvae were transferred into a plastic cylinder topped with gauze (Ø 25 cm, height 30 cm) containing 15-30 clean chili leaves which were not dripped with culture filtrate. Every day the chili leaves were replaced with the new ones. Every day the dead larvae were recorded up to 12 days after the application.

Data analysis

The differences in mortality data and lethal time that killed 50% of test insects (LT₅₀) were analyzed using analysis of variance (ANOVA), after that further tests were carried out using the 5% Tukey's Honestly Significant Difference (HSD). All data were calculated using software of SAS University Edition 2.7 9.4 M5.

RESULTS AND DISCUSSION

Insecticidal activities of entomopathogenic fungi

The culture filtrate derived from twenty-five *B. bassiana* isolates tested for the insecticidal activity showed that the most toxic *B. bassiana* isolates were those coded BJgTs (98%) and were not significantly different, among others with BSwTd2 (94.67%) and BTmTs isolates (96%). All *B. bassiana* isolates showed the ability to kill the test larvae (Table 3). The BJgTs, BSwTd2, and BTmTs isolates were the most toxic because they produced the highest mortality and also their ability to kill the shortest as evidenced by the short LT₅₀ (6.35, 5.92, and 7.78 days) isolates. The most toxic isolates were shown with the highest mortality and their shortest LT₅₀. Consequently, for *B. bassiana*, the most toxic isolate was BSwTd2.

The culture filtrate of the twenty *M. anisopliae* isolates which were tested for their insecticidal activity showed that the most toxic isolates were that coded MKbTp2 (96%) and MPdB (85.33%) and the mortality of both isolates was significantly different from the other isolates (Table 4). The BSwTd2 isolate was the most toxic because they produced the highest mortality as well as the ability to kill the shortest as evidenced by the shortest LT₅₀ (7.36 days), and then followed by LT₅₀ isolate MPdB (8.09 days). The most

toxic *M. anisopliae* isolate was proven by the highest mortality and the shortest LT₅₀ which was MKbTp2 isolate.

The color of the fungal broth (Figure 3) and culture filtrate (Figure 4) of *B. bassiana* varied among the isolates. All fungal broth isolates were generally transplanted brown, but there were among those isolates dark brown in color, for example, isolates BSwTd2, BSwTd3, BPdR, TS1d2. There was a tendency for the darker color of the fungal broth and culture filtrate to cause higher mortality. The color of fungal broth (Figure 5) and culture filtrate (Figure 6) of *M. anisopliae* differed from the colors of fungal broth and culture of *B. bassiana*. The *M. anisopliae* fungal broth was generally dark greenish-brown, but the culture filtrate tended to have a more varied color, from greenish dark brown to light brown. The same tendency was also produced in *M. anisopliae* isolates, i.e., the color of the older culture filtrate resulting in higher mortality. The isolates having a darker colored culture filtrate were MPdB, MPdMs1, MPdMs2, MPdMs3, MKbTp1, MKbTp2.

Effect of entomopathogenic fungal culture filtrate on test larvae

Spodoptera litura larvae fed with chili leaves applied with *B. bassiana* or *M. anisopliae* culture filtrate exhibited the same behavior, i.e., they increasingly ate less, while the controls not given culture filtrate continued to eat greedily. The chili leaves given to the larvae which were not given culture filtrate generally only left a leaf bone, whereas in the larvae given culture filtrate there were still leaves remaining (Figures 7 and 8). Thus, the culture filtrate could reduce the appetite of *S. litura* larvae.

The larvae *S. litura* has given culture filtrate besides being lazier to eat showed lazier and lazier to move. After 2 and 3 days being given the culture filtrate, the larvae began to change the color of integument which was previously bright green to become dull and the larvae were discovered to be dead. Four days after being given the culture filtrate, the larvae increasingly began to shrink and die. The dead larvae were wrinkled, dry, black, and odorless. The larvae not given the culture filtrate had larger body sizes and the integument colors were greener, and brighter (Figures 9 and 10).

Table 3. Mortality of larval *Spodoptera litura* after being infested with *Beauveria bassiana* culture filtrates

Isolate codes	Mortality ± SE*(%)	LT ₅₀ (days)± SE
BJgTs	98.00 ± 1.41 ^b	6.35 ± 0.49
BSmMs	50.00 ± 32.53 ^{ab}	11.57 ± 4.08
BswTd1	49.33 ± 15.36 ^{ab}	12.03 ± 3.52
BswTd2	94.67 ± 4.35 ^{ab}	5.92 ± 0.46
BswTd3	73.33 ± 18.51 ^{ab}	8.86 ± 1.78
BswTd4	26.00 ± 1.41 ^{ab}	16.07 ± 0.97
BPdR	86.67 ± 3.93 ^{ab}	8.17 ± 0.61
BKbTp	62.67 ± 10.39 ^{ab}	10.71 ± 0.91
BKKPp2	58.00 ± 7.07 ^{ab}	10.68 ± 0.34
TS1d3	10.67 ± 2.88 ^a	17.45 ± 0.76
BTmPe	46.67 ± 17.01 ^{ab}	14.32 ± 3.72
BTmTr	29.33 ± 9.68 ^{ab}	14.31 ± 1.79
TS1d2	88.00 ± 8.22 ^{ab}	7.87 ± 1.11
BTmTs	96.00 ± 0.00 ^{ab}	7.78 ± 0.09
BLePd2	70.00 ± 21.21 ^{ab}	9.59 ± 1.43
BTmKt	80.00 ± 14.73 ^{ab}	8.80 ± 1.41
BPcMs	81.33 ± 9.49 ^{ab}	7.87 ± 1.32
BMkMs	72.00 ± 17.99 ^{ab}	10.82 ± 2.46
BTmGa	66.67 ± 22.34 ^{ab}	12.27 ± 3.60
BTmSr	21.33 ± 9.30 ^{ab}	16.80 ± 2.67
BPcPd2	18.67 ± 7.85 ^{ab}	9.43 ± 3.86
BTmSo	80.00 ± 14.14 ^{ab}	7.55 ± 1.35
BTmMa	52.00 ± 19.69 ^{ab}	11.09 ± 1.62
BTmPe	70.00 ± 18.38 ^{ab}	9.30 ± 1.94
BLePd	94.00 ± 4.24 ^{ab}	9.30 ± 0.45
ANOVA F-value	2.09 [*]	1.20 ^{ns}
P value (0.05)	0.02	0.30
Tukey's HSD test	67.09	-

Note: ns = not significantly different; * = significantly different; values within a column (the data of each isolate) followed by the same letters were not significantly different at P < 0.05 according to Tukey's HSD test. Original data were transformed using Arcsin transformation prior to statistical analysis

Table 4. Mortality of larval *Spodoptera litura* after being infested with *Metarhizium anisopliae* culture filtrates

Isolate codes	Mortality ± SE*(%)	LT ₅₀ (days)± SE
MPdB	85.33 ± 14.67 ^b	8.09 ± 1.88 ^a
MPdR1	57.33 ± 19.37 ^{ab}	11.87 ± 2.56 ^a
MPdR2	65.33 ± 13.33 ^{ab}	10.43 ± 0.51 ^a
MPdPe	42.67 ± 9.61 ^{ab}	12.67 ± 0.89 ^a
MJgMs1	53.33 ± 23.25 ^{ab}	13.67 ± 4.52 ^a
MJgMs2	25.33 ± 3.53 ^{ab}	14.50 ± 0.56 ^a
MPdMs1	73.33 ± 10.41 ^{ab}	9.90 ± 0.97 ^a
MPdMs2	84.00 ± 9.24 ^{ab}	8.70 ± 1.26 ^a
MPdMs3	68.00 ± 16.17 ^{ab}	10.08 ± 0.86 ^a
MJgKeTs	42.67 ± 10.67 ^{ab}	12.68 ± 1.05 ^a
MPdMs4	16.00 ± 10.07 ^{ab}	53.70 ± 23.11 ^b
MJgTs2	46.67 ± 13.92 ^{ab}	14.29 ± 2.23 ^a
MKbTp1	84.00 ± 10.07 ^{ab}	8.69 ± 0.95 ^a
MSwTp1	56.00 ± 12.22 ^{ab}	14.64 ± 3.24 ^a
MSwTp2	74.67 ± 13.92 ^{ab}	10.62 ± 1.39 ^a
MSwTp3	61.33 ± 16.38 ^{ab}	10.91 ± 1.99 ^a
MSwTp4	61.33 ± 23.13 ^{ab}	14.60 ± 5.54 ^a
MKKPp1	44.00 ± 10.07 ^{ab}	12.33 ± 0.90 ^a
MKbTp2	96.00 ± 4.00 ^b	7.36 ± 0.62 ^a
MagPd	33.33 ± 3.53 ^{ab}	12.98 ± 0.23 ^a
ANOVA F-value	2.39 [*]	50.13 [*]
P value (0.05)	0.01	0.00
Tukey's HSD test	53.35	2.65

Note: ns = not significantly different; * = significantly different; values within a column (the data of each isolate) followed by the same letters were not significantly different at P < 0.05 according to Tukey's HSD test. Original data were transformed using Arcsin transformation prior to statistical analysis




Figure 3. *Beauveria bassiana* isolates cultured in PDB: BJgTs (A), BSmMs (B), BSwTd1 (C), BSwTd2 (D), BSwTd3 (E), BSwTd4 (F), BPdR (G), BKbTp (H), BKKPp2 (I), Ts1d3 (J), BTmPc (K), BTmTr (L), (M) Ts1d2 (M), BTmTs (N), BlePd2 (O), BTmKt (P), BPCmS (Q), BMkMs (R), BtmGa (S), BTmSr (T), BPCPd2 (U), BtmSo (V), BtmMa (W), BtmPe (X), Blepd (Y)

Discussion

The *B. bassiana* culture filtrate, particularly the isolates coded BSwTd2, were the most toxic compared to other isolates with LT_{50} of only 5.91 days. The BSwTd2 isolate was from peat soils overgrown with oil palm plants. For *M. anisopliae*, the most toxic MKbTp2 isolate was from the cabbage in highland. The larvae began to die after 3x24 hours after the treatment either due to *B. bassiana* or *M. anisopliae*. The larvae death resulted from the 6 weeks of incubation of entomopathogenic fungi causing the toxic culture filtrate. According to Pinnamaneni et al. (2010), the culture of filtrates of the toxic *B. bassiana* was caused by the incubation in liquid culture media in which the fungus produced chitinolytic exochitinase enzymes so that when they entered the body the insect was able to degrade the cuticle. In addition, the fungal broth during the incubation could produce protease enzymes (Qazi 2008).

The time of death of the larvae by this culture filtrate was faster than that of by conidia. El Husseini (2019) stated that the death of *Spodoptera* larvae by *M. anisopliae* conidia began to occur at the fourth day (4x24 hours) or the fifth day after the treatment (post-treatment), whereas in this study the *M. anisopliae* culture filtrate began to kill *S. litura* at the third day after the treatment. The lethal time difference between these was due to the mode of action of fungal conidia being different from the culture filtrate. To kill the host insect, the fungus conidia got contact first with it and then produced toxins to kill the host, while the culture filtrate directly killed the host insect. El-Ghany (2015) states that entomopathogenic fungus conidia kill its host insect in two stages: parasitic and saprophytic phases. The parasitic phase began with the fungal conidia attaching to the host insect cuticle (Augustyniuk-Kram and Kram 2012). Then, if the humidity was high, the conidia started to germinate on the host cuticle (El-Ghany 2015). The

conidia germinating by forming germ tubes continued to grow sticking out to find a soft integument position to facilitate entry into the cuticle (Fernandes et al. 2007). The infection occurs when the germ tubes are able to penetrate insect cuticles and the ability to infect them is a determining factor for the fungus virulence (Altre and Vandenberg 2001). After the germ tubes penetrated the cuticle and reached the hemocoel, and then they produced specific infection hyphae originating at appressoria (El-Ghany 2015). Furthermore, the hyphae spread to the hemolymph and developed to produce blastospores, and the blastospores produced fungal cytotoxic, for example, destruxins by *M. anisopliae* which killed the host insects

(Mancillas-Paredes et al. 2019), but the death of these host insects was not caused only by the toxin but also due to the mechanical damage by the penetration of fungi into the body of insects (El-Ghany 2015).

After the host insect died, it entered into the saprophytic phase which was influenced by favorable environmental conditions (Peña-Peña et al. 2015). In the body of the dead insect, the fungus formed mycelia and hyphae which continued to grow covering the body of the host insect, and then the hyphae formed conidiogenous cells and the conidia were produced by utilizing the nutrients/fluids of the host insect and finally, the infection process was complete (El-Ghany 2015).



Figure 4. *Beauveria bassiana* culture filtrate: BJgTs (A), BSmMs (B), BSwTd1 (C), BSwTd2 (D), BSwTd3 (E), BSwTd4 (F), BPcR (G), BKbTp (H), BKKPp2 (I), Ts1d3 (J), BTmPc (K), BTmTr (L), (M) Ts1d2 (M), BTmTs (N), BlePd2 (O), BTmKt (P), BPCmS (Q), BMkMs (R), BtmGa (S), BTmSr (T), BPcPd2 (U), BtmSo (V), BtmMa (W), BtmPe (X), Blepd (Y)

In contrast to the mode of action of fungal conidia, a fungal culture filtrate directly kills the host insect due to the toxins produced by fungal broth so that the time needed to kill the host insect is shorter. The entomopathogenic fungus cultured in liquid media can produce conidia in the form of blastospores (Mascarin et al. 2015) which can produce toxins (Mascarin and Jaronski 2016). In this study, the result of calculating conidia density after being incubated for 6 weeks was generally more than 1×10^9 conidia mL^{-1} . Fungal conidia and propagules were separated from the filtrate through two stages of filtration to produce toxic fungal culture filtrate. The *B. bassiana* and *M. anisopliae* culture filtrates were able to kill up to 94.67% and 96% of the larvae of *S. litura*, respectively. The short time of death by this culture filtrate resulted from the death of the direct host insect by consuming feed (chili leaves) moistened with the culture filtrate. Soesanto et al. (2019) stated that entomopathogenic fungal culture filtrate contains secondary metabolites. *B. bassiana* culture filtrate

containing the secondary metabolites can weaken the host insect's immune system (Zibae et al. 2011). The secondary metabolites produce toxins, for example, destruxins and efraptens (Zibae et al. 2009). *B. bassiana* also produces a protease enzyme that can kill host insects by dissolving their body proteins (Mancillas-Paredes et al. 2019).

The color of fungal broth and culture of *B. bassiana* and *M. anisopliae* varied between isolates. The darker color of fungal broths and culture filtrates tend to cause higher mortality. In line with the results obtained by Ayudya et al. (2019), the darker culture filtrates tended to be more toxic than the light-colored ones. The darker culture filtrates show higher production of secondary metabolites (Luo et al. 2017) and the activity of extracellular enzymes (Khachatourians et al. 2007) and the enzymes produced such as proteases are able to dissolve integument insects that result in death (Mancillas-Paredes et al. 2019).



Figure 5. *Metarhizium anisopliae* isolates cultured in PDB: MPdB (A), MPdR1 (B), MPdR2 (C), MpdPe (D), MJgMs1 (E), MJgMs2 (F), MPdMs1 (G), MPdMs2 (H), MPdMs3 (I), MjgKeTs (J), MPdMs4 (K), MJgTs2 (L), MKbTp1 (M), MSwTp1 (N), MSwTp2 (O), MSwTp3 (P), MSwTp4 (Q), MKKp1 (R), MKbTp2 (S), MagPd (T)




Figure 6. *Metarhizium anisopliae* culture filtrate: MPdB (A), MPdR1 (B), MPdR2 (C), MpdPe (D), MJgMs1 (E), MJgMs2 (F), MPdMs1 (G), MPdMs2 (H), MPdMs3 (I), MjgKeTs (J), MPdMs4 (K), MJgTs2 (L), MKbTp1 (M), MSwTp1 (N), MSwTp2 (O), MSwTp3 (P), MSwTp4 (Q), MKKp1 (R), MKbTp2 (S), MagPd (T)




Figure 7. The leaf damage: control (A) and larval *Spodoptera litura* applied with *Beauveria bassiana* culture filtrate (B)



Figure 8. The leaf damage: control (A) and larval *Spodoptera litura* applied with *Metarhizium anisopliae* culture filtrate (B)



Figure 9. The healthy (A) and dead larvae (B) of *Spodoptera litura* caused by *Beauveria bassiana* culture filtrate



Figure 10. The healthy (A) and dead larvae (B) of *Spodoptera litura* caused by *Metarhizium anisopliae* culture filtrate

The symptoms of sick host insects by culture filtrate in this study differed from the deaths by fungal conidia. The insects getting sick by the culture filtrate were not covered by mycelia or conidia on the integument. However, they got sick because of the fungal conidia infection which was generally covered in mycelia (Sumikarsih et al. 2019). The body shape of the sick *S. litura* larvae due to the culture filtrate was shrunk, wrinkled, dry, and odorless. Similar to the results of the study of Ayudya et al. (2019), the *S. litura* larvae treated with *B. bassiana* culture filtrate decreased their body weight and became dried, odorless, and the integument was not overgrown with fungal mycelia because in the culture filtrate it no longer contained conidia but contained toxic compounds for host insects. Therefore, the mortality of *S. litura* larvae in this study was caused by the culture filtrate containing toxic compounds. Zibae et al. (2011) state that toxic compounds in the culture filtrate are secondary metabolites. Zibae et al. (2009) stated that the secondary metabolites are easier to apply in the field because they can be integrated with other control techniques including synthetic insecticides. Consequently, the culture filtrate which is toxic due to the high content of secondary metabolites has the potential to be further developed for the active ingredient of mycoinsecticide.

The most toxic isolates of the culture filtrate were BSwtD2 of *B. bassiana* and MKbTp2 of *M. anisopliae* resulting in the mortality of above 90%. The two isolates that were found each came from the lowlands and highlands so that in the future there is an opportunity for the isolates to be developed and applied in low and highland ecosystems. Entomopathogenic fungi that produce toxic filtrate cultures are capable of killing their hosts in a short time, so that they have the potential to be active ingredients in the development of mycoinsecticides technology.

ACKNOWLEDGEMENTS

Thank you to the sponsors of the donors of this research, the Individual National Strategic Research (PSNI) program for the 2018 fiscal year based on the Research Contracts of the Directorate of Research and Community Service (DRPM), Directorate General of Research and Development Strengthening, Ministry of Technology Research and Higher Education, Contract number: 093/SP2H/LT/DRPM/IV/2018 chaired by SH.

REFERENCES

- Altre JA, Vandenberg JD. 2001. Factors influencing the infectivity of isolates of *Pacilomyces fumosoroseus* against diamondback moth, *Plutella xylostella*. *J Invertebr Pathol* 78: 31-36. DOI: 10.1006/jjpa.2001.5043
- Augustyniuk-Kram A, Kram KJ. 2012. Entomopathogenic fungi as an important natural regulator of insect outbreaks in forests review. *Forest Ecosystem, More than Just Trees*. In: Blanco JA. (ed.). In Tech, Croatia. DOI: 10.5772/30596.
- Ayudya D, Herlinda S, Suwandi S. 2019. Insecticidal activity of culture filtrates from liquid medium of *Beauveria bassiana* isolates from South Sumatra (Indonesia) wetland soil against larvae of *Spodoptera litura*. *Biodiversitas* 20: 2101-2109. DOI: 10.13057/biodiv/d200802
- Bandani AR, Khambay BPS, Faull L, Newton R, Deadman M, Butt TM. 2000. Production of efrapeptins by *Tolypocladium* species and evaluation of their insecticidal and antimicrobial properties. *Mycol Res* 104: 537-544. DOI: 10.1017/S0953756299001859.
- Bueno RC, Bueno AF, Moscardi F, Parra JR, Hoffmanncampo CB. 2011. Lepidopteran larva consumption of soybean foliage: basis for developing multiple-species economic thresholds for pest management decisions. *Pest Manag Sci* 67 (2): 170-174.
- Darlan N, Pradiko I, Winarna, Siregar HH. 2016. Effect of el niño 2015 on oil palm performance in central and Southern Sumatera. *J Tanah dan Iklim* 40: 113-120. DOI: 10.2017/jti.v40n2.2016.113-120. [Indonesian]
- El-Ghany TMA. 2015. Entomopathogenic Fungi and their Role in Biological Control. *Biology Department Faculty of Science Jazan University KSA, Cairo*. DOI:10.4172/978-1-63278-065-2-66
- El-Husseini MMM. 2019. Efficacy of the entomopathogenic fungus, *Metarhizium anisopliae* (Metsch.), against larvae of the cotton leafworm, *Spodoptera litoralis* (Boisd.) (Lepidoptera: Noctuidae), under laboratory conditions. *Egypt J Biol Pest Control* 29: 4-6. DOI: 10.1186/s41938-019-0156-2
- Fattah A, Syam S, Daud ID, Dewi VS, Rahman A. 2018. The intensity of leaf damage caused by attack of *Spodoptera litura* F and seed yield on some soybean varieties in South Sulawesi Indonesia. *Sci Res J* 6: 55-60.
- Fernandes EKK, Rangel DEN, Moraes AM, Bittencourt VREP, Roberts DW. 2007. Variability in tolerance to UV-B radiation among *Beauveria* spp. isolates. *J Invertebr Pathol* 96: 237-243. DOI: 10.1016/j.jip.2007.05.007
- Garrido-Jurado J, Torrent J, Barrón V, Corpas A, Quesada-Moraga E. 2011. Soil properties affect the availability, movement, and virulence of entomopathogenic fungi conidia against puparia of *Ceratitis capitata* (Diptera: Tephritidae). *Biol Control* 58: 277-285. DOI: 10.1016/j.biocontrol.2011.05.017
- Herlinda S, Yudha S, Thalib R, Khodijah, Suwandi, Lakitan B, Verawaty M. 2018a. Species richness and abundance of spiders inhabiting rice in fresh swamps and tidal lowlands in South Sumatera, Indonesia. *J ISSAAS* 24: 82-93.
- Herlinda S, Rizkie L, Suwandi, Susilawati, Lakitan B, Verawaty M, Hasbi. 2018b. Effects of high temperature and ultraviolet-C irradiance on conidial viability and density of *Beauveria bassiana* and *Metarhizium anisopliae* isolated from soils of lowland ecosystems in Indonesia. *Eurasian J Anal Chem* 13: 209-216.
- Johari A, Herlinda S, Pujiastuti Y, Irsan C, Sartiami, D. 2016. Morphological and genetic variation of *Thrips parvispinus* (Thysanoptera: Thripidae) in chili plantation (*Capsicum annum* L.) in the lowland and highland of Jambi Province, Indonesia. *Am J Biol Sci* 2: 17-21. DOI: 10.11648/j.abio.s.2014020601.14
- Juhriah, Azrai M, Tambaru E, Rahayu JE. 2019. Phenotypic characteristics and grouping of hybrid waxy corn *Zea mays* L. *J Ilmu Alam dan Lingkungan* 10: 51-60. DOI: 10.20956/jal.v10i1.6624 [Indonesian]
- Karenina T, Herlinda S, Irsan C, Pujiastuti Y. 2019. Abundance and species diversity of predatory arthropods inhabiting rice of refuge habitats and synthetic insecticide application in freshwater swamps in South Sumatra, Indonesia. *Biodiversitas* 20 (8): 2375-2387. DOI: 10.13057/biodiv/d200836
- Kartika K, Lakitan B, Wijaya A, Kadir S, Widuri L.I, Siaga E. 2018. Effects of particle size and application rate of rice-husk biochar on chemical properties of tropical wetland soil, rice growth and yield. *Aust J Crop Sci* 12: 817-826. DOI: 10.21475/ajcs.18.12.05.PNE1043
- Khachatourians GG, Qazi SS, Reid WR. 2007. Biochemistry and molecular biology of entomopathogenic fungi. In: Brakhage AA, Zipfer PF (eds.). *The Mycota VI: Human and animal relationships*. Springer-Verlag, Berlin.
- Liu SF, Wang GJ, Nong XQ, Liu B, Wang M, Li SL, Cao GC, Zhang ZH. 2017. Entomopathogen *Metarhizium anisopliae* promotes the early development of Peanut Root. *Plant Protect Sci* 53: 101-107. DOI: 10.17221/49/2016-PPS
- Luo Z, Ren H, Mousa JJ, Rangel DEN, Zhang Y, Bruner SD, Keyhani NO. 2017. The PacC transcription factor regulates secondary metabolite production and stress response, but has only minor effects on virulence in the insect pathogenic fungus *Beauveria bassiana*. *Environ Microbiol* 19: 788-802. DOI: 10.1111/1462-2920.13648
- Mancillas-Paredes JM, Hernández-Sánchez H, Jaramillo-Flores ME, García-Gutiérrez C. 2019. Proteases and chitinases induced in

- Beauveria bassiana* during infection by *Zabrotes subfasciatus*. Southwestern Entomol 44: 125-137. DOI: 10.3958/059.044.0114
- Maqsood S, Sabri, MA, Ali A, Abbas M, Aziz A. 2017. Comparative toxicity of some insecticides against army worm, *Spodoptera litura* L. (Lepidoptera: Noctuidae) under laboratory conditions. J Entomol Zool Stud 5: 770-773.
- Maryanto M, Sukiyono K, Priyono B. 2018. Technical efficiency analysis and determinants in potato farming (*Solanum tuberosum* L.) in Pagar Alam City, South Sumatra Province. Agraris 4: 1-8. DOI: 10.18196/agr.4154 [Indonesian]
- Mascarin GM, Jackson MA, Behle RW, Kobori NN, Júnior ÍD. 2016. Improved shelf life of dried *Beauveria bassiana* blastospores using convective drying and active packaging processes. Appl Microbiol Biotechnol 100: 8359-8370. DOI: 10.1007/s00253-016-7597-2
- Mascarin GM, Jackson MA, Kobori NN, Behle RW, Delalibera Júnior Í. 2015. Liquid culture fermentation for rapid production of desiccation-tolerant blastospores of *Beauveria bassiana* and *Isaria fumosorosea* strains. J Invertebr Pathol 127: 11-20. DOI: 10.1016/j.jip.2014.12.001
- Munir J, Herman W. 2019. Various physical and soil chemical properties phenomena support food security in West Sumatera. Ziraa'ah 44: 146-153. DOI: 10.31602/zmp.v44i2.1736. [Indonesia]
- Nagal G, Verma K, Rathore L. 2016. Management of *Spodoptera litura* (Fabricius) through some novel insecticides and biopesticides on bell pepper under polyhouse environment. Adv Life Sci 5: 1081-1084.
- Nasution K, Respatjarti. 2019. Yield trial on eight potential lines F7 generation of chili pepper (*Capiscum Annuum* L.) compact type in Low Land. J Produksi Tanaman 7: 464-473. [Indonesian]
- Oreste M, Baser N, Bubici G, Tarasco E. 2015. Effect of *Beauveria bassiana* strains on the *Ceratitidis capitata* - *Psytalia concolor* system. Bull Insectol 68: 265-272.
- Peña-Peña AJ, Santillán-Galicia MT, Hernández-López J, Guzmán-Franco AW. 2015. *Metarhizium pingshaense* applied as a seed treatment induces fungal infection in larvae of the white grub *Anomala cincta*. J Invertebr Pathol 130: 9-12. DOI: 10.1016/j.jip.2015.06.010
- Pinnamaneni R., Kalidas P, Rao KRSS. 2010. Cloning and Expression of Bbchit1 gene of *Beauveria bassiana*. Open Entomol J 4: 30-35. DOI: 10.2174/1874407901004010030
- Prabawati G, Herlinda S, Pujiastuti Y. 2019. The abundance of canopy arthropods in South Sumatra (Indonesia) freshwater swamp main and ratooned rice applied with bioinsecticides and synthetic insecticide. Biodiversitas 20: 2921-2930. DOI: 10.13057/biodiv/d201021
- Qazi SS, Khachatourians GG. 2008. Addition of exogenous carbon and nitrogen sources to aphid exuviae modulates synthesis of proteases and chitinase by germinating conidia of *Beauveria bassiana*. Arch Microbiol 189 (6): 589-596.
- Safitri A, Herlinda S, Setiawan, A. 2018. Entomopathogenic fungi of soils of freshwater swamps, tidal lowlands, peatlands, and highlands of South Sumatra, Indonesia. Biodiversitas 19: 2365-2373. DOI: 10.13057/biodiv/d190647
- Sagrim M, Sumule AI, Iyai DA, Baransano M. 2017. Prime potency of agriculture commodities on highland of Arfak Mountains Regency, Papua Barat. J Ilmu Pertanian Indonesia 22: 141-146. [Indonesian]. DOI: 10.18343/jipi.22.3.141
- Situmorang S, Haryono D, Soelaiman A. 2019. Analysis of business performance and welfare of vegetable farmers in Pagar Alam City South Sumatera Province. Indon J Socio Econ 1: 84-92. [Indonesian]
- Soesanto L, Mugiastuti E, Manan A. 2019. Raw secondary metabolites application of two *Trichoderma harzianum* isolates towards vascular streak dieback on cocoa seedlings. Pelita Perkebunan 35: 22-32. DOI: 10.22302/iccj.jur.pelitaperkebunan.v35i1.346
- Sumikarsih E, Herlinda S, Pujiastuti Y. 2019. Conidial density and viability of *Beauveria bassiana* isolate from java and Sumatra and their virulence against *Nilaparvata lugens* at different temperatures. Agrivita 1: 335-349. DOI: 10.17503/agrivita.v41i2.2105
- Turnip A, Rantuboi DYP, Lantang D. 2018. Effectivity test of *Beauveria bassiana* fungus and application time of *Spodoptera litura* in mustard greens (*Brassica juncea*). J Biologi Papua 10: 26-31. DOI: 10.31957/jbp.131. [Indonesian]
- Vijaya M, Rani PU. 2017. Defensive responses in *Capiscum annuum* (L) plants, induced due to the feeding by different larval instars of *Spodoptera litura* (F). Arthropod-Plant Interact 11: 193-202.
- Zamhari A, Sitorus S, Pravitasari A. 2017. Analysis of the competitive commodities and its direction of development plan in Pagar Alam City, South Sumatera Province. Tata Loka 19: 218-229. DOI: 10.14710/tataloka.19.3.218-229. [Indonesian]
- Zibae A, Bandani AR, Talaei-Hassanlouei R, Malagoli D. 2011. Cellular Immune Reactions of the Sunn Pest, *Eurygaster integriceps*, to the Entomopathogenic Fungus, *Beauveria bassiana* and its secondary metabolites. J Insect Sci 11: 1-16. DOI: 10.1673/031.011.13801
- Zibae A, Bandani AR, Tork M. 2009. Effect of the entomopathogenic fungus, *Beauveria bassiana*, and its secondary metabolite on detoxifying enzyme activities and acetylcholinesterase (AChE) of the Sunn pest, *Eurygaster integriceps* (Heteroptera: Scutellaridae). Biocontrol Sci Technol 19: 485-498. DOI: 10.1080/09583150902847127.

Toxicity of entomopathogenic fungal culture filtrate of lowland and highland soil of South Sumatra (Indonesia) against *Spodoptera litura* larvae

ORIGINALITY REPORT

7%

SIMILARITY INDEX

MATCH ALL SOURCES (ONLY SELECTED SOURCE PRINTED)

★fr.scribd.com
Internet

4%

EXCLUDE QUOTES ON
EXCLUDE BIBLIOGRAPHY ON

EXCLUDE MATCHES < 1%

**FORMAT PENILAIAN (VALIDASI & PEER REVIEW)
LEMBAR
HASIL PENILAIAN SEJAWAT SEBIDANG ATAU *PEER REVIEW*
KARYA ILMIAH : JURNAL ILMIAH**

Jurnal Artikel Ilmiah : Toxicity of entomopathogenic fungal culture filtrate of lowland and highland soil of South Sumatra (Indonesia) against Spodoptera litura larvae

Penulis Artikel Ilmiah : Arum Setiawan

Identitas Jurnal Artikel Ilmiah : a. Nama Jurnal : Biodiversitas
b. Nomor/Volume/Hal : 5/21/1839-1849
c. Edisi (bulan/tahun) : Mei 2020
d. Penerbit : Jurusan Biologi FMIPA UNS
e. Jumlah Halaman : 11

Kategori Publikasi Jurnal Ilmiah : Jurnal Ilmiah Internasional Bereputasi
(beri \surd pada kategori yang tepat) Jurnal Ilmiah Internasional
 Jurnal Ilmiah Nasional Terakreditasi S1, S2
 Jurnal Ilmiah Nasional Terakreditasi S3, S4
 Jurnal Ilmiah Nasional Tidak Terakreditasi

I. Hasil Penilaian Validasi :

No.	ASPEK	URAIAN/KOMENTAR PENILAIAN
1.	Indikasi Plagiasi	7 %
2.	Linearitas	Paper kurang linier dengan bidang biologi konservasi

II. Hasil Penilaian *Peer Review* :

Komponen Yang Dinilai	Nilai Maksimal Jurnal Ilmiah (isikan di kolom yang sesuai)					Nilai Akhir Yang Diperoleh
	Internasional Bereputasi (Maks 40)	Internasional (Maks 20)	Nasional Terakreditasi S1, S2 Maks 25	Nasional Terakreditasi S3, S4 Maks 20	Nasional tidak Terakreditasi (maks 10)	
Kelengkapan dan Kesesuaian unsur isi jurnal (10%)	4					3
Ruang lingkup dan kedalaman pembahasan (30%)	12					11
Kecukupan dan Kemutakhiran data/informasi dan metodologi (30%)	12					12
Kelengkapan unsur dan kualitas penerbit (30%)	12					12
Total = (100%)	40					38
Kontribusi Pengusul (Penulis Pertama /Anggota Utama)	Anggota Utama $(0,4 \times 38) / 9 = 1,69$					1,69
KOMENTAR/ULASAN <i>PEER REVIEW</i>						
• Kelengkapan dan Kesesuaian Unsur:	Paper terkait toksisitas entomopathogenic kultur jamur. Isi paper sudah memenuhi kaidah-kaidah karya ilmiah, namun kurang sesuai dengan bidang biologi konservasi.					
• Ruang Lingkup dan Kedalaman Pembahasan:	Hasil penelitian dibahas cukup komprehensif dengan penyampaian perbandingan dari temuan-temuan penelitian lainnya dan teori terkait. Referensi yang diacu dalam pembahasan sudah cukup update untuk bidang kajian ini.					
• Kecukupan & Kemutakhiran Data & Metodologi:	Data-data hasil penelitian cukup baik dan didukung peta lokasi sampling dan gambar yang ditampilkan menarik. Data didapatkan dengan menggunakan metode yang standard..					
• Kelengkapan Unsur & Kualitas Penerbit:	Penerbit Jurusan Biologi FMIPA UNS berkualitas baik, tidak termasuk predatory publisher, dan jurnal terindeks di scopus Q3.					

Surabaya, 13 Mei 2020
Penilai 1



Prof. Hery Purnobasuki, M.Si., Ph.D.
NIP 196705071991021001
Unit Kerja : Jurusan Biologi FST Unair
Bidang Ilmu : Biologi
Jabatan/Pangkat : Guru Besar/ Pembina Utama Madya

FORMAT PENILAIAN (VALIDASI & PEER REVIEW)

LEMBAR

HASIL PENILAIAN SEJAWAT SEBIDANG ATAU PEER REVIEW

KARYA ILMIAH : JURNAL ILMIAH

Jurnal Artikel Ilmiah : Toxicity of entomopathogenic fungal culture filtrate of lowland and highland soil of South Sumatra (Indonesia) against *Spodoptera litura* larvae

Penulis Artikel Ilmiah : Arum Setiawan

Identitas Jurnal Artikel Ilmiah :

a. Nama Jurnal : Biodiversitas

b. Nomor/Volume/Hal : 5/21/1839-1849

c. Edisi (bulan/tahun) : Mei 2020

d. Penerbit : Jurusan Biologi FMIPA UNS

e. Jumlah Halaman : 11

Kategori Publikasi Jurnal Ilmiah (beri V pada kategori yang tepat) :

Jurnal Ilmiah Internasional Bereputasi

Jurnal Ilmiah Internasional

Jurnal Ilmiah Nasional Terakreditasi S1, S2

Jurnal Ilmiah Nasional Terakreditasi S3, S4

Jurnal Ilmiah Nasional Tidak Terakreditasi

I. Hasil Penilaian Validasi :

No.	ASPEK	URAIAN/KOMENTAR PENILAIAN
1.	Indikasi Plagiasi	7 %
2.	Linearitas	V

II. Hasil Penilaian Peer Review :

Komponen Yang Dinilai	Nilai Maksimal Jurnal Ilmiah (isikan di kolom yang sesuai)					Nilai Akhir Yang Diperoleh
	Internasional Bereputasi (Maks 40)	Internasional (Maks 20)	Nasional Terakreditasi S1, S2 Maks 25	Nasional Terakreditasi S3, S4 Maks 20	Nasional tidak Terakreditasi (maks 10)	
Kelengkapan dan Kesesuaian unsur isi jurnal (10%)	4					4
Ruang lingkup dan kedalaman pembahasan (30%)	12					10
Kecukupan dan Kemutakhiran data/informasi dan metodologi (30%)	12					10
Kelengkapan unsur dan kualitas penerbit (30%)	12					12
Total = (100%)	40					36
Kontribusi Pengusul (Penulis Pertama /Anggota Utama)	Biodiversitas Vol. 21(5) Mei 2020. Q3 SJR 0,27. Penulis ke 7 dari 10. Nilai maksimal 90 %. Nilai pengusul $(0,4 \times 0,9 \times 40)/9 = 1,6$					1,6
KOMENTAR/ULASAN PEER REVIEW						
• Kelengkapan dan Kesesuaian Unsur:	Format lengkap dari abstrak sampai dengan referensi.					
• Ruang Lingkup dan Kedalaman Pembahasan:	Masih dalam ruang lingkup tetapi pembahasan kurang detail.					
• Kecukupan & Kemutakhiran Data & Metodologi:	Data layak dan baru. Metode cukup baik dan dapat dipahami.					
• Kelengkapan Unsur & Kualitas Penerbit:	Penerbit cukup berkualitas.					

Yogyakarta, 16 Juni 2020

Penilai 2

tanda tangan

Prof. Dr. Suwarno Hadisusanto

NIP 195411161983031002

Unit Kerja : Fakultas Biologi UGM

Bidang Ilmu : Biologi/Ekologi

Jabatan/Pangkat : Guru Besar/ Pembina Utama Madya