

# Plasma Levels of the Engulfment and Cell Motility Protein-1 are Associated with Kidney Damage in Diabetic Nephropathy: A Single-Center Pilot Study in Indonesia Population

*by* Zulkhair Ali

---

**Submission date:** 05-Apr-2021 01:48PM (UTC+0700)

**Submission ID:** 1550779749

**File name:** Plasma\_Levels\_of\_the\_Engulfment\_and\_Cell\_Motility\_Protein-1.pdf (386.15K)

**Word count:** 3876

**Character count:** 20736



# Plasma Levels of the Engulfment and Cell Motility Protein-1 are Associated with Kidney Damage in Diabetic Nephropathy: A Single-Center Pilot Study in Indonesia Population

Elfiani Elfiani<sup>1,2\*</sup>, Ellyza Nasrul<sup>1</sup>, Yanwirasti Yanwirasti<sup>1</sup>, Zulkhair Ali<sup>3</sup>, Anggelia Puspasari<sup>4</sup>

<sup>1</sup>Doctoral Degree of Biomedical Sciences, Faculty of Medicine, Universitas Andalas, Padang, Indonesia; <sup>2</sup>Department of Internal Medicine, Faculty of Medicine and Health Science, Universitas Jambi, Jambi, Indonesia; <sup>3</sup>Department of Internal Medicine, Faculty of Medicine, Universitas Sriwijaya, Palembang, Indonesia; <sup>4</sup>Department of Biochemistry, Faculty of Medicine and Health Science, Universitas Jambi, Jambi, Indonesia

## Abstract

**Citation:** Elfiani E, Nasrul E, Yanwirasti Y, Ali Z, Puspasari A. Plasma Levels of the Engulfment and Cell Motility Protein-1 are Associated with Kidney Damage in Diabetic Nephropathy: A Single-Center Pilot Study in Indonesia. *Open Access Macedonian Journal of Medical Sciences*. 2020 Jul 25; 8(A):418-422. <https://doi.org/10.3889/oamjms.2020.4276>  
**Key words:** Engulfment and cell motility protein-1; Kidney damage; Type 2 diabetes mellitus; Diabetic nephropathy; Albumin to creatinine ratio; Glomerular filtration rate  
**\*Correspondence:** Elfiani Elfiani, Department of Internal Medicine, Faculty of Medicine and Health Science, Universitas Jambi, Jambi, Indonesia. Tel.: +62811745164.  
 E-mail: [elfiani@gmail.com](mailto:elfiani@gmail.com)  
**Received:** 28-Jan-2020  
**Revised:** 22-Apr-2020  
**Accepted:** 07-Jul-2020  
**Copyright:** © 2020 Elfiani Elfiani, Ellyza Nasrul, Yanwirasti Yanwirasti, Zulkhair Ali, Anggelia Puspasari  
**Funding:** This research was partially funded by Universitas Jambi research annually funding for 2018 and 2019.  
**Competing Interests:** The authors have declared that no competing interests exist.  
**Open Access:** This is an open access article distributed under the terms of the [Creative Commons Attribution-NonCommercial 4.0 International License \(CC BY-NC 4.0\)](https://creativecommons.org/licenses/by-nc/4.0/)

**BACKGROUND:** Diabetic nephropathy (DN) is a chronic kidney disease with microvascular complications which influence a patient's quality of life and poses a potentially significant economic burden. The discovery of new biomarkers is needed to gain a better understanding and find additional treatment approaches beyond controlling plasma glucose and blood pressure. The engulfment and cell motility protein called engulfment and cell motility protein (ELMO)-1 is known to play a role in tissue remodeling and is associated with DN progression in animal models. Human studies correlating this protein with DN have been limited, although a genetic variation of ELMO-1 has been associated with DN.

**AIMS:** This research aimed to investigate the association and correlation of circulating ELMO-1 with the level of kidney damage in DN.

**METHODS:** The research design was cross-sectional. In this study, 60 patients with type 2 diabetes mellitus (T2DM) participated. Fasting blood pressure, plasma glucose, and renal function parameters were measured to confirm the diagnosis. Albumin to creatinine ratio (ACR) was used to assess the level of kidney damage and determines DN group.

**RESULTS:** Analyses shown the severity of albuminuria which is associated with plasma ELMO-1 when DN is evident. Furthermore, in all study participants, plasma ELMO-1 was correlated with increased proteinuria, ACR, and decline of glomerular filtration rate.

**CONCLUSIONS:** Our research findings provide support for ELMO-1 as a promising marker of kidney damage level in patients with T2DM, especially when DN has occurred. Larger sample sizes and multicenter research in the Indonesian population are needed to confirm these results.

## Introduction

The incidence of diabetic nephropathy (DN) has increased worldwide, especially in Asian populations. This condition leads to end-stage renal diseases which often decrease a patient's quality of life and cause potentially significant economic burden [1], [2]. Additional approaches are needed to prevent and treat DN beyond controlling hyperglycemia and high blood pressure. Research of new biomarkers is important to gain a better understanding of DN progression and further develop new treatment approaches to prevent the progression of DN [3], [4], [5].

Increased microalbuminuria and decreased glomerular filtration rate (GFR) are clinical markers of DN [6]. Tissue remodeling, including alteration of matrix deposition, podocytes, and endothelial injury, which increases cell apoptosis, may happen before clinical markers of DN are detected [7]. These events are

associated with alteration of some circulating protein expressions, which are promising as new biomarkers before the commonly used clinical markers of DN appear [3], [7], [8].

The engulfment and cell motility protein called engulfment and cell motility protein (ELMO)-1 is known to play a role in tissue remodeling. The exact role of this protein in human DN pathophysiology is still unknown, although some genetic association studies reported that the ELMO-1 gene is a strong candidate as a DN associated gene. Polymorphism in the intronic region of ELMO-1 was associated with DN in Asia and America populations. None have described any association of its polymorphisms or DN with systemic or local ELMO-1 expression, although translation studies in animal models have reported that the ELMO-1 protein does play a role in the progression of DN [9], [10], [11], [12].

Diabetic rat models which have different systemic levels of ELMO-1 shown higher ELMO-1 expression that was associated with severity of renal fibrosis and amount

of urinary albumin excretion through aggravated lipid peroxidation and increased tumor growth factor beta-1 (TGFβ-1) levels, which are both markers of extracellular matrix expansion [13]. Another animal translation study that used a genetically modified zebrafish model shown that ELMO-1 is an important factor for glomerular protection and renal cell survival through decreasing apoptosis, especially under diabetic conditions [14].

This research was conducted because studies in humans about ELMO-1 expression in DN are limited, and recent findings in animal models briefly describe the promising potential of ELMO-1 as a new biomarker which plays a role in DN progression. This research aimed to investigate the association and correlation of circulating ELMO-1 in the progression of DN in the Indonesian population for the 1<sup>st</sup> time.

## Materials and Methods

### Study design

This research was a cross-sectional study. A total of 60 patients with type II diabetes mellitus (T2DM) were recruited from the Jambi province who fulfilled eligible criteria for this research. Inclusion criteria were as follows: Patient with T2DM who was suffering from DM for at least 5 years, with an age range of 30–60 years old. We excluded patients with urinary infections based on clinical examination and routine urine analysis, history of other kidney diseases based on the medical record, pregnant women, and immunocompromised subjects. Patients' T2DM diagnosis was based on fasting plasma glucose >126 mg/dL and 2 h postprandial plasma glucose >200 mg/dL, as listed in medical records. In all patients, we measured albumin to creatinine ratio (ACR), GFR, and plasma creatinine as kidney function parameters and plasma ELMO-1. We also measured blood pressure and plasma glucose levels. All research subjects signed informed consent forms after receiving a detailed explanation about the study objectives and design. The research protocol was arranged based on the Helsinki Declaration and was approved by the research ethics committee of the Faculty of Medicine, Andalas University, with approval registration number 706/KEP/FK/2019.

### Kidney function parameters

Random spot urine samples as much as 30 mL were collected from the patients. Quantitative measurements of urine albumin were made with immunoturbidimetric. Urine creatinine was measured with enzymatic colorimetric. Then, the ACR was calculated based on the ratio of quantitative creatinine to albumin in random spot urine sampling. DN was diagnosed according to ACR, where DN was defined as ACR ≥30 mg/g. Furthermore, we also grouped degree of

kidney damage level according to ACR, with normal to mildly albuminuria as the A1 group (ACR <30 mg/g), the moderately increased albuminuria as the A2 group (ACR 30–300 mg/g), and the severely increased albuminuria as the A3 group (ACR >300 mg/g). GFR was calculated based on chronic kidney disease-epidemiology collaboration equation, based on estimated creatinine serum, gender equation coefficient, and age. Plasma creatinine levels were also measured with enzymatic colorimetric.

### Measurement ELMO-1

Blood samples as much as 5 ml were collected from the antecubital vein. The centrifuge was performed within 30 min after collection and samples stored in –20°C before performing ELISA. Sandwich ELISA protocol for human ELMO-1 was used to measure ELMO-1 (MyBiosource<sup>®</sup>) with catalog number MBS9321199. Random twice repetition of ELISA for some samples was used to confirm the results of ELISA.

## Results

### Baseline subject characteristics

Baseline subject characteristics are shown in Table 1. A total of 60 subjects who fulfilled the research inclusion criteria participated in this study, with 20 subjects in each group.

Demographic characteristics shown that the mean age was younger in the A1 group than the other groups. Female subjects were more frequent in the A1 and A2 groups, but it was not statistically significant.

We measured kidney function for each subject. The A1 group had lower urine creatinine, but it was not statistically significant. Lower plasma creatinine, urine albumin, ACR, and higher GFR were found in the A1 group compared to the other groups, and the differences were statistically significant. These baseline subject characteristics shown that the A1 (non-ND) group had better kidney function than the A2 and A3 groups (ND).

All patients had been diagnosed with T2DM. The A1 group had lower fasting and postprandial plasma glucose than the A2 and A3 groups, but it was not statistically significant. Lower systolic and diastolic blood pressures were also found in the A1 group compared to the other groups, and the result was statistically significant.

### Plasma ELMO-1 level was associated with the severity of kidney level damage based on ACR when DN occurs

The mean of ELMO-1 plasma level based on the grade of ACR is shown in Figure 1. ELMO-1 plasma

**Table 1: Baseline subject characteristic**

Characteristic	A1 (n = 20)	A2 (n = 20)	A3 (n = 20)	p-value
Age and years old	48.05 ± 9.09	50.35 ± 7.36	51.95 ± 7.64	0.314**
Gender				0.619
Male, n	8	7	10	
Female, n	12	13	10	
Fasting plasma glucose, mg/dL	140.95 ± 35.49	154.80 ± 53.98	164.95 ± 57.69	0.513*
Postprandial plasma glucose, mg/dL	242.7 ± 96.09	224.55 ± 50.66	257.10 ± 57.79	0.321*
Systolic blood pressure, mmHg	120.00 (100.00 – 160.00)	130.00 (110.00 – 150.00)	150.00 (110.00 – 170.00)	<0.001**
Diastolic blood pressure, mmHg	80.00 (50.00 – 90.00)	80.00 (60.00 – 90.00)	80.00 (70.00 – 100.00)	0.042**
Plasma creatinine, mg/dL	0.88 ± 0.38	0.90 ± 0.22	1.92 ± 1.06	0.001*
Urine creatinine, mg/dL	86.21 (13.43 – 377.97)	114.04 (19.69 – 351.55)	79.31 (11.43 – 192.99)	0.217**
Urine albumin, mg/dL	14.7 ± 10.77	89.25 ± 83.71	2015.80 ± 2755.93	<0.001*
Albumin creatinine ratio	12.82 (4.06 – 28.44)	47.36 (30.68 – 203.77)	2263.00 (349.96 – 14403.00)	<0.001**
Glomerular filtration rate, mL/min	92.66 ± 23.59	84.17 ± 22.55	50.34 ± 37.10	<0.001*

\*One-way ANOVA was performed due to normally distributed data. \*\*Non-parametric test was performed due to not normally distributed data even after transformation with log 10.

level for each group was not normally distributed even after transformation with log 10, so the non-parametric test was performed. The mean value of the ELMO-1 plasma level of the A1 group was lower than the mean value of ELMO-1 plasma level of A2, but it was not statistically significant ( $p = 0.779$ ). With the limited number of samples, this result has shown that ELMO-1 was not associated with the level of kidney damage of microalbuminuria (A1 group) to early macroalbuminuria (A2 group).

**Table 2: ELMO-1 level based on clinical characteristics affected by DN**

p parameter	ELMO-1 level
Glomerular filtration rate	
GFR ≥60 mL/min, n = 41	134.22 (29.10 – 3726.78)
GFR <60 mL/min, n = 19	128.54 (61.77 – 3446.93)
Fasting plasma glucose	
FPG ≤130 mg/dL, n = 25	121.44 (33.36 – 3446.93)
FPG >130 mg/dL, n = 35	134.22 (29.10 – 3726.78)
Postprandial plasma glucose	
PPG ≤180 mg/dL, n = 9	144.16 (94.45 – 3131.57)
PPG >180 mg/dL, n = 51	128.54 (29.10 – 3726.78)
Blood pressure	
SBP <140 mmHg and DBP <90 mmHg, n = 26	134.22 (29.10 – 1976.67)
SBP ≥140 mmHg and/or DBP ≥90 mmHg, n = 34	126.41 (61.77 – 3726.78)

The mean value of ELMO-1 plasma level of A1 group was lower than the mean value of ELMO-1 plasma level of A3, and it was statistically significant ( $p = 0.005$ ). The mean value of ELMO-1 plasma level of the A2 group was lower than mean value of ELMO-1 plasma level of the A3 group, and it was statistically significant ( $p = 0.009$ ). This result shown that ELMO-1 plasma level was associated with level of kidney damage based on ACR when DN has occurred (A2 and A3 groups were confirmed DN).

**Table 3: Correlation of ELMO-1 with clinical parameter**

Kidney function	Serum ELMO-1 (ng/mL)	p-value
R		
Plasma creatinine, mg/dL	0.078	0.554
Urine creatinine, mg/dL	-0.100	0.447
Urine albumin, mg/dL	0.344	0.007
Fasting plasma glucose, mg/dL	0.013	0.919*
Postprandial plasma glucose, mg/dL	0.110	0.403
Systolic blood pressure, mmHg	0.152	0.245
Diastolic blood pressure, mmHg	0.043	0.745
Glomerular filtration rate, mL/min	-0.293	0.023*
Albumin creatinine ratio	0.383	0.003
Age, years old	0.197	0.132*

\*Pearson correlation test was performed due to normally distributed data of one variable. Otherwise, the Spearman correlation was performed.

Table 2 shows that the mean value of ELMO-1 plasma level was higher in variables affected by DN. Higher plasma ELMO-1 levels were found in the lower

GFR group, in higher plasma glucose fasting and 2 h postprandial plasma glucose and higher blood pressure groups. Although plasma ELMO-1 level difference were not significant statistically between these groups.

### Plasma ELMO-1 level was correlated with kidney function in all subjects with T2DM

That correlations of ELMO-1 with some clinical parameters which play a role in the progression of DN are shown in Table 3. Urine albumin and ACR have a positive weak correlation with ELMO-1 plasma level, and the result was statistically significant. GFR had a negative weak correlation with ELMO-1 plasma level, and the result was statistically significant.

Plasma creatinine, plasma glucose, blood pressure, and age had very weak positive correlations with ELMO-1 plasma level. Urine creatinine had a very weak negative correlation with ELMO-1 plasma level. This correlation was not statistically significant.


## Discussion

DN, also known as diabetic kidney disease, causes structural changes and declining kidney function. Albuminuria and GFR are clinical markers widely used to monitor the progression of DN. Both markers are endpoints of many pathways, so prevention of decreases of the two markers is not easy. Identifying additional new markers which are directly involved in certain dominant pathways can assist in gaining better understanding of DN pathophysiology and help develop new approaches in DN treatment [3], [4], [5].

Analyses in our research shown the severity of albuminuria were associated with plasma ELMO-1 when DN was evident (Figure 1). Furthermore, in all study, plasma ELMO-1 was correlated with declining of kidney function in patients with T2DM based on levels of proteinuria, ACR, and GFR (Table 2). This finding demonstrates that the ELMO-1 plasma level is



a promising biomarker of the severity level of kidney damage in DN in the Indonesia population. In addition, ELMO-1 plasma level is also a promising marker of declining kidney function in patients with T2DM.



**Figure 1:** Mean of engulfment and cell motility protein-1 plasma level based on the grade of albumin to creatinine ratio

Studies of ELMO-1 plasma or tissue specifically associated with diabetes or DN in humans are still limited. One study found kidney expression of ELMO-1 in the non-diabetic compared to diabetic patient remained unchanged in podocytes and tubules, but it did not define whether the diabetic patient was suffering DN or not [14]. Phenotype genotype association studies have shown that the ELMO-1 gene is a strong gene candidate associated with DN. Single nucleotide polymorphism in the intronic region of ELMO-1 gene was associated with DN in Asia and American populations, but none described the correlation with systemic or local ELMO-1 expression [9], [10], [11], [12].

One supporting study about the role of ELMO-1 related to DN described the association using an animal model. Using a zebrafish animal model, the results shown that ELMO-1 knockout model has a more detrimental effect on pronephric structure and function under hyperglycemia conditions compared to controls. This finding related to the detrimental effect on the processes of podocytes and glomerular ultrafiltration. Injection of ELMO-1 mRNA to create ELMO-1 overexpression and pan-caspase inhibitor rescue altered the renal structure and function. This research shown that ELMO-1 was an important factor for glomerular protection and renal cell survival that decreasing apoptosis, especially under diabetic conditions [14]. Another study found that a rat model with a mutation of in insulin-2 gene had different levels of ELMO-1. The research reported that higher ELMO-1 expression was associated with the severity of renal fibrosis and amount of urinary albumin excretion. The increased expression of ELMO-1 was followed by the aggravated marker of oxidative stress and NAD(P)H oxidase 4 and reduced glutathione, and the aggravated markers of fibrogenesis, TGF $\beta$ -1, and endothelin-1 [13].

Baseline subject characteristics shown that blood mean systolic and diastolic blood pressures were associated with the degree of albuminuria, but

plasma glucose was not, although lower plasma glucose was found in the A1 group compared to the A2 and A3 groups. The previous study shown controlling glucose plasma and blood pressure influences the progression of DN, although the duration of diabetes also contributes to DN [2], [6], [7]. Age and gender were not confounding factors for this research, and in the baseline characteristics, it did not differ statistically between the three groups.

Random spot urine sampling for ACR is used as one of the markers of DN progression. In this research, we used this marker as a reference to know DN progression because a previous study reported that it was representative and more practical to assess proteinuria related to diabetic complications than 24-h urine samples. In the next step, we grouped ACR into three groups. There were A1 or mildly albuminuria, A2 or moderately increased or microalbuminuria, and A3 or severe increased or macro-albuminuria. DN was confirmed at 6 diagnosed with ACR in groups A2 and A3 [7], [15]. To the best of our knowledge, this is the first study of the association of circulating ELMO-1 with DN in the Indonesia population. Some limitations of our study were that it was conducted in only one center and had a limited number of participants which reduced the generalizability of the findings. To confirm the finding of ELMO-1 as a progression marker of severity in DN, further multicenter studies with larger numbers of participants are needed.

## Conclusions

Plasma ELMO-1 level is associated with the severity level of kidney damage based on ACR when DN has occurred. Albuminuria and ACR had positive correlations with plasma ELMO-1 and GFR had a negative correlation with plasma ELMO-1 in all subjects who suffered T2DM. These findings shown that plasma ELMO-1 is a promising marker of the level of kidney damage, especially when DN has occurred in the Indonesia population. According to the limitations of this study, larger numbers of subjects and multicenter research are needed.

## Acknowledgment

This study was a part of the author's Master thesis. The authors would like to acknowledge the contribution of staff at the biotechnology laboratory of the Medical Faculty University of Andalas, Prodia laboratory and Genetica Science laboratory, for their help in laboratory measurements.

## References

1. Koye DN, Magliano DJ, Nelson RG, Pavkov ME. The global epidemiology of diabetes and kidney disease. *Adv Chronic Kidney Dis.* 2018;25(2):121-32. <https://doi.org/10.1053/j.ackd.2017.10.011>  
PMid:29580576
2. Gheith O, Farouk N, Nampoory N, Halim MA, Al-Otaibi T. Diabetic kidney diseases. World wide difference of prevalence and risk factors. *J Nephropharmacol.* 2016;5(1):49-56.  
PMid:28197499
3. Persson F, Rossing P. Diagnosis of diabetic kidney disease: State of the art and future perspective. *Kidney Int Suppl.* 2018;8(1):2-7.  
PMid:30675433
4. Lee C, Lam KS. Biomarkers of progression in diabetic nephropathy: The past, present and future. *J Diabetes Investig.* 2015;6(3):247-9. <https://doi.org/10.1111/jdi.12329>  
PMid:25969706
5. Currie G, Mckay G, Delles C. Biomarkers in diabetic nephropathy: Present and future. *World J Diabetes.* 2014;5(6):763-76.  
PMid:25512779
6. Alicic RZ, Rooney MT, Tuttle KR. Diabetic kidney disease: Challenges, progress and possibilities. *Clin J Am Soc Nephrol.* 2017;12(12):2032-45. <https://doi.org/10.2215/cjn.11491116>  
PMid:28522654
7. Fioretto P, Mauer M. Histopathology of diabetic nephropathy. *Semin Nephrol.* 2007;27(2):195-207. <https://doi.org/10.1016/j.semnephrol.2007.01.012>  
PMid:17418688
8. Battle D. Clinical and cellular markers of diabetic nephropathy. *Kidney Int.* 2003;63(6):2319-30. <https://doi.org/10.1046/j.1523-1755.2003.00053.x>  
PMid:12753325
9. Shimazaki A, Kawamura Y, Kanazawa A, Sekine A, Saito S, Tsunoda T, *et al.* Genetic variations in the gene encoding ELMO1 are associated with susceptibility to diabetic nephropathy. *Diabetes.* 2005;54(4):1171-8. <https://doi.org/10.2337/diabetes.54.4.1171>  
PMid:15793258
10. Pezzolesi MG, Katavetin P, Kure M, Poznik GD, Skupien J, Mychaleckyj JC, *et al.* Confirmation of genetic associations at ELMO1 in the GoKinD collection supports its role as a susceptibility gene in diabetic nephropathy. *Diabetes.* 2009;58(11):2698-702. <https://doi.org/10.2337/db09-0641>  
PMid:19651817
11. Bodhini D, Chidambaram M, Liju S, Revathi B, Laasya D, Sathish N, *et al.* Association of rs11643718 SLC12A3 and rs741301 ELMO1 variants with diabetic nephropathy in South Indian population. *Ann Hum Gen.* 2016;80(6):1-6. <https://doi.org/10.1111/ahg.12174>  
PMid:27699784
12. Mehrabzadeh M, Pasalar P, Karimi M, Abdollahi M, Daneshpour M, Asadolahpour E, *et al.* Association between ELMO1 gene polymorphisms and diabetic nephropathy in an Iranian population. *J Diabetes Metab Disord.* 2016;15:43. <https://doi.org/10.1186/s40200-016-0265-3>  
PMid:27761430
13. Hathaway CK, Chang AS, Grant R, Kim H, Madden VJ, Bagnell CR, *et al.* High Elmo1 expression aggravates and low Elmo1 expression prevents diabetic nephropathy. *Proc Natl Acad Sci U S A.* 2016;113(8):2218-22. <https://doi.org/10.1073/pnas.1600511113>  
PMid:26858454
14. Shama KR, Heckler K, Stoll SJ, Hillebrands J, Kynast K, Herpel E, *et al.* ELMO1 protects renal structure and ultrafiltration in kidney development and under diabetic conditions. *Sci Rep.* 2016;6:37172. <https://doi.org/10.1038/srep37172>
15. Roshan B, Stanton RC. A story of microalbuminuria and diabetic nephropathy. *J Nephropathol.* 2013;2(4):234-40.  
PMid:24475455

# Plasma Levels of the Engulfment and Cell Motility Protein-1 are Associated with Kidney Damage in Diabetic Nephropathy: A Single-Center Pilot Study in Indonesia Population

## ORIGINALITY REPORT

9%

SIMILARITY INDEX

6%

INTERNET SOURCES

8%

PUBLICATIONS

1%

STUDENT PAPERS

## PRIMARY SOURCES

1

[eprint.ncl.ac.uk](http://eprint.ncl.ac.uk)

Internet Source

<1%

2

[www.researchsquare.com](http://www.researchsquare.com)

Internet Source

<1%

3

Katherine R Tuttle, Mark C Lakshmanan, Brian Rayner, Robert S Busch, Alan G Zimmermann, D Bradley Woodward, Fady T Botros.

"Dulaglutide versus insulin glargine in patients with type 2 diabetes and moderate-to-severe chronic kidney disease (AWARD-7): a multicentre, open-label, randomised trial", The Lancet Diabetes & Endocrinology, 2018

Publication

<1%

4

Chih-Hung Lin, Yi-Cheng Chang, Lee-Ming Chuang. "Early detection of diabetic kidney disease: Present limitations and future perspectives", World Journal of Diabetes, 2016

Publication

<1%

5	Yao Chen, Shu-ying Fang. "Potential genetic polymorphisms predicting polycystic ovary syndrome", Endocrine Connections, 2018 Publication	<1 %
6	ijms.sums.ac.ir Internet Source	<1 %
7	mafiadoc.com Internet Source	<1 %
8	www.dovepress.com Internet Source	<1 %
9	Anchalee Tantiwettrueangdet, Ravat Panvichian, Pattana Sornmayura, Surasak Leelaudomlipi. "PCNA-associated Factor (KIAA0101/p15PAF) over expression and Gene Copy Number Alterations in Hepatocellular Carcinoma Tissues", Research Square, 2020 Publication	<1 %
10	M.R. Manaviat, M. Rashidi, M. Afkhami-Ardekani. "Four Years Incidence of Diabetic Retinopathy and Effective Factors on its Progression in Type II Diabetes", European Journal of Ophthalmology, 2018 Publication	<1 %
11	Qijin Zhu, Xiangming Qi, Yonggui Wu, Kun Wang. "Clinical study of total glucosides of paeony for the treatment of diabetic kidney	<1 %



disease in patients with diabetes mellitus",  
International Urology and Nephrology, 2016

Publication

12

[www.j3.jstage.jst.go.jp](http://www.j3.jstage.jst.go.jp)

Internet Source

<1 %

13

[ir.uiowa.edu](http://ir.uiowa.edu)

Internet Source

<1 %

14

[www.unboundmedicine.com](http://www.unboundmedicine.com)

Internet Source

<1 %

15

Irena Kostovska, Katerina Tosheska Trajkovska, Svetlana Cekovska, Sonja Topuzovska et al. "Role of urinary podocalyxin in early diagnosis of diabetic nephropathy", Romanian Journal of Internal Medicine, 2020

Publication

<1 %

16

[cdltu.edu.np](http://cdltu.edu.np)

Internet Source

<1 %

17

Sangyeoup Lee, Yun Jin Kim, Tae Yong Jeon, Hyoung Hoi Kim, Sang Woo Oh, Yongsoon Park, Sung Soo Kim. "Obesity is the only independent factor associated with ultrasound-diagnosed non-alcoholic fatty liver disease: A cross-sectional case-control study", Scandinavian Journal of Gastroenterology, 2009

Publication

<1 %

18

Xue, Hao, Zhuang Lu, Wen Lu Tang, Lu Wei

Pang, Gan Mi Wang, Gavin WK Wong, James M Wright, and Wen Lu Tang. "First-line drugs inhibiting the renin angiotensin system versus other first-line antihypertensive drug classes for hypertension", Cochrane Database of Systematic Reviews, 2015.

Publication

<1 %

19

[www.hindawi.com](http://www.hindawi.com)

Internet Source

<1 %

20

[www.rssdi.org](http://www.rssdi.org)

Internet Source

<1 %

21

Aya Mikdache, Laura Fontenas, Shahad Albadri, Celine Revenu et al. "Elmo1 function, linked to Rac1 activity, regulates peripheral neuronal numbers and myelination in zebrafish", Cellular and Molecular Life Sciences, 2019

Publication

<1 %

22

Godfrey Katamba, David Collins Agaba, Francis Muzaale, Agnes Namaganda et al. "Neck circumference as marker of overweight and obesity among secondary adolescents aged 12-19 years in southwestern Uganda: a cross-sectional survey", Research Square, 2021

Publication

<1 %

23

Mariana de Ponte, Vanessa Cardoso, Juliana Costa-Pessoa, Maria Oliveira-Souza. "Early Diabetes Aggravates Renal

<1 %

# Ischemia/reperfusion-induced Acute Kidney Injury ", Research Square, 2021

Publication

24

R. Walz, S.T. Terracciano, M. Bianchin, M.L.F. Chaves, M.R. Cerski, L.C. Severo, A.T. Londero. "2-04-16 Cerebral phaeohyphomycosis due to cladophialophora bantiana. Report of Brazilian case and comments on the pertinent literature", Journal of the Neurological Sciences, 1997

Publication

<1 %

25

The Genetics of Type 2 Diabetes and Related Traits, 2016.

Publication

<1 %

26

[makperiodicallibrary.com](http://makperiodicallibrary.com)

Internet Source

<1 %

27

[test.dovepress.com](http://test.dovepress.com)

Internet Source

<1 %

28

[www.ejode.eg.net](http://www.ejode.eg.net)

Internet Source

<1 %

29

Jun Ishizaki, Ayako Takemori, Kenta Horie, Daisuke Hiraoka et al. "Usefulness of tissue inhibitor of metalloproteinase 1 as a predictor of sustained remission in patients with antineutrophil cytoplasmic antibody-associated vasculitis", Research Square, 2020

Publication

<1 %

---

30

[www.acc.org](http://www.acc.org)

Internet Source

<1%

---

31

Junlin Zhang, Yiting Wang, Rui Zhang, Hanyu Li, Qianqian Han, Ruikun Guo, Tingli Wang, Li Li, Fang Liu. "Implication of decreased serum complement 3 in patients with diabetic nephropathy", *Acta Diabetologica*, 2017

Publication

<1%

---

32

Yujing Wang, Yuqin Gao, Yang Xun. "Work Engagement and Associated Factors Among Dental Nurses in China", *Research Square*, 2021

Publication

<1%

---

---

Exclude quotes

Off

Exclude matches

Off

Exclude bibliography

On