

Red Wood (*Pterocarpus Indicus* Wild) and Bread Fruit (*Artocarpus Communis*) Bark Sap as Attractant of Stingless Bee (*Trigona* Spp)

By Salni Salni



8
Red Wood (*Pterocarpus Indicus* Wild) and Bread Fruit (*Artocarpus Communis*) Bark Sap as Attractant of Stingless Bee (*Trigona* Spp)

Hanifa Marisa and Saini

11
13
Biology Department of Faculty of Science, The University of Sriwijaya, Km 32 Indralaya, South Sumatera Province, Indonesia 30662

Received 15 November 2011, Revised 29 December 2011, Accepted 1 February 2012, Available online 26 February 2012

ABSTRACT

10
An investigation about the power of redwood bark sap to invite stingless bee arrival had been done at Biology campus area, The University of Sriwijaya, Indralaya, South Sumatera, Indonesia; during Nopember 2010. Redwood breadfruit bark have been hurt for outcoming the red sap, and duration time for stingless bee arrival be measured then. Meanly, in 17 minutes, stingless bee had came for red sap; and in 5 minutes for breadfruit. Chemical test of redwood sap after soluted in methanol and put on gelsilicon chromatography with ethil-acetate and methanol (1:1), and sprayed by sulphic acid indicate that the compounds is terpenid (tannin) for red wood sap, and tannin and alcaloid for bread fruit sap.

| *Pterocarpus indicus* | *Artocarpus communis* | *Trigona spp* | bark sap | gum |

4
© 2012 Ibnu Sina Institute. All rights reserved.
<http://dx.doi.org/10.11113/mjfas.v8n2.132>

1. INTRODUCTION

At an afternoon, one year ago, a neighbor cut the red wood trunks in front of the house. Many red elastic sap/gum came out from the bark and lot of insects, stingless bee, hold the gum and keep it in their leg baskets. These phenomena were interested. What were the chemical compounds in the red sap? And what for stingless bee to keep it? The study on *Trigona* and *Pterocarpus* begin at that day. Even, *Trigona sp.* also be found at bread fruit sap too 12 any months later.

At least there are 100.000 chemical compounds produced by 200.000 species of flowering plants in their growth and development. Most of them were not essential for physiological process of that plant.

The structure of more than 6000 alkaloids, 3000 terpenoids, some thousands of phenylpropanoid, 1000 flavonoid, 500 qu inons, 650 pol yacetylene and 400 of amino acidshad been known. Most of chemical compounds of plant identified, were functioned as volatile attractant for insects. Attraction of those compounds processed by diffusion in the air (Metcalf, 1992).

In Philippines, stingless bees are mass produced for pollination and honey production. During the swarming season (February-March) strong colonies are divided. Broad chambers containing queen cells are removed along with same adult bees, and placed 1-2 metres away from the original colony. These so called 'starter' colonies are then allowed to rear their own queen (Fajardo and Cervancia, 2011).

2
Heard (1999) find that stingless bee are common visitors to flowering plants in tropics, but evidence for

their importance and effectiveness as crop pollinators is lacking for most plant species. They are known to visit the flowers of 90 crop species. They were confirmed to be effective and important pollinator of 9 species. They make a contribution to the pollination of 60 other species, but there is insufficient information to determine their overall effectiveness or importance. They have been recorded from another 20 crops, but other evidence suggests that they do not have an important role because these plants are pollinated by other means.

The nest of stingless bees usually consist of an external tube, internal tunnel, resin dumps, wax dumps, foot pots for storing pollen and honey, brood and nest envelopes like involucre and batumen. Brood cells, honey pots and pollen pots are arranged in separate clusters. Brood cells and food pots made of cerumen which is a mixture of wax and resin. (Pooley and Michener, 1969 in Danaraddi, 2007). Eltz *et al.* (2002) in Danaraddi (2006) reported the nest density of *Trigona collina* Smith, was generally high in sites located in the Sipilok forest fragment (mean 8.4 nest/ha). Nest densities in continuous forest were all low (between 0 and 2.1 nest/ha, mean 0.5 nest / ha).

7
Boogert *et al.* (2006) found that spesies *Trigona corvina*, scent marked a high quality food source and that they used their own scent marks to relocate it in choice experiment. Boogert studied for some species; were bumblebees, honey bees and stingless bees.

Stingless bees need flower nectar and pollen for their food. Stingless bees could find the food until 500 metres (Baconawa 1999 cit. Abdillah 2008). Nectar of rubber plant and Cassava even were got by *Trigona* from their stem (Abdillah, 2008). Compared with Apis, population of *Trigona* colony were larger (Anonymous 2001 cit. Abdillah 2008).

Corresponding author at:
E-mail addresses: gndiqhan2002@yahoo.com (Hanifa Marisa)

MacArthur and Pianka argued that to obtain food, any predator (herbivore) must expend time and energy, first in searching for its object, and then in handling it (i.e. Pursuing, subduing and consuming it). Searching is bound to be directed, to some extent, toward particular food/prey type, but while searching, a predator/herbivore is nevertheless likely to encounter a wide variety of food items. (Begon, M., Harper, J.L. and Colin R Towsend, 1986).

Table 1 Time that were needed by *Trigona sp1* to find and arrived at bark hurt of *P indicus*.

Number of replication	REDWOOD	BREADFRUIT
1	19	3
2	15	4
3	18	5
4	16	4
5	17	9

2. EXPERIMENTAL

Redwood and breadfruit bark have been hurt for outcoming the red sap, and duration time for stingless bee arrival be measured then by watch (searching time). Five hurt done, and mean of data were measured. . Chemical test of redwood sap after soluted in methanol and put on gelsilicon chromatography with ethil-acetate and methanol (1:1) for redwood sap; and *n*-hexan for breadfruit, and sprayed by sulphic acid 10 % after dried. The colour of spot were watched then (Thin Layer Chromatography) as published by Fransworth (1966)

3. RESULT AND DISCUSSION

As Begon *et al.* (1986) explained, it is needed a certain time for insect to find the foot object (searching time). *Trigona sp1* (smaller) need about 17 minutes to find the red wood gum. Furthermore, *Trigona sp2* (bigger) need about 5 minutes to find the object. Because the sap of those trees were not nectar and pollen organ; it must be used of *Trigona spp* for nest built (see Baconawa 1999 cit. Abdillah 208). But, Roubik and Patino (2009) reported, that *Trigona* nest were built from pollen too. Seventy two plants species pollen were found in *Trigona corvine* nest scutellum. Lemberg *et al.* (2008) reported that *Trigona* used resin-like substances as glue against intruders to their nest.

Body size, as could saw at **Figure 4**, inferences of flight range and possible ecological implications included the finding of food of course. Data suggested that maximum flight distance in Meliponini, -stingless bee group- is a function of body size, especially generalized wing size, which can be estimated through principal

component analysis, as had been done by Araujo *et al.* (2004).

Worker bee size reflects an adaptation to environmental conditions; a major part of the morphological variation in Meliponini occurs independently of phylogeny due to the fact that, for social bees, worker body size has been generally considered as an adaptation to foraging activity and floral resource exploitation (Pignata and Dimiz-Filho, 1996 cit. Arraujo *et al.* 2004).

Table 2 Result of chemical compound test by sulphic acid reagent

NO	SAMPLE (SAP)	COLOUR AFTER REAGENT TEST
1	RED WOOD	BROWN (TERPENOID)
2	BREAD FRUIT	VIOLET-BROWN (TANNIN) AND YELLOW (ALCALOID)

Furthermore, it is known that stingless bee is social insect that have specific framework in activity of foraging. The foraging is a complex process involving large numbers of individuals collecting food from many different sources. The state of foraging behavioral were included 5 points; waiting at the nest and available to start foraging, searching the food source, exploiting food source, recruiting nest mates to food source and follow recruiters to food source. By those mechanisms, time of finding food source be influenced (Sumpter and Pratt, 2003).



Fig. 1 (from above to below): Red wood bark hurt and red gum/sap came out

Syah (2005) has isolated the chemical compounds of *Artocarpus communis* and found 50 phenolic compounds those were calcon, flavanon, 3-phenylflavon and stilbene. Most of those chemical compounds were characterized as cytotoxic, anti-inflammation, anti-malariae, anti-microbe and inhibitor for some enzymes. Specific for tree bark, the chemical compounds that were isolated are: 3-phenylflavon, Dihydrobenzoxanton, Xuinonodihydrobenzoxanton, furanodihydrobenzoxanton and tetrahydroxanton. Other compounds were found at flower, root and root epiderm. *Trigona sp* nest flavonoid (propolis) reported had the antibacterial activity against *Streptococcus mutans* (in vitro) (Sabir, 2005). Luttrell (2010) wrote, plants resin added for honey and pollen pots where the antifungal, antibacterial and other biological activity of the plants resins aid in the preservation of honey and pollen.



Fig. 2 Thin Layer Chromatography test of Redwood gum by H_2SO_4 reagent



Fig. 3 Spot of gum drop of red wood



Fig. 4 Two species of *Trigona* spp, the bigger (left) attracted to bread fruit sap and the smaller (right) attracted to red wood gum



Fig. 5 *Trigona* crowded at Red wood gum. Many *Trigona* sp are taking the gum

It could be seen, that the species of *Trigona sp* show difference in liking the sap of tree bark. The smaller, came to red wood gum and the bigger came to bread fruit sap. In niche concept of organism, every species has own specific status of life, where tend to different to the other. If two or more species have the same niche, they should compete each other, and one of them should change their life niche.

At other hand, some facts may influence the time to find out the trees sap. Distance to the nest, air flow condition and capability of catch the smell of gum could do the main role in this situation. Even, carbon dioxide, as a content of air, may influence the attraction of insect to object. In bed bug (*Cimex lectularius L*), for example, carbon dioxide, heat and chemical lure, influence the attraction to pitfall trap (Wang *et al.*, 2009). In bats species, ultrasound emissions of wind turbines show the potential attractant for them (Szewczak, 2006).

ACKNOWLEDGEMENT

We thank the rector of The University of Sriwijaya for facilitation and funding; and to RAFSS 2011 UTM for giving the presentation chance.



Fig. 6 *Trigona sp* (big size body), attacking the tip of arm that closed to their nest, a hundred meter from bread fruit gum. The nest made at an *Acacia sp* tree (left above corner at picture)

4. CONCLUSION

Red wood and bread fruit bark sap attract *Trigona spp* for nest material. It took time about 17 minutes to find red wood gum by *Trigona sp1* and 5 minutes to find breadfruit sap by *Trigona sp2*. Differentiation of time to finding the gum may be because of the different species niche, included body size and air condition.

REFERENCES

- [1] Abdillah, Habib. Effect of volume of stup to weight colony and activity bee of klanceng (*Trigona sp*). Faculty of Animal Husbandry The University of Brawijaya. Malang. 2008
- [2] Araujo, E D., Costa M, Chaud-Netto J, and Fowler H G., Braz. J. Biol. 64 (3 B); (2004)563-568.
- [3] Begon, Michael, Harper John L, and Colin R Townsend. 1986. Ecology. Blackwell Scientific Publications. Oxford
- [4] Boogert, Neeltje Janna., Frouke Elisabeth Hofstede., and Ingrid Aguilar Monge., Apidologie 37 (2006) 366-375
- [5] Danaraddi, C. S. Studies on stingless bee, *Trigona iridipennis* Smith with special reference to foraging behaviour and melissopalynology at Dhaward, Kamataka. Master of Science Thesis. College of Agricultural Science, Dharwad. 2007
- [6] Fajardo A C and Cleo RC Cervancia., Bees for Development Journal. (67) 2011; p 3-5
- [7] Fransworth, NS., J Pharmaceutical Science Vol 55 (3). 1966.
- [8] Heard, Tim A., Rev. Entomol. 1999 (44); 183-206
- [9] Lemberg, Lars., Dworscak K, and Nico Blutgen., J Agr Research and Bee World. 47 (1); (2008)p17-21
- [10] Luttrell, Bob. The Role of Plant Resins and the Cadaghi Enigma. Bob the Beeman. 2010 Page 1-4.
- [11] Metcalf, Robert Lee. Plant Kairomons In Ecology and Insect Control. Chapman and Hall Inc. London. 1992.
- [12] Roubik, David. W and J Enrique Moreno Patino. *Trigona corvine*: An Ecological Study Based on Unusual Nest Structure and Pollen Analysis. Hindawi Publishing Corporation Psyche. Vol 2009. ID 268756.
- [13] Sabir, Ardo., Dent. J. Vol 38 No 3, (2005) P 135-141.7.
- [14] Sumpter, D.J.T and S. C. Pratt., Behav. Ecol. Sociobiol. 53; (2003)131- 144.
- [15] Syah, Yana Maolana., Bulletin of The Indonesia Society of natural products Chemistry. Vol 5 No 2 (2005)pp 33-50.
- [16] Szewczak, Joseph M. and Ed Arnett. Ultrasound emissions from wind turbines as a potential attractant to bats; a preliminary investigation. Foote Creek Rim Wind Facility. Arlington, WY. 2006.
- [17] Wang, Changlu., Timothy Gibb, Gary W Bennett and Susan Mc Knight., J. Econ. Entomol. 102 (4); 2009. pp 1580-1585.

Red Wood (*Pterocarpus Indicus* Wild) and Bread Fruit (*Artocarpus Communis*) Bark Sap as Attractant of Stingless Bee (*Trigona* Spp)

ORIGINALITY REPORT

17%

SIMILARITY INDEX

PRIMARY SOURCES

1	myjurnal.my Internet	40 words — 2%
2	link.springer.com Internet	34 words — 2%
3	acervodigital.unesp.br Internet	32 words — 2%
4	repo.uum.edu.my Internet	30 words — 2%
5	www.authorstream.com Internet	27 words — 1%
6	Thomas Eitz, Carsten A. Brühl, Sander van der Kaars, Eduard K. Linsenmair. "Determinants of stingless bee nest density in lowland dipterocarp forests of Sabah, Malaysia", <i>Oecologia</i> , 2002 Crossref	26 words — 1%
7	mafiadoc.com Internet	22 words — 1%
8	sinta3.ristekdikti.go.id Internet	22 words — 1%
9	www.control.utoronto.ca Internet	16 words — 1%

10	pinpdf.com Internet	16 words — 1%
11	repository.unib.ac.id Internet	14 words — 1%
12	Turna, Michael T. "Extraction of a phagostimulant and classification of the feeding recognition template for larvae of the moth Malacosoma americanum", Proquest, 20111108 ProQuest	12 words — 1%
13	hal.archives-ouvertes.fr Internet	11 words — 1%
14	D. Sumpter, S. Pratt. "A modelling framework for understanding social insect foraging", Behavioral Ecology and Sociobiology, 2003 Crossref	9 words — < 1%
15	Christoph Grüter. "Stingless Bees", Springer Science and Business Media LLC, 2020 Crossref	7 words — < 1%

EXCLUDE QUOTES OFF
EXCLUDE BIBLIOGRAPHY ON

EXCLUDE MATCHES OFF