

The Effectivity of Jambu Akasia (Bellucia Pentamera Naudin) fruit extract for killing the Aedes aegypti L. Larvae

by Salni Salni

Submission date: 03-May-2020 09:39AM (UTC+0700)

Submission ID: 1314341548

File name: The_effectivity_of_jambu.pdf (736.61K)

Word count: 1245


Character count: 6500

PAPER • OPEN ACCESS

The Effectivity of Jambu Akasia (*Bellucia Pentamera* Naudin) fruit extract for killing the *Aedes aegypti* L. Larvae

1
To cite this article: H Marisa *et al* 2018 *J. Phys.: Conf. Ser.* **1116** 052041

View the [article online](#) for updates and enhancements.



IOP ebooks™

Bringing together innovative digital publishing with leading authors from the global scientific community.

Start exploring the collection—download the first chapter of every title for free.

The Effectivity of Jambu Akasia (*Bellucia Pentamera* Naudin) fruit extract for killing the *Aedes aegypti* L. Larvae

H Marisa*, Salni and F Salfamas

Department of Biology, Faculty of Mathematics and Natural Sciences, Universitas Sriwijaya, KM 32 Indralaya, South Sumatra, Indonesia 30662

*E-mail: gmdiqhan2002@yahoo.com

Abstract. A research about effectivity of *B pentamera* fruits extract on mortality of *Ae.aegypti* larvae had been conducted during November 2017–January 2018 at P2B2 Research Center Baturaja and Biology Laboratory Faculty of Mathematics and Natural Sciences, Universitas Sriwijaya. The aims of this study were to gain the effectivity and LC₅₀ value of extract on larvae mortality. Completely Randomized Design was used as experimental design. Seven treatments were tested (Abate 0.0001 %; Negative control; 1%; 1.5%; 2%; 2.5% and 3% methanolic extract). Extraction of fruit material with methanol as maceration process followed with vacuum rotary evaporator concentration. One-way ANOVA was used as data analysis and LC₅₀ obtained from Probit Analysis. The study found that concentration of methanolic fruit extract at 2.5% killed 52% larvae during 24 hr experiment and 63% during 48 hr. LC₅₀ values of fruits extract were 3.34% for 24 hours experiment and 3.32% for 48 hours. Thin Layer Chromatography methods was used for bioautography test of fruit extract, and three spots were found and identified as terpenoid, tannin and flavonoid compounds, based on colour appearance. In conclusion, fruit extract of *B. pentamera* is considered effective in killing *Ae aegypti* larvae yet a potential larvicide for future study.

1. Introduction

It has been known that *B pentamera* leaves extract could kill 17.5% larvae of *Ae.aegypti* during 24 hours and 37.5% during 48 hr of exposure [1]. The study was based on the fact that leaves extract could inhibit the growth of some bacterial strains [2]. Furthermore, fruits of *B.pentamera* have long been used as anti-helminthic therapy [3], yet promoting the possibility of further potential to be tested as potential larvicides, especially against *Ae.aegypti* larvae. To prove this assumption, while based on previous report, research on finding the effectivity of methanolic fruit extract on *Ae. aegypti* larvae has been conducted at Laboratory of Biology, Faculty of Mathematics and Natural Sciences, Universitas Sriwijaya and P2B2 Research Center, Baturaja, South Sumatra.

2. Material and Methods

Fruits of *B. pentamera* were sampled from Tanjung Batu villages, Ogan Ilir, South Sumatra. Completely Randomized Design was used in data analysis and as experimental design. Seven treatments were tested (Abate 0.0001 %; negative control; 1%; 1.5%; 2%; 2.5% and 3% fruit extract concentration with methanol as solvent). Extraction with methanol as maceration process solvent was further concentrated by using vacuum rotary-evaporator equipment. One-way ANOVA was used as data analysis and LC₅₀ value was obtained from Probit analysis.



Content from this work may be used under the terms of the Creative Commons Attribution 3.0 licence. Any further distribution of this work must maintain attribution to the author(s) and the title of the work, journal citation and DOI.

Published under licence by IOP Publishing Ltd

3. Results and Discussion

Mortality effect of fruits extract of *B. pentamera* on *Ae.aegypti* is shown in table below. The results showed that the increase of concentration of fruits extract from 0.0 to 3.0 influences the mortality significantly, but addition of the highest concentration until 3.0% did not show any significant mortality. The phenomena of hormesis was observed from this experiment. Previous authors reported the same case in *Pistia stratiotes* affected by petroleum liquid waste which made the number of *Pistia* profile of toxicity responded as hormesis too [4].

Table 1. Effect of *B pentamera* fruits extract on *Ae.aegypti* larvae mortality during 24 and 48

Concentration (%)	24 hours		48 hours	
	Mortality percentage	Mean values	Mortality percentage	Mean values
0.0	0	0.00 a	0	0.00 a
1.0	0	0.00 a	1	0.25 a
1.5	2	0.50 a	2	0.50 a
2.0	19	4.75 a	26	6.50 b
2.5	52	5.50 b	63	15.75 c
3.0	22	13.00 a	26	6.50 b
Abate 0.0001	96	24.00 c	99	24.75 d

Note; mean values notation at $\alpha = 5\%$

Tabel 2 about LC_{50} below, show that the value of fruits extract at 24 hours treatment was 3.34 and 48 hours treatment was 3.02%. The results explained that LC_{50} of fruits extract is obtained at lower potency than Abate 0.0001% as positive control. Previous study reported that fruits of *B. pentamera* contained vitamin C and could kill some viruses, but the ability of those compounds is not yet enough to kill *Ae. aegypti* in small concentration [5].

Table 2. LC_{50} *B pentamera* fruits extract on larvae mortality of *Ae. aegypti*

Duration of experiment	LC_{50}
24 hours	3.34
48 hours	3.02

The picture below shows the qualitative result of thin layer chromatography test of methanolic extract. Based on the color appearance, several compounds were detected such as flavonoid, tannin and terpenoid [6].



Figure 1 Color appearance in Thin Layer Chromatography;

1. Terpenoid (blue) 2. Tannin (brown) 3. Flavonoid (yellow)

4. Conclusion

Methanolic extract from fruits of *B. pentamera* possessed potential larvicidal activity by killing 52% at 2.5 concentration during 24 hr and 63% during 48 hr. Probit analysis of found that LC_{50} value of 3.34 for 24 hr and 3.02 for 48 hr experiment. Hormesis phenomenon was observed by an increase of mortality along addition of concentration and decrease later at 2.5 concentration. It is proposed to analyze LC_{50} by quadratic regression as well.

References

- [1] Marisa H and Salni 2017 *Bellucia pentamera* Naudin potency as natural medicine: Change the status from dangerous invasive to useful plant In *Paper of International Symposium on Natural Medicine* 23–24 August 2017 (Bogor: Indonesia)
- [2] Sari Y 2017 *Uji aktivitas antibakteri fraksi dan senyawa aktif ekstrak daun Kardia (Bellucia pentamera Naudin) terhadap Escherichia coli dan Staphylococcus aureus* (Palembang: Universitas Sriwijaya)
- [3] Slik F 2016 *Ann. Sci. Nat. ser.* **6** 105
- [4] Marisa H and Estuningsih SP 2017 *BIOVALENTIA: Biological Research Journal* **3** 45
- [5] Marisa H, Salni, F Salfamas and Y Oktariansyah 2017 Studi terhadap *Bellucia pentamera* Naudin: Perubahan status invasif menjadi bermanfaat larvasida In *Prosiding Seminar Nasional 2017 Fakultas Pertanian-UMJ* 8 November 2017 pp 44–52 (Jakarta: Indonesia)
- [6] Kristanti, H and Tunjung WAS 2013 Detection of alkaloid, flavonoid and terpenoid compounds in bread (*Artocarpus communis* Forst) leaves and pulps In *The 3rd International Conference on Biological Science, Faculty of Biology-Universitas Gadjah Mada* 20–21 2013 (Yogyakarta: Indonesia)

The Effectivity of Jambu Akasia (Bellucia Pentamera Naudin) fruit extract for killing the Aedes aegypti L. Larvae

ORIGINALITY REPORT

18%

SIMILARITY INDEX

15%

INTERNET SOURCES

6%

PUBLICATIONS

17%

STUDENT PAPERS

PRIMARY SOURCES

1	Submitted to Universitas Riau Student Paper	11%
2	eprints.covenantuniversity.edu.ng Internet Source	4%
3	www.e3s-conferences.org Internet Source	2%
4	Shaum Shiyan, Triana Hertiani, Ronny Martien, Akhmad Kharis Nugroho. "OPTIMIZATION AND VALIDATION OF RP-HPLC/UV DETECTION FOR SEVERAL COMPOUNDS SIMULTANEOUSLY IN SEMI-PURIFIED EXTRACT OF WHITE TEA", Rasayan Journal of Chemistry, 2019 Publication	1%

Exclude quotes Off

Exclude bibliography On

Exclude matches < 1%