

Supplementation of Solid Ex-Decanter on Performance of Cattle Fed Palm Fruit By-Products

By Armina Fariani

Supplementation of Solid Ex-Decanter on Performance of Cattle Fed Palm Fruit By-Products

Lili Warly¹, Suyitman¹, Evitayani¹, and A. Fariani²

¹Departement of Animal Nutrition, Faculty of Animal Science, Andalas University
Kampus Unand, Limau Manis, Padang, West Sumatera, Indonesia

²Departement of Animal Nutrition, Fakulty of Agriculture, Sriwijaya University,
South Sumatera, Indonesia

Abstract : The objectives of this study was to evaluate the potential of palm fruit by-products as feed supplement to increase weaned calves performance. Solid ex-decanter were combined with multi mineral and utilized as feed suplement to form a lick block. Ten weaned calves 6 month old (122.34±20.25 kg) were randomly categorized into 2 groups. Control weaned calves receive no solid Ex-Decanter Multi-Nutrient Block (SEDMB) supplementation and treated calf were given continuously. The experiment was conducted at rural farming areas in Darmasraya and calves fed total mix ratio formulated with palm fruit by product based. Measured parameter were daily weight gain (DWG), body condition score (BCS), feed consumption, feed conversion, *in vivo* dry matter and organic matter digestibility. The treatment were significantly different ($p < 0.05$) on daily wight gain, body condition score and feed conversion but did not significantly different ($P < 0.05$) on feed consumption, dry matter and organic matter digestibility. SEDMB supplementation on weaned calves performance of Simbrah Breed showed the effectiveness of feed consumption.

9

Key words : Solid ex-decanter, calves, supplement, palm fruit by-products.

INTRODUCTION

The world's palm oil production was 36.85 million metric tons, while indonesia as lead of oil palm producers in the word with an estimated planted area 6.07 million hectares (USDA report, 2007). The oil palm industry offers two opportunities to promote animal production; firstly feed source from oil palm industry by-product, palm plantation, the foragesin in the inter-rows and oil p¹² frounds after reguler pruning. These are potential feeds source as the yields, palatability and nutritive values are adequate for cattle. The objectives of systematic integration of beef cattle and palm fruit agroindustry by product through optimal use of resouse and also utilized cattle feces as fertilizer to reduce the use of synthetic fertilizers.

Under appropriate conditions and systematic management, cattle can be effectively used for weed control mechanism in oil palm plantation allows the establishments of a harmonious relationship between cattle, the udergrowth

and oil palm. Reduced herbicides usages are environmentally healthy and simultaneously help to reduce total weeding cost through lower volume of chem¹³al use and reduced and extra labor. Reduced herbicides usages means reduced maintenance cost and less environmental contamination and pollution (Azid, 2004). Several study shown that integrated farm system between cattle and oil palm plantation under systematic management shows good potential as a livestock production system (Jalaludin, 1997; 5)ahab, 2003; Devandra and Leng, 2011). Livestock production in developing countries is largely dependent on fibrous feeds-mainly crop residues and low quality pasture that are deficient in protein, minerals and 2)itamins (Makkar, 2007). The advantages of the use of multi-nutritional blocks, in diets based on crop by-products or pastures of typical low quality, are well knows in terms of providing adequate non-protein nitrogen in the rumen, improving both function and efficiency,

which is reflected in higher voluntary intake by the animal and better digestibility fibre. Nutrition programs are requirement to minimize use of purchased mixed rations that the cost is generally higher and finally calves get to consume some feed provide protein and minerals. Later calves provide nutrition to promote some gain but maintain health status.

MATERIALS AND METHODS

Ten weaned calves 6 month old (122.1 ± 19.27 kg) were randomly categorized into 2 groups. Control weaned calves received no Solid Ex-Decanter Multi-Nutrient Block (SEDMB) supplementation and treated calf were given continuously or *ad libitum*. The design used in this study was t-test assuming a two-way where if $p < (+/-) 0.05$ is significantly different results, while if $p < (+/-) 0.05$ then the result is not significantly different. The experiment was conducted in July to September 2012 and calf fed total mix ratio formulated with palm fruit by-product based. Parameters observed were daily weight gain (DWG), body condition score (BCS), feed consumption, feed conversion, *in vivo* dry matter and organic matter digestibility. All calves were weighed before feeding on d-0 and at d-7 intervals thereafter. On weigh days, drinkers were turned off and emptied. Number of days on test required to attain the predetermined market weight was recorded. Total average daily gain (ADG) was computed as the difference between initial body weight (BW) and final BW divided by number of days on test. Each morning the feed bunks were observed and adjustments were made in the amount of residual feed in the feed bunks. Body condition was scored using standard procedures based on a scale of 1 to 9, as describe by Eversole *et al.* (2009). Calves were scored for body condition every 2 weeks.

RESULTS AND DISCUSSION

Daily Weight Gain (DWG) : Beef producers face the challenge of remaining economically viable despite continuously changing paradigms in beef cattle production. Cow calf producers may want to consider early weaning as a management practice when traditional production system cannot effectively address adverse conditions such as lack of forage, unfavorable market trends, noncompetitive weight rates and poor weather conditions.

Weaning weight can be increased by genetic crossbreeding, growth potential and environmental (creep feeding) manipulations or by age of calf (calving earlier and/or weaning later). Production efficiency can be enhanced by using nonharvested forage (Lamb *et al.*, 1996), but nutrient availability in forage can be limiting as forage matures. Nutritional status of cattle grazing mature native range forage can be increased with appropriate supplementation (Kartchner, 19; Adams *et al.*, 1994; Sanson *et al.*, 199).

As shown in table 2, the results of daily weight gain of weaned calves after supplemented by ex-decanter multi-nutrient block (SEDMB) were significantly higher ($p < 0.05$) compared with non supplemented calves (0.563 Vs 0.37 kg/day). Leadley and Sodja (2003) reported that rates of weight gain up through weaning are lost of variations. The lowest rate of gain reported was 0.7 pounds (0.317 kg) per day for 42 days. The highest rate was 1.3 pounds (0.589 kg) per day for 56 days. Fluharty *et al.* (2000) also reported about daily weight gain of weaned calves after 36-7 day after weaning ± 0.8 kg/day on normally weaned (205 day old).

Table 1 : ration composition based on palm agroindustry by-product

No	Feed ingredients	Composition (%)
1	Native pasture	75
2	Solid ex-decanter	10
3	Palm midrib mill	14.5
4	Palm press fibre	0.5
	Total	100

Body condition score (BCS) : Body condition score (BCS) are numbers used to suggest the relative fatness or body composition of the cow. For BCS to be most helpful, producers need to calibrate the system under their own conditions with their own cattle. A nine-grade system is commonly used by researchers in the United states. Body condition was scored using standard procedures based very thin body condition and 9 extreme fatness (Eversole *et al.*, 2009).

SEDMB supplementation on weaned calves were significantly different ($p < 0.05$) than control (Table 3). BCS obtained were in line with increase of daily weight gain of Simbrah breed weaned calves. Fluharty *et al.* (2000) reported body conditional score of weaned calves with normally weaned (205 day old) were 4.4 after 99 day weaned. This score were lower than calves with early weaned (100 day old).

Body condition at weaning also is related to reproductive performance. Bowman and Sowell (1998) reported nine-year summary of data from more than 77,000 cows clearly shows that cows that are thin at weaning are less likely to become pregnant during the following breeding season. Mathis *et al.* (2002) suggested that every beef operation

is different and producers using BCS as a tool should set BCS targets based on their willingness to sort out thin cows at weaning and provide them with additional energy directly after weaning when their requirements are low.

6 Feed consumption and feed conversion :

The key variables affecting the profitability of feedlots are : store cattle purchase price; finished cattle sale price; cost of feed consumed and live-weight gain. This fact sheet deals with feed consumption and live-weight gain. As shown in table 3, the level of consumption on treated weaned calves were not significantly different ($p = 0.294$) control calves were higher average feed consumption (12.71 kg/day). Although the average feed consumption of treated calves are lower than control, the daily weight gain of treated weaned calves shows higher value (Table 2) and gives significantly effect (statistical test). SEDMB supplementation on weaned calves also affecting feed conversion. Higher feed conversion (Table 3) shows on control weaned calves (35.22) than treated weaned calves (23.39). T-test results shows a significantly different ($p = 0.042$). it suggest that solid

Table 2 : Body weight gain of simbrah crossbred weaned calves on integrated farm system between cattle and palm fruit plantation.

Calves	Control			Treatment		
	Initial wight (kg)	Final weight (kg)	Daily weight gain (kg)	Initial eight (kg)	Final weight (kg)	Daily weight gain (kg)
1	87	107	0.408	100	128	0.571
2	110	135	0.510	143	170	0.551
3	119	135	0.326	146	182	0.734
4	125	143	0.367	125	140	0.306
5	116	128	0.244	150	182	0.653
SD	111.4±11.9	129.9±11.2	0.37±0.08	132.8±16.8	16.4±2.3	0.563±0.13*

* : Significantly difference ($p < 0.05$). SD : Standard deviation

Tabel 3. Average body condition score, Feed consumption and Feed Conversion of Simbra breed on integrated farm system between cattle and oil palm plantation

Measured Parameters	Control	Treatment	p-value
---------------------	---------	-----------	---------

Body condition Score	3.00 ^a	3.75 ^a	0.00021*
Feed Consumption, kg	12.71	12.44	0.294 ^{ns}
Feed conversion	35.22	23.39	0.042*

* : Significantly difference ($p < 0.05$), ns : not significantly difference ($p > 0.05$).

10 Body condition score, 1 : several emaciated, 2 : Emaciated, 3 : Very thin, 4 : Thin, 5 : moderate, 6 : Good, 7 : Very good, 8 : Obese, 9 : Very obese.

9
Tabel 4 *In vivo* dry matter and organic matter digestibility

Measured parameters	Control	Treatment	p-value
Average dry matter digestibility, (%)	76.92	74.16	0.095 ^{ns}
Average organic matter digestibility, (%)	63.39	64.48	0.296 ^{ns}

* : Significantly difference (p<0.05), ns : not significantly difference (p>0.05).

ex-decanter multi-nutrient block gives feed efficiency. Calves will instinctively lick a solid ex-decanter multi-nutrient block if needed mineral intake.

Dry matter and organic matter digestibility : Apparent digestibility is a naturally feed digestibility occurring in the digestive tract in the body of the ruminant livestock. The process of digestion occurs in the rumen were assisted by microorganisms in it. Digestion by microorganisms also performed by enzymatic that the enzyme produced by the cells of microorganisms in the rumen (Tillman *et al.*, 1991).

Commodities used in feedlot rations vary considerably in dry matter content (DM). Hay and grain are approximately 90% DM, molasses 75% DM and silage 40% DM. a basic guide for estimating dry matter consumption of feedlot animals is to calculate 2.7 to 3.0% of their live weight (in kilograms). therefore, an animal consuming a grain based diet of 90% DM, would have an estimated intake of fresh feed between 3.0% (2.7% x 100/90) and 3.33% (3.0% x 100/90) of their live weight. The dry matter content of ration refers to the amount of dry material available in a given ration. A number of factors influence the average daily dry matter consumption of lot-fed cattle. These include, live weight (their required maintenance energy requirements), body condition, energy concentration of the ration, health status and ration palatability (Sarah, 2012).

16 Based on the data in table 4, the treatment were not statistically different (p>0.05) on

dry matter and organic matter digestibility. Dry matter digestibility of control weaned calves were higher than treated weaned calves. However, organic matter digestibility of weaned calves supplemented with SEDM 17 were higher even the statistically test were not significantly different (p>0.05). it was rume microbial activity to reach optimal digestibility with lower dry matter intake.

14
Conclusion : Based on the results of the study it can be that the supplementation of solid ex-decanter multi-nutrient block as *ad libitum* can 15 prove weaned calves performance. The treatment were significantly improved daily weight gain, body condition score and feed conversion. However, solid ex-decanter multi-nutrient block did not affect the apparent digestibility of the rotation.

7 ACKNOWLEDGEMENT

Authors are grateful to The Directorate General of Higher Education, the Ministry of Education and Culture Republic of Indonesia for financial support, Grant Number 02/UN.16/PL-USN/2014.

REFERENCE

- Adams, D.C., R.T. Clark, S.A. Coady, J.B. Lamb and M.K. Nielsen, 1994. Extended grazing systems for improving economic returns from Nebraska Sandhills cow/calf operations. *J. Range Manage.* 47:258
- Azid, M.K., 24. Study on Cattle farming at Boreo Samudera Sdn. Bhd (BBSB Report submitted to Borneo Samudera sdn. Bhd. 17 November 2004.
- Bowman, J. And B. Sowell, 1998. Feeding the beef cow herd. In: Kellems, R.O. and D.C. Church (Ed.) *Livestock Feeds and Feeding* (4th Ed.). pp. 243. Prentice-Hall, Upper Saddle River, N.J.
- Devendra, C. And R.A. Leng, 2001. *Feed Resources for Animal in Asia: Issues,*

- Strategies for use, Intensification and Integration for Increased Productivity. Asian-Aust. J. Anim. Sci., 24: 303-321.
- Eversole, D.D., M.F. Browne, J.B. Hall and R.E. Djetz, 2009. Body Condition Scoring Beef Cows. Virginia Polytechnic Institute and State University. Publication, 400-791.
- Fluharty, F.L., S.C. Loerch, T.B. Turner, S.J. Moeller and G.D. Loe, 2000. Effect of weaning age and diet on growth and carcass characteristic in steers. J. Anim. Sci., 78: 1759-1767.
- Jalaludin, S., 1997. Integrated animal production in the oil palm plantation. Malaysian Agriculture University, Selangor.
- Kartchner, R.J., 1980. Effect of protein and energy supplementation of cows grazing native winter range forage on intake and digestibility. J. Anim. Sci., 51:432.
- Lamb, J.B., D.C. Adams, T.J. Klopfenstein, W.W. Stroup and G.P. Lardy, 1996. Range or meadow regrowth and weaning effect on two year-old cows. J. Range Manage. (In press).
- Leandley, S and P. Sodja, 2003. Growth Rates for Preweaned Calves-A review. www.calfnotes.com/pdf/CNCE0898. [3 April 2013].
- Makkar, H.P.S., 2007. Feed supplement block technology-past, present and future. FAO Animal Production and Health. Rome.
- Mathis, C.P., J.E. Sawyer and Parker, 2002. Managing and Feeding Beef Cows Using body Condition Score. Collage of Agriculture and Home Economics. New Mexico University Press.
- Sanson, D.W., D.C. Clanton and I.G. Rush, 1990. Intake and digestion of low-quality meadow hay by steers and performance of cows on native range when fed protein supplements containing various levels of corn. J. Anim. Sci., 68:595.
- Srah, J.F., 2012. Feed Consumption and Live weight Grain. <http://futurebeef.co.au>. [3 April 2013].
- Tillman, A.D., H. Hartadi, S. Reksohadiprodjo, S. Prawirokusumo and S. Lebdoesoekojo, 1991. Ilmu Makanan Ternak Dasar (Fundamental Feed Science). Gadjah Mada University Press, Yogyakarta.
- USDA Statistic Report, 2007. www.IPOB.com.id.
- Wahab, H.A., 2003. Pasture and Sheep Production under Soursop (Annona muricata) Orchard. Malaysian Agriculture Research Development Institute.

Supplementation of Solid Ex-Decanter on Performance of Cattle Fed Palm Fruit By-Products

ORIGINALITY REPORT

19%

SIMILARITY INDEX

PRIMARY SOURCES

1	www.larri.ars.usda.gov Internet	67 words — 3%
2	www.fao.org Internet	53 words — 2%
3	cowbcs.info Internet	48 words — 2%
4	animalscience.tamu.edu Internet	30 words — 1%
5	nexusacademicpublishers.com Internet	24 words — 1%
6	neirong.net Internet	22 words — 1%
7	media.neliti.com Internet	21 words — 1%
8	www.veterinaryworld.org Internet	21 words — 1%
9	repository.unsri.ac.id Internet	19 words — 1%
10	F L Fluharty, S C Loerch, T B Turner, S J Moeller, G D Lowe. "Effects of weaning age and diet on growth and carcass characteristics in steers.", Journal of Animal Science, 2000	18 words — 1%

11	scialert.net Internet	17 words — 1%
12	T S M Widi, N Widyas, R G M F Damai. "Weaning weight of Brahman cross (BX) and Bali cattle under intensive and oil palm plantation-cattle integrated systems", IOP Conference Series: Earth and Environmental Science, 2019 Crossref	15 words — 1%
13	www.ijpaes.com Internet	13 words — 1%
14	agrosym.ues.rs.ba Internet	12 words — 1%
15	linknovate.com Internet	11 words — 1%
16	akademik.unsoed.ac.id Internet	11 words — 1%
17	eprints.undip.ac.id Internet	9 words — < 1%

EXCLUDE QUOTES OFF
EXCLUDE BIBLIOGRAPHY ON

EXCLUDE MATCHES OFF