

2_2130906_AMM V391_Leila Kasum_Ngu_MF_AHPKS_Contr olling CO2 and CH4 emission.pdf

by Muhammad Faizal 18

Submission date: 06-Aug-2018 10:49AM (UTC+0800)

Submission ID: 987823781

File name: 91_Leila_Kasum_Ngu_MF_AHPKS_Controlling_CO2_and_CH4_emission.pdf (378.44K)

Word count: 2576

Character count: 12468

Controlling CO₂ and CH₄ Emission in a Degraded Peat Swamp Forest Related to Water Table and Peat Characteristics

Leila Kalsum^a, Ngudiantoro^b, M.Faizal^c, and A.Halim PKS^d

^aStudent of S-3 Environmental Science Study Program University of Sriwijaya Palembang Indonesia.

^aelak87chem@yahoo.com, ^bngudiantoro@unsri.ac.id, ^cFaisal_ga@yahoo.com, ^dhalimpks@gmail.com

Keywords: CO₂ and CH₄ emission, water table, peat characteristics

Abstract. This study focuses on factors controlling CO₂ and CH₄ emission in a peat swamp forest related to water table and peat characteristics such as peat depth, C-organic, pH, ash content and N-total. This study was conducted in the dry season at a Merang peat swamp forest that has degraded due to logging activities, forest fires and canal opening. Emission of CO₂ and CH₄ was measured by using a closed chamber made by PVC material (60 cm x 60 cm x 40 cm). This close chamber was completed with a fan inside the chamber to stir the gas, a thermometer inside the chamber to measure the gas temperature and a syringe to sample gas. This study has shown that the highest CO₂ emission is at an average of 438.93 mg/m²/hr occurring in land cover type (1) ferns and grasses (open burned area) and the lowest is at average of 44.45 mg/m²/hr in the *gelam* and *belidang*-dominated land. The emission of CH₄ is very low between 0.0018 to 0.0069 mg/m²/hr. the main controlling factor on CO₂ and CH₄ emission is concluded to be the water table. The emission of CO₂ will be greater if water table, pH and C-organic increase.

Introduction

Peatlands are a natural carbon reservoir and cover 3% of the surface area of the earth [1]. Globally, peatlands store approximately 329-525 Giga tonnes (Gt) of carbon, and approximately 86% (455Gt) of them are stored in temperate regions (Canada and Russia), while approximately 14% (70 Gt) is found in the tropics [2]. The peatlands in Indonesia are mainly spread in Papua (7.6 million acres), Sumatra (6.9 million acres), Kalimantan (5.8 million acres) and the rest of approximately 2.2 million acres [3]. The peatlands in the South-Sumatra region reach 1.4 million acres, however the area is gradually decreasing [4]. They cannot be separated from the movement of the world economy with their uses such as industrial timber plantations, agriculture, forestry, conservation, and plantations. However, their management is not easy because they contain complex physical, chemical and hydrological peat issues. The peatlands have long time been considered as carbon sink ecosystem. Related to climate change, they may become carbon source ecosystem with potential to lose carbon either as trace gas carbon dioxide (CO₂) and methane (CH₄) and fluvial dissolved organic carbon [5].

Carbon is one of the peatland elements that undergo recycling in an ecosystem. The carbon cycle is the movement of carbon from the air to the ground and back again into the air. The cycle begins when CO₂ in the atmosphere is absorbed by plants and then converted into carbohydrates through photosynthesis. The photosynthetic reaction will occur if there is sunlight and chlorophyll. Carbohydrates are a source of formation of other organic compounds in plants such as protein and lignin. If the leaves fall or crops die, there will be a process of decomposition and mineralization of soil carbon; they will return to the air in the form of CO₂ gas. Decomposition and mineralization is an overview of enzymatic oxidation and reduction reaction. [6]

1 Greenhouse gas emissions from the peatland consist of CO₂, CH₄, and N₂O. However, CO₂ emissions are much higher than others and CO₂ emissions are strong enough to represent emissions from the peatland, when measuring gas such as CH₄ and N₂O is difficult [7]. According to [8] - [9], CO₂ and CH₄ emission in peatlands are caused by decomposition of carbon compounds under aerobic

and anaerobic condition. The condition of geography, substrate availability and vegetation significantly affect carbon emission [10]. In the exploitation of peatlands, it is usually accompanied by the opening of a canal or drainage, and the drainage in the peat causes changes in the nature of biotic and abiotic degradation and impact on peat degradation [11]. A study shows that the peat soil degradation increases with an increase in pH and ash content [12]. Water table is an important factor in the management of peatlands, a deeper water table will increase carbon emission [13]- [14]- [15]. This study was conducted on the peat swamp forest that has degraded due to logging activities, forest fires and canal opening; it focuses on factors controlling CO_2 and CH_4 emission in the Merang peat swamp forest related to water table and peat characteristics such as peat depth, C-organic, pH, ash content and N-total of peat soil.

Material and Method

Study Site

The study was conducted at the Merang peat swamp forest in South-Sumatra (refer with Fig. 1a). The majority of these peatlands have been damaged by illegal logging activities, canal opening, and forest fires. This study site had previously been identified the level of damage. The peat swamp forest can generally be grouped into three classification: primarily degraded peat swamp forests, secondary peat swamp forests and the peat swamp forest degraded with land cover < 10 % [16]. However, in this study, the classification of the land cover developed more specifically by adapting to the current conditions in the field as follows 1. land dominated by ferns and grasses (open burned area); 2. land dominated by *mahang*; 3. land dominated by *gelam* and *belidang*; 4. land with medium canopy. This classification is in the same line with the conditions of land cover classification by Project MRPP based on its reports [17]. At this location there was a canal located in the middle of the peatland with a depth of approximately 2 meters with an average width of 2.5 meters. This canal was built by illegal loggers for passing their timber from the Merang peat swamp forest. Observation points or sampling points in this study followed the direction of the canal towards the Merang peat swamp forests (refer with Fig. 1b).




Figure 1.a. Location of Merang peat swamp forest, b. Location of sampling points

Measurement of CO_2 and CH_4 Emission

CO_2 and CH_4 emission have been measured by using a closed chamber method. This method was used for measuring the magnitude of emissions from peat soil [18]- [19]- [20]- [21]-[22]-[23]. This study followed the path of the canal. The starting point of each study site was set at 100 meters from the side of the canal. Furthermore, observation points were set at intervals of 200 meters. At each study site, three observation points were established. The close chamber was a size of 60 x 60 x 40 cm. The close chamber was equipped by a fan and a thermometer. When gas emission was measured, the close chamber was placed in the ground and inserted about 3 cm in the soil to prevent a gas leakage. The gas

concentration in the chamber was monitored at any given time. The time intervals for measuring the emissions of CO_2 and CH_4 were a three-minutes period for every 24 minutes [24]. Gaseous emissions were taken from the close chamber by using special tools such as the 10 ml syringe. Furthermore, the gas was analyzed using gas chromatography (GC). Increased concentration of the gas was used to calculate the rate of gas emissions from the soil surface.

Measurement of Water table and Peat Soil Characteristic

Water table measurements were performed using a piezometer to measure the depth of ground water in each of the study area. Peat depth was also measured at each site of study using a drill of peat and drilling was done until it reached the mineral soil layer. Along with the drilling, peat samples were taken at various depths to analyze the characteristics of the peat soil. Each of 500 gram soil samples from the peat drilling was taken and stored in the plastic bags. Furthermore these soil samples were taken to the laboratory for analysis. The analysis in the laboratory was aimed to find out the characteristics of peat soil such as C-organic, N-total, pH and ash content. For measuring C-organic, the Walkey and Black method was used, the pH by using pH H_2O , N-total using the Kjeldhal method and ash content by the loss ignition method.

Result and Discussion


Emission of CO_2 and CH_4

The study results show that the magnitude of emissions of CO_2 and CH_4 at some land cover types and the result of peat analysis (refer with table 1). The highest emission of CO_2 occurred at land cover type 1 or land dominated by fern and grasses as much as $486.39 \text{ mg/m}^2/\text{hr}$ with water table 1.65 m, peat depth 4.8 m, C-organic 45 %, pH 4.22, ash content 33.12 and N-total 0.49 %. The lowest emission occurred at the lower type 3 or land dominated by *gelam* and *belidang* as much as $44.45 \text{ mg/m}^2/\text{hr}$ with water table 0.6 m, peat depth 0.08 m, C-organic 8.42 %, pH 3.2, ash content 60.56 and N-total 0.31 %. However the pattern of CH_4 emission could not be seen clearly. It was obtained that the smaller the water table the greater CH_4 emission was and the magnitude of CH_4 emission was very low, that is, between 0.0018 to $0.0069 \text{ mg/m}^2/\text{hr}$.

Table 1. Peat Emission and Characteristic

Land cover	CH_4 emission, $\text{mg/m}^2/\text{hr}$	CO_2 emission, $\text{mg/m}^2/\text{hr}$	Peat Depth, m	Water Tab. m	C-Org (%)	pH	Ash Cont.	N tot (%)
4	0.0069	117.89	4.3	1.10	30	3.1	15.58	0.7
4	0.0069	116.69	4.25	1.10	30	3.2	0.95	0.97
4	0.0069	119.94	4.6	1.10	32	3.25	13.33	0.67
4	0.0062	181.86	4.45	1.20	35	3.32	17.13	0.77
4	0.0055	228.76	4.35	1.30	36	3.3	12.69	0.76
4	0.0054	226.7	4.5	1.30	35	3.23	15.83	0.75
3	0.0018	44.94	0.1	0.60	8.16	3	57.78	0.35
3	0.0019	44.45	0.08	0.60	8.42	3.2	60.56	0.31
3	0.0031	45.43	0.1	0.60	8.11	3.15	55	0.32
2	0.0068	134.34	2	1.10	33	3.35	46.55	0.4
2	0.0068	128.93	1.99	1.10	32.4	3.3	34.18	0.38
2	0.0065	136.79	1.99	1.15	33	3.2	59.22	0.4
1	0.0034	474.91	4.8	1.60	42.14	4.1	23.84	0.53
1	0.0033	486.39	4.8	1.65	45	4.22	33.12	0.49
1	0.0032	355.49	4.5	1.45	37	4	26	0.49

An overview of the average CO₂ emissions for 4 types of land cover can be seen in Figure 2: the highest emissions occurred in land cover type (1) and the lowest in cover type (3).



1
Figure 2. Emission of CO₂ under different land cover

1 Water Table and Peat Characteristics

The greater the water table, the greater CO₂ emissions increased (refer with table 1); this similarly applied to the C-Organic and pH. Based on statistical analysis, the relationship between CO₂ emissions with water table, C-organic and pH was significant but distance from the channel, ash content and total nitrogen content were. The overall results of the analysis show that at the Merang peat swamp forest the % C organic content was between 8-37%, pH 3.15 - 4.22, ash content 0.95 - 60.5, N total 0.31 - 0.97 %, peat depth between 0.08 - 4.8 m and the water table between 0.6 - 1.65 m. Finally, that most important factor in controlling CO₂ emissions was the water table, followed by C-organic and pH.

Summary

In brief, the water table was found to be a very important controlling factor; it was followed by pH and C-organic content. The water table could be controlled and the exploitation of peats was fruitful if the drainage or canal was not opened excessively. The opening of the canal accelerated the oxidation of the carbon stored in the peatlands.

References

- [1] Limpens, J *et al.* Peatlands and Carbon Cycle: From Local Process to Global Implications. *Biogeosciences*, 5, Hal 1475-1491. (2008)
- [2] Mubekti. Pola Penurunan Tanah di Lahan Gambut pada Berbagai Karakteristik Tanahgambut (Konsolidasi, Kompaksi, Dekomposisi) dan pengaruhnya terhadap emisi karbon. Technical Note. BPPT. (2010)
- [3] Hooijer, A., M Silvius, H Wösten, S Page. Peat-CO₂ Assessment of CO₂ emissions from Drained Peatlands in SE Asia. Delft Hydraulics report **Q3943**. (2006)
- [4] Tajudin, EK. Program dan Skema Penurunan Laju Deforestasi dan Degradasi Hutan Rawa Gambut di Sumatera- Selatan . Workshop ITTO Project RED. Palembang. (2012)
- [5] Michael L.Welloc., Brian Redidy., Cristina. Soil Organic Carbon Stoc. *Forestry Journal*. (2011)
- [6] Kim. H Tan. *Soil physic*. (2000)

-
- [7] Page, *et.al.* Review of Peat Surface Green House Gas emission from Oil Palm Plantations in Southeast Asia. ICCT. (2010)
- [8] Hillel, D. Introduction to Environmental Soil Physics. Elsevier Academic Press. (2004)
- [9] Reddy Ramesh, Biogeochemistry of Wetland, CRC Press. (2008)
- [10] Martikainen. Flux carbon. (1995)
- [11] Szaljak, L *et al.* Chemical Properties of Different Peat Moorsh Soils From The Biebrza River Valley, Agronomy Research 5(2), 165-174. (2007)
- [12] Murayama, S., Z.A, Bakar. Decomposition Of Tropical Peat Soils, Estimation of in Situ Decomposition by Measurement of CO₂ Flux. JARQ 30, 153-158. (1996)
- [13] Hooijer, A., M Silvius, H Wösten, S Page,. Peat-CO₂ Assessment of CO₂ emissions from Drained Peatlands in SE Asia. Delft Hydraulics report **Q3943**. (2006)
- [14] Huistedent, V. J., Van Den Bos, R., Alvares M. Modeling the Effect of water Table management on CO₂ and CH₄ fluxes from peat soils. Netherland Journal of Geosciences : 3-18.(2004)
- [15] Vien, D.M. *et al.* Carbon Dioksida Emission From Peatland in Relation to Hydrology, Peat moisture, humification at the Vo Doi National Park Vietnam. World Congress of Soil Science. Brisbane. Australia. (2010)
- [16] Barkah, S.B. Survey Vegetasi dan Kerusakan Hutan Rawa Gambut Areal MRPP. Laporan Kegiatan Survey. MRPP-GIZ. (2009)
- [17] Solichin. Laporan Merang REDD Pilot Project.(2010)
- [18] Hucinson and Livingstone. Gas Flux Measurement. (2002)
- [19] Pumpanen, J *et al.* Comparison of Different Chamber Technique For Measuring Soil CO₂ Efflux, Agricultural and Forest Meteorology :page 159-176. (2004)
- [20] Widen, B., Lindroth, A. Callibration System For Soil Carbon Dioksida Measurement Chamber: Description and Application, Soil Science Soc, 67, 327-334.(2003)
- [21] Agus, F., *et al.*, 2010. Root Respiration Interferes with Peat CO₂ Emission Measurement. Word Congress of Soil Science. Brisbane, Australia.
- [22] Madset, R. Consideration for Making Chamber –based Soil CO₂ Flux Measurement, World Congress Soil Science, Australia.(2010)
- [23] Page, *et.al.* Review of Peat Surface Green House Gas emission from Oil Palm Plantations in Southeast Asia. ICCT. (2011)
- [24] Saad, N,S. Pengukuran Emisi CO₂ . Balai Penelitian Lingkungan Pertanian Departemen Pertanian. (2007)

2_2130906_AMM V391_Leila
Kasum_Ngu_MF_AHPKS_Controlling CO2 and CH4
emission.pdf

ORIGINALITY REPORT

85%	%	85%	0%
SIMILARITY INDEX	INTERNET SOURCES	PUBLICATIONS	STUDENT PAPERS

PRIMARY SOURCES

1	Kalsum, Leila, null Ngudiantoro, M. Faizal, and A. Halim Pks. "Controlling CO ₂ and CH ₄ Emission in a Degraded Peat Swamp Forest Related to Water Table and Peat Characteristics", Applied Mechanics and Materials, 2013. Publication	85%
---	---	-----

Exclude quotes	On	Exclude matches	< 1 words
Exclude bibliography	On		