

PAPER • OPEN ACCESS

Sriwijaya International Conference on Earth Science and Environmental Issue (ICESEI) 2020

To cite this article: 2021 *IOP Conf. Ser.: Earth Environ. Sci.* **810** 011001

View the [article online](#) for updates and enhancements.

You may also like

- [Diversity and abundance model according to habitat characteristics of filariasis vector, *Mansonia* spp. in Banyuwasin, South Sumatera, Indonesia](#)
Rini Pratiwi, Chairil Anwar, Salni et al.
- [The effect of Leaf Salam Extracts \(*Syzygium polyanthum*\) in diabetes mellitus therapy on wistar albino rats](#)
Nita Parisa, Nia Savitri Tamzil, Debby Handayati Harahap et al.
- [Diversity, dominance and density of Tree Stage Plants of Sriwijaya University Forest Area Indralaya](#)
D Setiawan, H Marisa and D P Indriani



IOP Publishing
**ENVIRONMENTAL
RESEARCH
2021**
A VIRTUAL CONFERENCE
15-19 NOVEMBER

FREE TO
ATTEND

REGISTER
NOW

Sriwijaya International Conference on Earth Science and Environmental Issue (ICESEI) 2020



PREFACE

In the Name of Allah SWT, the Most Gracious, all the praises and thanks be to Him, once more give us the bountiful blessings to establish **Sriwijaya International Conference on Earth Science and Environmental Issue (ICESEI) 2020** formerly known as **The 2nd Sriwijaya International Conference On Environmental Issues 2020 (2nd SRICOENV)** especially in the pandemic Covid-19 which become biggest challenge in all over the world. This conference is organized by Graduate Program Universitas Sriwijaya, Indonesia and co-organized by Thai Nguyen University of Agriculture and Forestry, Vietnam; Graz University of Technology, Austria and Academy of Sciences Malaysia, Malaysia.

The objective of ICESEI 2020 is to perform current state of the arts related to Environmental and Climate Change. All the participants are global partners for future collaboration to exchange new ideas and experiences as their research on "Environment and Climate Change" issue. The conference provides an opportunity for delegations from different institutions to present their ideas and application directly, then build business relationships and research collaborations to create global partnership.

This international conference is a routine agenda of activities at the Graduate Program of Universitas Sriwijaya which initially started in 2018 (1st Sricoenv). Due to Covid-19 pandemic situation which globally occupied, safety and healthiness become priority besides most of traveling and gatherings are strictly prohibited, then the 2nd SRICOENV 2020 is conducted virtually on 21st October 2020.

At this moment, we have to conduct the international conference which flexibly adjusted which expected as the media to extend dissemination process of researchers around the world for their current investigation in the field of Agriculture, Agricultural Technology, Biodiversity, Ecosystem Service and Conservation, Biotechnology and Plant Improvement, Earth Science, Environmental Issue, Forest and Climate Change, Forestry and Environmental Technology, Food Security, Food and Chemical, Soil and Land Science, Food Science, Earth Science, Environmental Science and Environmental Issue. The research results dissemination also hopefully could be known and has positive impact on community.

The Graduate Program of Universitas Sriwijaya as host had holden the ICESEI 2020 from Palembang, South Sumatera-Indonesia. The ICESEI 2020 used ZOOM as platform media. Initially, the presentations are born from keynote speakers for 25-30 minutes each. Then, the ZOOM had split parallelly into four rooms. After that, each room has divided into four sessions. Each speaker has chance to present their research material for 10-15 minutes, then followed by Q&A forums for 15-20 minutes at the end of each session. All questions and answers are delivered directly on ZOOM channel (chat forum). The conference entirely took 9 hours.

We are pleased to inform that the ICESEI 2020 was attended by 200 participants. Those authors and participants came from inside and outside countries such as Austria, Malaysia, Bangladesh, U.S, Japan, Thailand, Philippine, Vietnam, Uganda (9 countries) and various regions in Indonesia (21 of 33 provinces).

The committee selected 63 papers of 94 pre-registered authors, through strict reviewing process to be prepared on publication process. Recently, all the papers have enhanced the quality based on the reviewer's suggestions, to meet the IOP EES requirements.

On behalf of the Graduate Program of Universitas Sriwijaya, we express our sincere gratitude for the participation of keynote speakers, invited speakers, and all sponsors; the Municipality of Palembang, the Belantara Foundation, BPJS Kesehatan, MKTI, APIKI, Ikadil, Bank Sumsel Babel, BNI, BRI. Furthermore, high appreciation to the whole committee team for their excellence in managed and organized all parts of this conference in pandemic circumstance. We also appreciate publisher to publish the proceedings. We hope people will achieve some important information from this proceeding. We re expecting more and more experts and scholars from all over the world to join this international event next year. We look for more scientists and expertise from everywhere to join on next **3rd Sricoenv**.

Palembang, 21 October 2020
Director of Graduate School Universitas Sriwijaya

Prof. Dr. Ir. Amin Rejo, M.P.

Advisory Committee

Anis Saggaff (Universitas Sriwijaya, Indonesia)
Zainuddin Nawawi (Universitas Sriwijaya, Indonesia)
Amin Rejo (Universitas Sriwijaya, Indonesia)
Duong Van Thao (Thai Nguyen University of Agriculture and Forestry, Vietnam)
Zakaria Hossain (Mie University, Japan)

Scientific Committee

Hiroki Tokuda (Nagoya University, Japan)
Prosono (Institut Teknologi Bandung, Indonesia)
Nunung Nuryantono (Institut Pertanian Bogor, Indonesia)
Budi Haryanto (Universitas Indonesia, Indonesia)
Supriharyono (Universitas Diponegoro, Indonesia)
Jatna Supriatna (Universitas Indonesia, Indonesia)
Juahir Hafizan (Universiti Sultan Zainal Abidin, Malaysia)
Philipp Wiedemann (Mannheim University of Applied Sciences, Germany)

Editors

Robbi Rahim
Siti Hanggita Rachmawati
Phan Thi Cam Tu
Nurul Huda
Dessy Adriani
Tengku Zia Ulqodri

Table of contents

[Volume 810](#)

2021

[Previous issue](#)[Next issue](#)

Sriwijaya International Conference on Earth Science and Environmental Issue 21 October 2020, Palembang, Indonesia

Accepted papers received: 23 June 2021

Published online: 09 August 2021

[Open all abstracts, in this issue](#)

Preface

011001

THE FOLLOWING ARTICLE IS OPEN ACCESS

[Sriwijaya International Conference on Earth Science and Environmental Issue \(ICESEI\) 2020](#)

[Open abstract](#), Sriwijaya International Conference on Earth Science and Environmental Issue (ICESEI)

2020 [View article](#), Sriwijaya International Conference on Earth Science and Environmental Issue (ICESEI)

2020 [PDF](#), Sriwijaya International Conference on Earth Science and Environmental Issue (ICESEI) 2020

011002

THE FOLLOWING ARTICLE IS OPEN ACCESS

[Peer review declaration](#)

[Open abstract](#), [Peer review declaration](#) [View article](#), [Peer review declaration](#) [PDF](#), [Peer review declaration](#)

Papers

012001

THE FOLLOWING ARTICLE IS OPEN ACCESS

[Hydroponically Planting Ipomoea Aquatica Vegetables Using Planting Media from Used Goods](#)

M A Nurwahyudi and H Hatta

[Open abstract](#), Hydroponically Planting Ipomoea Aquatica Vegetables Using Planting Media from Used

Goods [View article](#), Hydroponically Planting Ipomoea Aquatica Vegetables Using Planting Media from Used

Goods [PDF](#), Hydroponically Planting Ipomoea Aquatica Vegetables Using Planting Media from Used Goods

012002

THE FOLLOWING ARTICLE IS OPEN ACCESS

[Novel Antihypertension Bioactive Compounds from Rusip](#)

R Rinto, I Widiastuti and B T Samudera

[Open abstract](#), Novel Antihypertension Bioactive Compounds from Rusip [View article](#), Novel

Antihypertension Bioactive Compounds from Rusip [PDF](#), Novel Antihypertension Bioactive Compounds from

Rusip

012003

THE FOLLOWING ARTICLE IS OPEN ACCESS

[Test of Physical Quality Compost and PH of Combination of Water Hyacinth \(*Eichornia Crassipes Mart. Solm*\) and Goat Manure Using Rument Liquid Mol as Activator](#)

A Napoleon, D P Sulistyani and Bakri

[Open abstract](#), Test of Physical Quality Compost and PH of Combination of Water Hyacinth (*Eichornia Crassipes Mart. Solm*) and Goat Manure Using Rument Liquid Mol as Activator [View article](#), Test of Physical Quality Compost and PH of Combination of Water Hyacinth (*Eichornia Crassipes Mart. Solm*) and Goat Manure Using Rument Liquid Mol as Activator [PDF](#), Test of Physical Quality Compost and PH of Combination of Water Hyacinth (*Eichornia Crassipes Mart. Solm*) and Goat Manure Using Rument Liquid Mol as Activator

012004

THE FOLLOWING ARTICLE IS OPEN ACCESS

[Moler Disease of Shallot in the Last Three Years at Brebes Central Java: The Intensity and Resulting Yields Losses is Increasing](#)

Supyani, SH Poromarto, Supriyadi and Hadiwiyono

[Open abstract](#), Moler Disease of Shallot in the Last Three Years at Brebes Central Java: The Intensity and Resulting Yields Losses is Increasing [View article](#), Moler Disease of Shallot in the Last Three Years at Brebes Central Java: The Intensity and Resulting Yields Losses is Increasing [PDF](#), Moler Disease of Shallot in the Last Three Years at Brebes Central Java: The Intensity and Resulting Yields Losses is Increasing

012005

THE FOLLOWING ARTICLE IS OPEN ACCESS

[Phytochemical and Antioxidant Analysis of Fermented 'Dayak' Wild Yam \(*Dioscorea hispida* Dennst\), Purple Yam \(*Dioscorea alata*\) and Air Potato \(*Dioscorea bulbifera* L.\) Tuber Flour](#)

Rudito, Suwanto, Yuli Witono, Bernatal Saragih and Enos Tangke Arung

[Open abstract](#), Phytochemical and Antioxidant Analysis of Fermented 'Dayak' Wild Yam (*Dioscorea hispida* Dennst), Purple Yam (*Dioscorea alata*) and Air Potato (*Dioscorea bulbifera* L.) Tuber Flour [View article](#), Phytochemical and Antioxidant Analysis of Fermented 'Dayak' Wild Yam (*Dioscorea hispida* Dennst), Purple Yam (*Dioscorea alata*) and Air Potato (*Dioscorea bulbifera* L.) Tuber Flour [PDF](#), Phytochemical and Antioxidant Analysis of Fermented 'Dayak' Wild Yam (*Dioscorea hispida* Dennst), Purple Yam (*Dioscorea alata*) and Air Potato (*Dioscorea bulbifera* L.) Tuber Flour

012006

THE FOLLOWING ARTICLE IS OPEN ACCESS

[The Potential of Pure Chitosan in Increasing Eggshell Thickness Silver Arabic Chicken \(*Silver brakel kriel*\)](#)

E Sahara, S Sandi, F Yosi, M L Sari, Riswandi and R Agustina

[Open abstract](#), The Potential of Pure Chitosan in Increasing Eggshell Thickness Silver Arabic Chicken (*Silver brakel kriel*) [View article](#), The Potential of Pure Chitosan in Increasing Eggshell Thickness Silver Arabic Chicken (*Silver brakel kriel*) [PDF](#), The Potential of Pure Chitosan in Increasing Eggshell Thickness Silver Arabic Chicken (*Silver brakel kriel*)

012007

THE FOLLOWING ARTICLE IS OPEN ACCESS

[The Effects of Probiotic from *Hymenachne Acutigluma* Silage in Feed to the Length of Small Intestine and Caeca in Pegagan Ducks](#)

S Sandi, E Sahara, A T Novitasari, A I Munawar Ali, A Susanda, F Yosi and Asmak

[Open abstract](#), [The Effects of Probiotic from *Hymenachne Acutigluma* Silage in Feed to the Length of Small Intestine and Caeca in Pegagan Ducks](#) [View article](#), [The Effects of Probiotic from *Hymenachne Acutigluma* Silage in Feed to the Length of Small Intestine and Caeca in Pegagan Ducks](#) [PDF](#), [The Effects of Probiotic from *Hymenachne Acutigluma* Silage in Feed to the Length of Small Intestine and Caeca in Pegagan Ducks](#)

012008

THE FOLLOWING ARTICLE IS OPEN ACCESS

[Physical and Chemical Quality of Bento Rayap Grass Haylage \(*Leersia Hexandra*\)](#)

[Supplemented with Water Mimmosa \(*Neptunia Oleracea*\)](#)

Riswandi, A I M Ali, Muhakka, A Imsya, A Abrar, E Sahara, B Hamzah, A Supriadi and P Alpian

[Open abstract](#), [Physical and Chemical Quality of Bento Rayap Grass Haylage \(*Leersia Hexandra*\) Supplemented with Water Mimmosa \(*Neptunia Oleracea*\)](#) [View article](#), [Physical and Chemical Quality of Bento Rayap Grass Haylage \(*Leersia Hexandra*\) Supplemented with Water Mimmosa \(*Neptunia Oleracea*\)](#) [PDF](#), [Physical and Chemical Quality of Bento Rayap Grass Haylage \(*Leersia Hexandra*\) Supplemented with Water Mimmosa \(*Neptunia Oleracea*\)](#)

012009

THE FOLLOWING ARTICLE IS OPEN ACCESS

[Phylogenetic of Featherback *Chitala sp* from South Sumatra Based on Cytochrome C Oxidase Subunit I \(COI\) Gene](#)

Y Anjarsari, M Syaifudin, D Jubaedah, F H Taqwa and D Yonarta

[Open abstract](#), [Phylogenetic of Featherback *Chitala sp* from South Sumatra Based on Cytochrome C Oxidase Subunit I \(COI\) Gene](#) [View article](#), [Phylogenetic of Featherback *Chitala sp* from South Sumatra Based on Cytochrome C Oxidase Subunit I \(COI\) Gene](#) [PDF](#), [Phylogenetic of Featherback *Chitala sp* from South Sumatra Based on Cytochrome C Oxidase Subunit I \(COI\) Gene](#)

012010

THE FOLLOWING ARTICLE IS OPEN ACCESS

[Farmer's Perception of Climate Change and the Impacts on Livelihood in South Sulawesi](#)

Arifah, D Salman, A Yassi and EB Demmallino

[Open abstract](#), [Farmer's Perception of Climate Change and the Impacts on Livelihood in South Sulawesi](#) [View article](#), [Farmer's Perception of Climate Change and the Impacts on Livelihood in South Sulawesi](#) [PDF](#), [Farmer's Perception of Climate Change and the Impacts on Livelihood in South Sulawesi](#)

012011

THE FOLLOWING ARTICLE IS OPEN ACCESS

[The Correlation of Climatic Factors with Incidence of Dengue Hemorrhagic Fever in Palembang Bari General Hospital](#)

A Ghiffari, C Anwar, M Soleha, T Prameswarie and D N Anggina

[Open abstract](#), [The Correlation of Climatic Factors with Incidence of Dengue Hemorrhagic Fever in Palembang Bari General Hospital](#) [View article](#), [The Correlation of Climatic Factors with Incidence of Dengue Hemorrhagic Fever in Palembang Bari General Hospital](#) [PDF](#), [The Correlation of Climatic Factors with Incidence of Dengue Hemorrhagic Fever in Palembang Bari General Hospital](#)

012012

THE FOLLOWING ARTICLE IS OPEN ACCESS

[Sustainability and Existential Risks with Special Reference to Climate Change - An Overview](#)

M B Awang

[Open abstract](#), Sustainability and Existential Risks with Special Reference to Climate Change - An Overview [View article](#), Sustainability and Existential Risks with Special Reference to Climate Change - An Overview [PDF](#), Sustainability and Existential Risks with Special Reference to Climate Change - An Overview

012013

THE FOLLOWING ARTICLE IS OPEN ACCESS

[Effect of Flue Gas Oxygen Content to Gas-Gas Heater Requirement in Limestone Forced Oxidation Desulfurizer System](#)

A M Reza, A B Heksaprilla, F Chariri, U Tasyrifah, M B A Gani and P D Suwondo

[Open abstract](#), Effect of Flue Gas Oxygen Content to Gas-Gas Heater Requirement in Limestone Forced Oxidation Desulfurizer System [View article](#), Effect of Flue Gas Oxygen Content to Gas-Gas Heater Requirement in Limestone Forced Oxidation Desulfurizer System [PDF](#), Effect of Flue Gas Oxygen Content to Gas-Gas Heater Requirement in Limestone Forced Oxidation Desulfurizer System

012014

THE FOLLOWING ARTICLE IS OPEN ACCESS

[Waste-to-Energy \(WTE\) Method to Mitigate Harmful Environmental and Health Consequences Due to LDPE Plastic Waste](#)

Novarini, S Kurniawan, Rusdianasari and Y Bow

[Open abstract](#), Waste-to-Energy (WTE) Method to Mitigate Harmful Environmental and Health Consequences Due to LDPE Plastic Waste [View article](#), Waste-to-Energy (WTE) Method to Mitigate Harmful Environmental and Health Consequences Due to LDPE Plastic Waste [PDF](#), Waste-to-Energy (WTE) Method to Mitigate Harmful Environmental and Health Consequences Due to LDPE Plastic Waste

012015

THE FOLLOWING ARTICLE IS OPEN ACCESS

[Increase the Toughness of Material of Construction Using Heat Treatment](#)

H Chandra, Warizal, H Dahlan, N Paramitha Eka Utami and A Tito

[Open abstract](#), Increase the Toughness of Material of Construction Using Heat Treatment [View article](#), Increase the Toughness of Material of Construction Using Heat Treatment [PDF](#), Increase the Toughness of Material of Construction Using Heat Treatment

012016

THE FOLLOWING ARTICLE IS OPEN ACCESS

[Analysis of the Effect of Stop Drilled Hole Diameter Variation on Fatigue Resistance in Medium Carbon Steel DIN HQ 705](#)

H Chandra and V Lestari

[Open abstract](#), Analysis of the Effect of Stop Drilled Hole Diameter Variation on Fatigue Resistance in Medium Carbon Steel DIN HQ 705 [View article](#), Analysis of the Effect of Stop Drilled Hole Diameter Variation on Fatigue Resistance in Medium Carbon Steel DIN HQ 705 [PDF](#), Analysis of the Effect of Stop Drilled Hole Diameter Variation on Fatigue Resistance in Medium Carbon Steel DIN HQ 705

012017

THE FOLLOWING ARTICLE IS OPEN ACCESS

[The Portrait of Women's Participation on Inland Fishery Activities in Lao PDR](#)

D Muthmainnah and A H Rais

[Open abstract](#), [The Portrait of Women's Participation on Inland Fishery Activities in Lao PDR](#) [View article](#), [The Portrait of Women's Participation on Inland Fishery Activities in Lao PDR](#) [PDF](#), [The Portrait of Women's Participation on Inland Fishery Activities in Lao PDR](#)

012018

THE FOLLOWING ARTICLE IS OPEN ACCESS

[Community Empowerment Model to Flooding Risk Reduction in Palembang City \(A Case Study of Gotong Royong Program\)](#)

A Sandy, Sriati, Azhar and A Siswanto

[Open abstract](#), [Community Empowerment Model to Flooding Risk Reduction in Palembang City \(A Case Study of Gotong Royong Program\)](#) [View article](#), [Community Empowerment Model to Flooding Risk Reduction in Palembang City \(A Case Study of Gotong Royong Program\)](#) [PDF](#), [Community Empowerment Model to Flooding Risk Reduction in Palembang City \(A Case Study of Gotong Royong Program\)](#)

012019

THE FOLLOWING ARTICLE IS OPEN ACCESS

[Model of Environmental Caring Character Development in SMA Lubuklinggau City](#)

Warah, Alfitri, Z Arifin and Riswani

[Open abstract](#), [Model of Environmental Caring Character Development in SMA Lubuklinggau City](#) [View article](#), [Model of Environmental Caring Character Development in SMA Lubuklinggau City](#) [PDF](#), [Model of Environmental Caring Character Development in SMA Lubuklinggau City](#)

012020

THE FOLLOWING ARTICLE IS OPEN ACCESS

[Placement of the Temples Site in Wetlands \(Case Study in Bumiayu Temples Site\)](#)

S M Siregar, E Sutriyono, A Siswanto and A A Munandar

[Open abstract](#), [Placement of the Temples Site in Wetlands \(Case Study in Bumiayu Temples Site\)](#) [View article](#), [Placement of the Temples Site in Wetlands \(Case Study in Bumiayu Temples Site\)](#) [PDF](#), [Placement of the Temples Site in Wetlands \(Case Study in Bumiayu Temples Site\)](#)

012021

THE FOLLOWING ARTICLE IS OPEN ACCESS

[Local Wisdom of the Wetland Swamps Agricultural System for a Sustainable Environment](#)

I M Sakir, Sriati, A Saptawan and R Juniah

[Open abstract](#), [Local Wisdom of the Wetland Swamps Agricultural System for a Sustainable Environment](#) [View article](#), [Local Wisdom of the Wetland Swamps Agricultural System for a Sustainable Environment](#) [PDF](#), [Local Wisdom of the Wetland Swamps Agricultural System for a Sustainable Environment](#)

012022

THE FOLLOWING ARTICLE IS OPEN ACCESS

[The Shape of Forest, Social and Economic Conditions of Community Living Around Production Forest with Industrial Plantation Forest Permit \(Case Study: Forest Management Unit of KPH Meranti\)](#)

Y Hidayat, Alfitri, D H Purnama and Riswani

[Open abstract](#), [The Shape of Forest, Social and Economic Conditions of Community Living Around Production Forest with Industrial Plantation Forest Permit \(Case Study: Forest Management Unit of KPH Meranti\)](#)

Meranti) [View article](#), The Shape of Forest, Social and Economic Conditions of Community Living Around Production Forest with Industrial Plantation Forest Permit (Case Study: Forest Management Unit of KPH Meranti) [PDF](#), The Shape of Forest, Social and Economic Conditions of Community Living Around Production Forest with Industrial Plantation Forest Permit (Case Study: Forest Management Unit of KPH Meranti)

012023

THE FOLLOWING ARTICLE IS OPEN ACCESS

[The Response of Water Level Depth to Vegetation Composition in Degraded Peatland: a Case Study of Sriwijaya Wetland Botanical Garden, Indonesia](#)

S Maryani and D Novriady

[Open abstract](#), The Response of Water Level Depth to Vegetation Composition in Degraded Peatland: a Case Study of Sriwijaya Wetland Botanical Garden, Indonesia [View article](#), The Response of Water Level Depth to Vegetation Composition in Degraded Peatland: a Case Study of Sriwijaya Wetland Botanical Garden, Indonesia [PDF](#), The Response of Water Level Depth to Vegetation Composition in Degraded Peatland: a Case Study of Sriwijaya Wetland Botanical Garden, Indonesia

012024

THE FOLLOWING ARTICLE IS OPEN ACCESS

[The Effect of Household's Age Structure to Waste Generation in Palembang City](#)

D Novriady, O Komalasari, H Hatta and R Oktarina

[Open abstract](#), The Effect of Household's Age Structure to Waste Generation in Palembang City [View article](#), The Effect of Household's Age Structure to Waste Generation in Palembang City [PDF](#), The Effect of Household's Age Structure to Waste Generation in Palembang City

012025

THE FOLLOWING ARTICLE IS OPEN ACCESS

[The Implementation of Character Building to Improve Resident Participation in Waste Management](#)

R S Y Zebua and Sunarti

[Open abstract](#), The Implementation of Character Building to Improve Resident Participation in Waste Management [View article](#), The Implementation of Character Building to Improve Resident Participation in Waste Management [PDF](#), The Implementation of Character Building to Improve Resident Participation in Waste Management

012026

THE FOLLOWING ARTICLE IS OPEN ACCESS

[Greenhouse Gas Inventory Standard for Cities: a Case of Jakarta](#)

RG Dewi, UWR Siagian, R Parinderati, GN Sevie, I Hendrawan, D Wati and R Rakhmana

[Open abstract](#), Greenhouse Gas Inventory Standard for Cities: a Case of Jakarta [View article](#), Greenhouse Gas Inventory Standard for Cities: a Case of Jakarta [PDF](#), Greenhouse Gas Inventory Standard for Cities: a Case of Jakarta

012027

THE FOLLOWING ARTICLE IS OPEN ACCESS

[Perceived Impact of Oil Palm Plantation on Agriculture and Environment](#)

H Saediman, I S Mboe, M Zani, L Daud and A M Padangaran

[Open abstract](#), Perceived Impact of Oil Palm Plantation on Agriculture and Environment [View article](#), Perceived Impact of Oil Palm Plantation on Agriculture and Environment [PDF](#), Perceived Impact of Oil Palm Plantation on Agriculture and Environment

012028

THE FOLLOWING ARTICLE IS OPEN ACCESS

[Gender Role in Farmers' Livelihood Strategies at Peatland Area of Fire-Prone in Ogan Komering Ilir Regency South Sumatra Province](#)

A Nurlia, M Rahmat, E A Waluyo, D H Purnama and Sabaruddin

[Open abstract](#), [Gender Role in Farmers' Livelihood Strategies at Peatland Area of Fire-Prone in Ogan Komering Ilir Regency South Sumatra Province](#) [View article](#), [Gender Role in Farmers' Livelihood Strategies at Peatland Area of Fire-Prone in Ogan Komering Ilir Regency South Sumatra Province](#) [PDF](#), [Gender Role in Farmers' Livelihood Strategies at Peatland Area of Fire-Prone in Ogan Komering Ilir Regency South Sumatra Province](#)

012029

THE FOLLOWING ARTICLE IS OPEN ACCESS

[Science of Climate Change in Agricultural Courses](#)

I P Handayani

[Open abstract](#), [Science of Climate Change in Agricultural Courses](#) [View article](#), [Science of Climate Change in Agricultural Courses](#) [PDF](#), [Science of Climate Change in Agricultural Courses](#)

012030

THE FOLLOWING ARTICLE IS OPEN ACCESS

[Interlinkage of Poverty-Hunger-Health-Wellbeing As Critically Seen For Effective Policy Anticipation on the Need for Wetland Ecosystem Reclamation: "An Argument on the Importance of Truly Sincere & Much Intense International Cooperation to Tackle Future Regional Economic Shortcoming"](#)

F Sjarkowi

[Open abstract](#), [Interlinkage of Poverty-Hunger-Health-Wellbeing As Critically Seen For Effective Policy Anticipation on the Need for Wetland Ecosystem Reclamation: "An Argument on the Importance of Truly Sincere & Much Intense International Cooperation to Tackle Future Regional Economic Shortcoming"](#) [View article](#), [Interlinkage of Poverty-Hunger-Health-Wellbeing As Critically Seen For Effective Policy Anticipation on the Need for Wetland Ecosystem Reclamation: "An Argument on the Importance of Truly Sincere & Much Intense International Cooperation to Tackle Future Regional Economic Shortcoming"](#) [PDF](#), [Interlinkage of Poverty-Hunger-Health-Wellbeing As Critically Seen For Effective Policy Anticipation on the Need for Wetland Ecosystem Reclamation: "An Argument on the Importance of Truly Sincere & Much Intense International Cooperation to Tackle Future Regional Economic Shortcoming"](#)

012031

THE FOLLOWING ARTICLE IS OPEN ACCESS

[Covid-19: People Activity, Monitoring Development and Environmental Issue](#)

Pantri Heriyati and Suryadiputra Liawatimena

[Open abstract](#), [Covid-19: People Activity, Monitoring Development and Environmental Issue](#) [View article](#), [Covid-19: People Activity, Monitoring Development and Environmental Issue](#) [PDF](#), [Covid-19: People Activity, Monitoring Development and Environmental Issue](#)

012032

THE FOLLOWING ARTICLE IS OPEN ACCESS

[Informality in Urban Water Supply: A Case of Tambaklorok, Semarang](#)

N R Putri and J S Setyono

[Open abstract](#), [Informality in Urban Water Supply: A Case of Tambaklorok, Semarang](#) [View article](#), [Informality in Urban Water Supply: A Case of Tambaklorok, Semarang](#) [PDF](#), [Informality in Urban Water Supply: A Case of Tambaklorok, Semarang](#)

012033

THE FOLLOWING ARTICLE IS OPEN ACCESS

[Geo Location Dataset Analysis for Identifying Fertilizer Dealers in Vellore district Tamilnadu, India](#)

Rajkumar Rajasekaran, Thipireddy Rishith Reddy, Palnati Gowtham Reddy and Jolly Masih
[Open abstract](#), [Geo Location Dataset Analysis for Identifying Fertilizer Dealers in Vellore district Tamilnadu, India](#) [View article](#), [Geo Location Dataset Analysis for Identifying Fertilizer Dealers in Vellore district Tamilnadu, India](#) [PDF](#), [Geo Location Dataset Analysis for Identifying Fertilizer Dealers in Vellore district Tamilnadu, India](#)

012034

THE FOLLOWING ARTICLE IS OPEN ACCESS

[The Differences of Nutritional Status of Children Living in Riverbank and in Mainland of Pedamaran I District of Ogan Komering Ilir Regency](#)

Nuryanto, R Oktarina, O Komalasari and O Juairiyah
[Open abstract](#), [The Differences of Nutritional Status of Children Living in Riverbank and in Mainland of Pedamaran I District of Ogan Komering Ilir Regency](#) [View article](#), [The Differences of Nutritional Status of Children Living in Riverbank and in Mainland of Pedamaran I District of Ogan Komering Ilir Regency](#) [PDF](#), [The Differences of Nutritional Status of Children Living in Riverbank and in Mainland of Pedamaran I District of Ogan Komering Ilir Regency](#)

012035

THE FOLLOWING ARTICLE IS OPEN ACCESS

[The Potential Contribution of Senior Citizens' Grant \(SCG\) Towards Nutrition Outcomes in Older Persons' Households in Uganda](#)

J Nambooze, M Mbabazi, L N Nyanzi, J Kakande, S Nahalamba and H Mayengo
[Open abstract](#), [The Potential Contribution of Senior Citizens' Grant \(SCG\) Towards Nutrition Outcomes in Older Persons' Households in Uganda](#) [View article](#), [The Potential Contribution of Senior Citizens' Grant \(SCG\) Towards Nutrition Outcomes in Older Persons' Households in Uganda](#) [PDF](#), [The Potential Contribution of Senior Citizens' Grant \(SCG\) Towards Nutrition Outcomes in Older Persons' Households in Uganda](#)

012036

THE FOLLOWING ARTICLE IS OPEN ACCESS

[Study of Black Cumin Seed Oil \(BCSO\) \(*Nigella sativa* L.\) as an Immunomodulator in The Healthy Active Smoker Volunteer](#)

T Hidayati, A Akrom, L Apriani, Indrayanti and S Sun
[Open abstract](#), [Study of Black Cumin Seed Oil \(BCSO\) \(*Nigella sativa* L.\) as an Immunomodulator in The Healthy Active Smoker Volunteer](#) [View article](#), [Study of Black Cumin Seed Oil \(BCSO\) \(*Nigella sativa* L.\) as an Immunomodulator in The Healthy Active Smoker Volunteer](#) [PDF](#), [Study of Black Cumin Seed Oil \(BCSO\) \(*Nigella sativa* L.\) as an Immunomodulator in The Healthy Active Smoker Volunteer](#)

012037

THE FOLLOWING ARTICLE IS OPEN ACCESS

[Ethanollic Extract of Black Cumin Seed Reduced Radical Reactive from Dimethylbenzanthracene Compounds](#)

A Akrom, Mustofa and L H Nurani
[Open abstract](#), [Ethanollic Extract of Black Cumin Seed Reduced Radical Reactive from Dimethylbenzanthracene Compounds](#) [View article](#), [Ethanollic Extract of Black Cumin Seed Reduced Radical Reactive from Dimethylbenzanthracene Compounds](#) [PDF](#), [Ethanollic Extract of Black Cumin Seed Reduced Radical Reactive from Dimethylbenzanthracene Compounds](#)

012038

THE FOLLOWING ARTICLE IS OPEN ACCESS

[Temulawak \(*Curcuma xanthorrhiza*\) Extract as a Cancer Chemopreventive Agent Via Up-Regulation p53 and Caspase-3 Gene](#)

I Indrayanti, K A Kamila, B A Hernowo, F A Haq and A Akrom

[Open abstract](#), [Temulawak \(*Curcuma xanthorrhiza*\) Extract as a Cancer Chemopreventive Agent Via Up-Regulation p53 and Caspase-3 Gene](#) [View article](#), [Temulawak \(*Curcuma xanthorrhiza*\) Extract as a Cancer Chemopreventive Agent Via Up-Regulation p53 and Caspase-3 Gene](#) [PDF](#), [Temulawak \(*Curcuma xanthorrhiza*\) Extract as a Cancer Chemopreventive Agent Via Up-Regulation p53 and Caspase-3 Gene](#)

012039

THE FOLLOWING ARTICLE IS OPEN ACCESS

[Jamu Reduce Oxidative Stress from Active Smokers in a Rural Area of Yogyakarta](#)

E Darmawan, A Akrom, E F Lerebulan and A Adnan

[Open abstract](#), [Jamu Reduce Oxidative Stress from Active Smokers in a Rural Area of Yogyakarta](#) [View article](#), [Jamu Reduce Oxidative Stress from Active Smokers in a Rural Area of Yogyakarta](#) [PDF](#), [Jamu Reduce Oxidative Stress from Active Smokers in a Rural Area of Yogyakarta](#)

012040

THE FOLLOWING ARTICLE IS OPEN ACCESS

[Antioxidant Profile of Heartwood and Sapwood of *Caesalpinia sappan* L. Tree's Part Grown in Imogiri Nature Preserve, Yogyakarta](#)

I S Arsiningtyas

[Open abstract](#), [Antioxidant Profile of Heartwood and Sapwood of *Caesalpinia sappan* L. Tree's Part Grown in Imogiri Nature Preserve, Yogyakarta](#) [View article](#), [Antioxidant Profile of Heartwood and Sapwood of *Caesalpinia sappan* L. Tree's Part Grown in Imogiri Nature Preserve, Yogyakarta](#) [PDF](#), [Antioxidant Profile of Heartwood and Sapwood of *Caesalpinia sappan* L. Tree's Part Grown in Imogiri Nature Preserve, Yogyakarta](#)

012041

THE FOLLOWING ARTICLE IS OPEN ACCESS

[Isolation and Identification of Phosphate Solubilising Bacteria from Potato Rhizosphere on Andisol](#)

A E Marpaung and D N Susilowati

[Open abstract](#), [Isolation and Identification of Phosphate Solubilising Bacteria from Potato Rhizosphere on Andisol](#) [View article](#), [Isolation and Identification of Phosphate Solubilising Bacteria from Potato Rhizosphere on Andisol](#) [PDF](#), [Isolation and Identification of Phosphate Solubilising Bacteria from Potato Rhizosphere on Andisol](#)

012042

THE FOLLOWING ARTICLE IS OPEN ACCESS

[Hygiene and Sanitation on Post-Harvest Coffee Processing in the South OKU Regency Community \(Case Study\)](#)

D Lilia, N Damiri, M Zulkarnain and Mulawarman

[Open abstract](#), [Hygiene and Sanitation on Post-Harvest Coffee Processing in the South OKU Regency Community \(Case Study\)](#) [View article](#), [Hygiene and Sanitation on Post-Harvest Coffee Processing in the South OKU Regency Community \(Case Study\)](#) [PDF](#), [Hygiene and Sanitation on Post-Harvest Coffee Processing in the South OKU Regency Community \(Case Study\)](#)

012043

THE FOLLOWING ARTICLE IS OPEN ACCESS

[Protein Energy Deficiency Increases the Risk of Anemia in Pregnant Women](#)

D K Mulyantoro and I Kusri

[Open abstract](#), Protein Energy Deficiency Increases the Risk of Anemia in Pregnant Women [View article](#), Protein Energy Deficiency Increases the Risk of Anemia in Pregnant Women [PDF](#), Protein Energy Deficiency Increases the Risk of Anemia in Pregnant Women

012044

THE FOLLOWING ARTICLE IS OPEN ACCESS

[Modification of Gluten-Free Starch and the Application on Cookies](#)

Nursanty, D Yesi and Y Sugiarti

[Open abstract](#), Modification of Gluten-Free Starch and the Application on Cookies [View article](#), Modification of Gluten-Free Starch and the Application on Cookies [PDF](#), Modification of Gluten-Free Starch and the Application on Cookies

012045

THE FOLLOWING ARTICLE IS OPEN ACCESS

[Relationship between Particulate Matter and Population of Airborne Microorganism in Goat Farm: A case study at Ladang Pasir Akar, Terengganu](#)

A A Mazlan, A Azid, M S Samsudin, M A Razik and S M Shahrudin

[Open abstract](#), Relationship between Particulate Matter and Population of Airborne Microorganism in Goat Farm: A case study at Ladang Pasir Akar, Terengganu [View article](#), Relationship between Particulate Matter and Population of Airborne Microorganism in Goat Farm: A case study at Ladang Pasir Akar, Terengganu [PDF](#), Relationship between Particulate Matter and Population of Airborne Microorganism in Goat Farm: A case study at Ladang Pasir Akar, Terengganu

012046

THE FOLLOWING ARTICLE IS OPEN ACCESS

[Prospect of Low Land Rice Sustainability in South Sumatera, Indonesia](#)

M Antoni, I Zahri, D Adriani and A K A Thoni

[Open abstract](#), Prospect of Low Land Rice Sustainability in South Sumatera, Indonesia [View article](#), Prospect of Low Land Rice Sustainability in South Sumatera, Indonesia [PDF](#), Prospect of Low Land Rice Sustainability in South Sumatera, Indonesia

012047

THE FOLLOWING ARTICLE IS OPEN ACCESS

[Food as a Basis for Good Health and Well-Being](#)

M Murkovic

[Open abstract](#), Food as a Basis for Good Health and Well-Being [View article](#), Food as a Basis for Good Health and Well-Being [PDF](#), Food as a Basis for Good Health and Well-Being

012048

THE FOLLOWING ARTICLE IS OPEN ACCESS

[Kinetic Study of Removal of Congo Red and Direct Green](#)

Muhammad Said, Lavini Indwi Saputri, Hasanudin, Poedji Loekitowati Hariani and Ng Law Yong

[Open abstract](#), Kinetic Study of Removal of Congo Red and Direct Green [View article](#), Kinetic Study of Removal of Congo Red and Direct Green [PDF](#), Kinetic Study of Removal of Congo Red and Direct Green

012049

THE FOLLOWING ARTICLE IS OPEN ACCESS

[Effectiveness of Proteins and Supernatants Isolated from *Bacillus Thuringiensis*-Based Bio-Insecticides Against Termites *Macrotermes Gilvus* \(Isoptera: Termitidae\)](#)

Y Pujiastuti, A Riskal, Suparman, A Arsi, B Gunawan and D P Sulistyani

[Open abstract](#), Effectiveness of Proteins and Supernatants Isolated from *Bacillus Thuringiensis*-Based Bio-Insecticides Against Termites *Macrotermes Gilvus* (Isoptera: Termitidae) [View article](#), Effectiveness of Proteins and Supernatants Isolated from *Bacillus Thuringiensis*-Based Bio-Insecticides Against Termites *Macrotermes Gilvus* (Isoptera: Termitidae) [PDF](#), Effectiveness of Proteins and Supernatants Isolated from *Bacillus Thuringiensis*-Based Bio-Insecticides Against Termites *Macrotermes Gilvus* (Isoptera: Termitidae)

012050

THE FOLLOWING ARTICLE IS OPEN ACCESS

[Fitness of Paraquat Resistant and Susceptible Biotypes Population of Goosegrass \(*Eleusine indica* \(L.\) Gaetrn.\)](#)

B S Napitupulu, E Purba and Mukhlis

[Open abstract](#), Fitness of Paraquat Resistant and Susceptible Biotypes Population of Goosegrass (*Eleusine indica* (L.) Gaetrn.) [View article](#), Fitness of Paraquat Resistant and Susceptible Biotypes Population of Goosegrass (*Eleusine indica* (L.) Gaetrn.) [PDF](#), Fitness of Paraquat Resistant and Susceptible Biotypes Population of Goosegrass (*Eleusine indica* (L.) Gaetrn.)

012051

THE FOLLOWING ARTICLE IS OPEN ACCESS

[Source of Carbon Dioxide Emissions Equivalent to Plant-Based Control Concept with Dynamic System in Prabumulih City, South Sumatra Province](#)

Y Pratiwi, A Rejo, A Fariani and M Faizal

[Open abstract](#), Source of Carbon Dioxide Emissions Equivalent to Plant-Based Control Concept with Dynamic System in Prabumulih City, South Sumatra Province [View article](#), Source of Carbon Dioxide Emissions Equivalent to Plant-Based Control Concept with Dynamic System in Prabumulih City, South Sumatra Province [PDF](#), Source of Carbon Dioxide Emissions Equivalent to Plant-Based Control Concept with Dynamic System in Prabumulih City, South Sumatra Province

012052

THE FOLLOWING ARTICLE IS OPEN ACCESS

[Source of apportionment of Air Quality Parameters at Federal Port of Malaysia with Emphasis on Ship Emission](#)

M S Samsudin, A Azid, M A Razik, M A Zaudi and S M Shaharudin

[Open abstract](#), Source of apportionment of Air Quality Parameters at Federal Port of Malaysia with Emphasis on Ship Emission [View article](#), Source of apportionment of Air Quality Parameters at Federal Port of Malaysia with Emphasis on Ship Emission [PDF](#), Source of apportionment of Air Quality Parameters at Federal Port of Malaysia with Emphasis on Ship Emission

012053

THE FOLLOWING ARTICLE IS OPEN ACCESS

[Erosion Measurement by Using Rainfall Simulator at Grass Soil and After Harvested Soil in Wanagama](#)

A Kusumandari, M C Satriagasa, R Hadi Purwanto and W T Widayanti

[Open abstract](#), Erosion Measurement by Using Rainfall Simulator at Grass Soil and After Harvested Soil in Wanagama [View article](#), Erosion Measurement by Using Rainfall Simulator at Grass Soil and After Harvested

Soil in Wanagama [PDF](#), Erosion Measurement by Using Rainfall Simulator at Grass Soil and After Harvested Soil in Wanagama

012054

THE FOLLOWING ARTICLE IS OPEN ACCESS

[Application of parijoto \(*medinilla speciosa* L.\) extract as body lotion](#)

A Kusumastuti and H S Rahma

[Open abstract](#), Application of parijoto (*medinilla speciosa* L.) extract as body lotion [View article](#), Application of parijoto (*medinilla speciosa* L.) extract as body lotion [PDF](#), Application of parijoto (*medinilla speciosa* L.) extract as body lotion

012055

THE FOLLOWING ARTICLE IS OPEN ACCESS

[Evaluation of Nutritional Content of Beef Rendang Using Wet and Dry Seasonings](#)

Anni Faridah and Rahmi Holinesti

[Open abstract](#), Evaluation of Nutritional Content of Beef Rendang Using Wet and Dry Seasonings [View article](#), Evaluation of Nutritional Content of Beef Rendang Using Wet and Dry Seasonings [PDF](#), Evaluation of Nutritional Content of Beef Rendang Using Wet and Dry Seasonings

012056

THE FOLLOWING ARTICLE IS OPEN ACCESS

[Fabric Coloring using Cuscuta as A Form of Environmental Sustainability](#)

Atika, Rodia Syamwil, Widowati, Siti Nurrohmah and Putri Zaida Zulfa

[Open abstract](#), Fabric Coloring using Cuscuta as A Form of Environmental Sustainability [View article](#), Fabric Coloring using Cuscuta as A Form of Environmental Sustainability [PDF](#), Fabric Coloring using Cuscuta as A Form of Environmental Sustainability

012057

THE FOLLOWING ARTICLE IS OPEN ACCESS

[Formulation and Evaluation of Secang Stem Extract \(*Caesalpinia Sappan* L \) in Decorative Cosmetics](#)

Ety Herawati and Lilies Yulastri

[Open abstract](#), Formulation and Evaluation of Secang Stem Extract (*Caesalpinia Sappan* L) in Decorative Cosmetics [View article](#), Formulation and Evaluation of Secang Stem Extract (*Caesalpinia Sappan* L) in Decorative Cosmetics [PDF](#), Formulation and Evaluation of Secang Stem Extract (*Caesalpinia Sappan* L) in Decorative Cosmetics

012058

THE FOLLOWING ARTICLE IS OPEN ACCESS

[The Effect of the Composition of Alginate Flour on the Making of Batik Simbut](#)

Roudlotus Sholikhah, Siti Nurrohmah and Atika

[Open abstract](#), The Effect of the Composition of Alginate Flour on the Making of Batik Simbut [View article](#), The Effect of the Composition of Alginate Flour on the Making of Batik Simbut [PDF](#), The Effect of the Composition of Alginate Flour on the Making of Batik Simbut

012059

THE FOLLOWING ARTICLE IS OPEN ACCESS

[A Great Food for Children Base on *Rinuak* Shredded Fish \(*Psilopsis* Sp\)](#)

Yuliana, Linda Rosalina, Kurnia Illahi Manvi and Heru Pramudia

[Open abstract](#), A Great Food for Children Base on Rinuak Shredded Fish (Psilopsis Sp) [View article](#), A Great Food for Children Base on Rinuak Shredded Fish (Psilopsis Sp) [PDF](#), A Great Food for Children Base on Rinuak Shredded Fish (Psilopsis Sp)

PAPER • OPEN ACCESS

Physical and Chemical Quality of Bento Rayap Grass Haylage (*Leersia Hexandra*) Supplemented with Water Mimmosa (*Neptunia Oleracea*)

To cite this article: Riswandi *et al* 2021 *IOP Conf. Ser.: Earth Environ. Sci.* **810** 012008

View the [article online](#) for updates and enhancements.

You may also like

- [Study on the observation of Eu²⁺ and Eu³⁺ valence states in low silica calcium aluminosilicate glasses](#)
J A Sampaio, M C Filadelpho, A A Andrade et al.
- [Modular quantum cosmology](#)
Orfeu Bertolami and Ricardo Schiappa
- [Observation Constraints on the Simplified GCG Model](#)
Dong Su-Mei and Wu Pu-Xun



ECS The Electrochemical Society
Advancing solid state & electrochemical science & technology

241st ECS Meeting

Vancouver, BC, Canada. May 29 – June 2, 2022

ECS Plenary Lecture featuring
Prof. Jeff Dahn,
Dalhousie University

Register now!

Physical and Chemical Quality of Bento Rayap Grass Haylage (*Leersia Hexandra*) Supplemented with Water Mimmosa (*Neptunia Oleracea*)

Riswandi^{1*}, A I M Ali¹, Muhakka¹, A Imsya¹, A Abrar¹, E Sahara¹, B Hamzah², A Supriadi³, P Alpian¹

¹Department of Animal Science, Faculty of Agriculture, University of Sriwijaya

²Department of Agricultural Product Technology, Faculty of Agriculture, University of Sriwijaya

³Department of Fishery Product Technology, Faculty of Agriculture, Universitas of Sriwijaya

*riswandi_dya@yahoo.com

Abstract. The effect of water mimmosa addition on the physical and chemical quality of Bento Rayap grass haylage was evaluated. The haylage material was fermented for 21 days and the physical quality of the resulting product was analyzed through organoleptic test and proximate analysis. Research variables consist of several treatments i.e. A0 (100% Bento Rayap grass), A1 (90% Bento Rayap grass + 10% water mimmosa), A2 (80% Bento Rayap grass + 20% water mimmosa) and A3 (70% Bento Rayap grass + 30% water mimmosa). The parameters observed were (1) physical quality includes color, aroma, texture, fungus and temperature while (2) chemical quality includes dry matter (DM) content, organic matter (OM), crude protein (CP), crude fiber (CF), and acidity (pH). The analysis result shows the addition of water mimmosa improved the physical quality of haylage from Bento grass. Water mimmosa addition also affects the chemical quality of Bento grass haylage confirmed by variance analysis result which gave statistically significant difference ($P < 0.05$). The property affected were DM, OM, CP, CF and pH. It was concluded that the addition of water mimmosa improved the physical and chemical quality of Bento Rayap grass haylage.

1. Introduction

The fluctuating availability of forage in the tropics is determined by the availability of land and season, in the rainy season the production is abundant and in the long dry season the production decreases so that there is a scarcity of forage sources [1]. In general, livestock business in Indonesia is still traditional and still relies on natural food sources so that it will affect the appearance of productivity that is not as expected. To overcome these problems, it is necessary to provide alternative forage sources from swamps.

Swamp forage is one of the feed ingredients that can be used to overcome the shortage of forage for livestock due to limited land availability. In addition, swamp forage can also provide a variety of feed ingredients for ruminants [2]. In connection with these conditions, technology is needed that can provide sustainable and quality animal feed. Several ways of forage preservation to provide forage throughout the year, such as making haylage (process of preserving forages using anaerobic fermentation technology). Anaerobic fermentation technology is more suitable to be applied in the provision of feed, this condition is because haylage has a low moisture content of 40-50% which



makes the haylage technology more durable. By making haylage, it can improve digestibility and feed efficiency and have good nutritional content [3].

Swamp grass forage that can be used as a feed ingredient for haylage is Bento Rayap grass (*Leersia hexandra*) and aquatic water mimmosa (*Neptunia oleracea* Lour). Bento Rayap grass (*Leersia hexandra*) is a source of fiber, has lower nutrients compared to swamp legumes [3]. Bento Rayap grass (*Leersia hexandra*) contains 11.98% crude protein (CP), 1.62% crude fat (EE), 27.39% crude fiber (CF), 82.15% NDF (Neutral Detergent Fiber), ADF (Acid Detergent Fiber) 51.60 % [4]. Water mimmosa legume is a type of swamp legume that has high nutritional content. The nutritional composition of water mimmosa (*Neptunia oleracea* Lour) consists of 28.02% CP, 2.02% EE and 17.25% CF [4]. Optimizing the use of swamp forage in the form of haylage can be done by combining swamp grass with protein sources such as legume, because the protein content of legume is quite high, so it is expected to be able to meet the nutritional needs of feed when given to livestock. Ridwan et al. [5] confirmed that Leguminosae addition on grass silage enhance physical quality and nutrient content. The aimed of this research are to evaluate the addition of water mimmosa (*Neptunia oleracea*) and its effect on the physical and chemical quality of Bento Rayap grass haylage and to obtain the effective formula on increasing haylage quality.

2. Methodology

2.1. Location and Duration

Place and Time his study was conducted in the Laboratory of Animal Nutrition and Feed, Faculty of Agriculture, University of Sriwijaya for 3 months.

2.2. Ensilage

The Bento Rayap grass (*Leersia hexandra*) and water mimmosa (*Neptunia oleracea*) was withered for 24 hours prior process to reduce its water content ($\pm 40\%$). Size reduction of samples were conducted by using chopper to obtain 3 cm in length. Molasses was acquired from sugar cane processing industry at Cinta Manis and added into silage as additive. The Bento Rayap grass combined with water mimmosa legume and molasses 5% was prepared according to research variables (w/w) design. Completely randomized design was used in silage making and comprise of four treatments and four replicates i.e. A0 (100% Bento Rayap grass), A1 (90% Bento Rayap grass + 10% water mimmosa), A2 (80% Bento Rayap grass + 20% water mimmosa) and A3 (70% Bento Rayap grass + 30% water mimmosa). All of materials prepared was placed in plastic tank (silo) of 5 kg capacity, compressed and closed to create anaerobic condition. The incubation process was conducted in room temperature (± 28 °C) for 21 days. During certain period of time, analysis on physical and chemical quality of product was carried out.

2.3. Haylage Sample Preparation

The haylage result after incubated for 21 days was split into two parts, the first part was dried and grinded to process further for proximate analysis while the second part was used for physical quality analysis (organoleptic) and pH determination.

2.4. Physical Quality Observation of Haylage Product

The physical quality of haylage was assessed through organoleptic test comprise of 15 panelists i.e. students and lecturer with food technology competence. The result of aroma, color and texture observation as well as fungus number on haylage were subjected to descriptive analysis [6].

2.5. Chemical Quality Determination of Haylage Product

The determination of pH were carried out using liquid extracted from haylage. The extraction procedure is as follows, samples (20 g) were placed in Erlenmeyer flask, 300 mL distilled water were added and then stirred for 30 minutes. The suspension was left for 24 hours [7]. Standard method was used to determine dry matter, organic matter, crude protein and crude fiber [7].

2.6. Data Analysis

Data obtained in this research was subjected to analysis of variance, the difference found out will be treated further using Duncan test using software SPSS 13.0 [8].

2.7. Observed Parameters

The observed parameters in this work were physical quality i.e. color, aroma, texture, fungus and temperature whereas chemical quality was assessed through dry matter (DM), organic matter (OM), crude protein (CP), crude fiber (CF) and pH.

3. Result and Discussion

3.1. The Physical Quality

Physical observation carried out by 15 panelists, the haylage product displays good physical quality indicates by its aroma (sour smell or typical fermentation aroma no bad odor), solid texture (no clot, no mucus or crumb), yellowish green, a view fungi, harvest temperature below the environment (Table 1).

Table 1. The effect of water mimmosa addition on the physical quality of Bento Rayap grass haylage

Treatment	Aroma	Color	Texture	Fungi (%)	Temperatur (° C)
A0	Fermentation aroma	Yellowish green	Solid	3.5	27
A1	Fermentation aroma	Yellowish green	Solid	3.7	27
A2	Fermentation aroma	Yellowish green	Solid	5.3	28
A3	Fermentation aroma	Yellowish green	Solid	5.6	28

Note: A0 = 100% bento rayap grass, A1 = 90% bento rayap grass + 10% water mimmosa, A2 = 80% bento rayap grass + 20% water mimmosa, A3 = 70% bento rayap grass + 30% water mimmosa.

a) Aroma

The results showed that the Bento Rayap grass haylage termites obtained all displayed a fermented aroma. Fermentation aroma is affected by lactic acid bacteria activity and other heterofermentative bacteria. According to McDonald et al. [9], heterofermentative silage produce not only lactic acid but also acetic, propionic, butyric and alcohol. Sour smell shows that the silage is in good quality. Saun and Heinrichs [10] reported that good silage smells like milk due to lactic contained does not has strong aroma. Another factor affects the silage aroma is the addition of 5% molasses. Molasses contains water soluble carbohydrate such as sucrose. This disaccharide compound is easy to consume by microbe during ensilage process to produce high lactic acid and creates strong fermentation aroma.

b) Color

Throughout ensilage process, forage experience discoloration which depends on raw material type and temperature effect. High temperature at the ensilage process convert haylage color into brown as the Maillard reaction occurred [11]. The Bento Rayap grass haylage obtained, all display yellowish green or almost as green as should be. Macaulay [12] recommended that good quality of silage has color of light green to yellow or brownish green depends on silage raw material. McDonald et al. [9] described that good visual display, sour smell, color as similar as its raw material and low pH (3.9-4.1) expresses well fermented silage.

c) Texture

Haylage texture observation on each treatment after 21 days ensilage show all result have solid texture, compact and no mucus. No damage observed such as dry (moldy) and broken texture means all treatment produced good quality of haylage. The Bento Rayap grass withering prior the treatment made it has suitable water content approximately 40-50% for fermentation process. Macaulay [12] reported that water content of raw material influences the initial step of ensilage. High water content (>80%) will result in slimy texture, soft and moldy whereas low water content (<30%) produce dry

texture and moldy. Despal et al. [13] explained silage added with accelerator (molasses) displays intact texture, soft and no mucus, which designate minor damage of silage result during ensilage process.

d) Fungi

The presence of fungi on haylage process is a good indication of a successful process. The analysis showed that a few fungi were found in the treatment, namely 3.5-5.6%. The water mimmosa legume addition in general able to maintain the good quality of haylage result. This result confirmed by Bento Rayap grass haylage pH at 3.23 – 4.13 (Table 2). At this pH, the growth of decay bacteria can be suppressed. Water mimmosa legume addition into Bento Rayap grass haylage made no drastic change in pH and still maintain the ideal condition for haylage process. The legume added in the raw material also enhance silage material density due to its crude fiber lower than the Bento Rayap grass, so it affected the solidification level. High solidification level prevent air trapped in a silo and repressed undesirable fungus and microbial spores to grow. According to Donald et al. [14], the presence of fungi is related to the existence of air trapped in silo both at initial stage of ensilage and the silo leaks during storage.

e) Temperature

Average value of silage temperature during harvest process is at 27-28 °C. This is a good condition for silage production because it below environment temperature (± 29 °C). Ridwan et al.[5] reported the temperature range of harvest silage result is between 26 to 28 °C and consider good as long as below open-air temperature. Temperature above the environment by 5-10 °C presumed silage contaminated by microorganism such as mold and fungus. As the ensilage process move forward, it triggers the air-tight within material and stimulates the lactic acid bacteria to produce lactic acid and prevent prolonged heat hence the temperature stabilized.

3.2. The Chemical Quality

The chemical quality observed on Bento Rayap grass haylage with the addition of water mimmosa is shown on Table 2.

Table 2. The effect of water mimmosa addition on the chemical quality of Bento Rayap grass haylage

Parameters	Treatments			
	A0	A1	A2	A3
Dry matter (%)	30.20 ^a ± 0.65	31.02 ^b ± 0.64	31.23 ^b ± 0.75	32.05 ^c ± 0.53
Organic matter (%)	28.09 ^a ± 0.71	28.73 ^b ± 0.62	29.19 ^b ± 0.89	29.85 ^c ± 0.39
Crude protein (%)	13.59 ^a ± 0.87	17.85 ^b ± 1.08	18.24 ^c ± 1.48	20.80 ^d ± 0.76
Crude fiber (%)	24.21 ^c ± 0.45	22.9 ^b ± 0.67	22.66 ^{ab} ± 0.64	22.33 ^a ± 0.49
pH	3,23 ^a ± 0.09	3,72 ^b ± 0.29	3.82 ^b ± 0.29	4.13 ^c ± 0.37

Note: The numbers in the same row followed by different letters indicate significantly different at test level 5%. A0 = 100% bento rayap grass, A1 = 90% bento rayap grass + 10% water mimmosa, A2 = 80% bento rayap grass + 20% water mimmosa, A3 = 70% bento rayap grass + 30% water mimmosa.

a) Dry Matter and Organic Matter Content

The addition of water mimmosa at 10% (A1), 20% (A2) and 30% (A3) significantly difference ($P < 0.05$) on DM and OM, while treatments between A1 and A2 shows no statistical significance different ($P > 0.05$). This result means that water mimmosa addition increase the DM and OM content of haylage due to the higher DM and OM content of water mimmosa compare to Bento Rayap grass. Ridwan et al. [5] reported that increase amount of *C. Colothyrsus* added in *P. Purpureum* grass would enhance DM and OM content and it is also reduced NDF and ADF of the silage product. McDonald et al. [9] suggested that decrease in DM and OM content during ensilage is affected by nutrient contained in raw material and microorganism involved.

b) Crude Protein Content

The addition of water mimmosa at 10% (A1), 20% (A2) and 30% (A3) affects CP content significantly ($P < 0.05$). Further test result indicates A1, A2 and A3 treatments statistically significance different with A0. The treatments also statistically significance among A1, A2 and A3. High nutrient content within water mimmosa legume improve microbial growth involved in haylage process hence increase its population. The microbe itself is single-cell protein resource thus it caused the increase of crude protein within silage as reported by Lima et al. [15].

c) Crude Fiber Content

The addition of water mimmosa at 10% (A1), 20% (A2) and 30% (A3) significantly difference ($P < 0.05$) on CF. Legume added in haylage material caused decrease in CF content compare with result without legume. High nutrient of water mimmosa legume as discussed previously enhance the ensilage process to produce haylage, this event was accompanied by increase of cellulose and hemicellulose degradation. Schroeder [16] suggested that heterofermentative bacteria convert simple glucose to organic acids (acetic, lactic, propionic and butyric) which cause the decrease in CF content.

Crude fiber content obtained by A3 treatment provide best result among other treatments. In general, Water mimmosa legume added to haylage forage decreases the CF content of the materials.

d) pH Value

Statistic calculation result confirmed the addition of water mimmosa significantly difference ($P < 0.05$) on the Bento rayap grass haylage pH (Table 2). Low pH shows better quality of silage [16]. Measurement result shows the average value of pH is 3.23 – 4.13. Low pH value obtained on haylage without the addition of water mimmosa legume (3.23) whereas high pH (4.13) obtained at legume addition by 30% (A3). The increase level of legume addition also tends to increase pH. water mimmosa legume has buffering capacity higher than grass. This buffering capacity climbs up during ensilage process due to the production of organic acids such as nitric and sulfuric acid which drove the pH to higher value [16]. Bento Rayap grass haylage in general has pH which resemble good quality of fermentation. According to Macaulay [12], silage quality can be classify based on pH by 4 types i.e. excellent (3.2 – 4.2), good (4.2 – 4.5), medium (4.5 – 4.8) and poor ($\text{pH} > 4.8$). Water mimmosa legume addition into Bento Rayap grass haylage made no drastic change in pH and still maintain the ideal condition for haylage process. The buffering capacity of legume can be suppressed by the presence of water-soluble carbohydrate from Bento Rayap grass. Lactic acid bacteria can still growth to produce lactic acid and lower the product pH [17].

4. Conclusion

It was concluded that the addition of water mimmosa improved the physical and chemical quality of Bento Rayap grass haylage.

Acknowledgment

The authors wish to acknowledge and thank you to the Sriwijaya University which provide research funding through Unggulan Kompetitif scheme year 2019 contract number: 0015/UN9/SK.LP2M.PT/2019.

References

- [1] Santoso B and Hariadi B T J 2009 *JITV* 13 3
- [2] Riswandi 2014 *Jurnal Pet. Sriwijaya* 3 2
- [3] Jaelani A T, Rostini, Zakir M I dan Jonathan 2014 *Jurnal Sain Pet.* 12 2
- [4] Ali A I M, Sandi S, Muhakka, Riswandi, Budianta D 2013 APCBEE Procedia ICAAA July 27-28 Moscow Russia
- [5] Ridwan R, I Rusmana, Y Widyastuti, K G Wiryawan, B Prasetya, M Sakamoto and M Ohkuma 2015 *Asian-Australas J. Anim. Sci.* 28
- [6] Rostini T 2014 *Int. J. of Biosci.* 5 12

- [7] AOAC 2003 *Official Methods of Analysis of the Association of Official Analytical Chemists*. 17th Ed. Association of Official Analytical Chemist, Arlington
- [8] Steel RGD dan JH Torrie 1993 *Prinsip dan Prosedur Statistika*. Gramedia Pustaka Utama. Jakarta
- [9] McDonald P, Edwards R A, Greenhalgh J F D, Morgan C A, Sinclair L A, Wilkinson R G 2010 *Animal Nutrition*. Seventh Ed. New York. C A Morgan, J F D Greenhalgh, L A Sinclair and R G Wilkinson, Inc.
- [10] Saun RJV and Heinrichs A J 2008 *Proc. of the Mid-Atlantic Conf.*; Pennsylvania, 26 May. Penn State's Collage
- [11] Gonzalez J, J Faria-M'armol, C A Rodriguez & A Mart'inez 2007 *Anim. Feed Sci. Technol.* 136
- [12] Macaulay A. *Evaluating Silage Quality* (2004). ca/departement/deptdocs.nsf/all/for4909.
- [13] Despal I G, Permana, S N Safarina dan A J Tatra. 2011 *J. Media Pet.* 43
- [14] Donald S, Fenlon DR, Seddon B 1993 *Proc. 10 th Silage Res. Conf.* Dublin City University, Dublin, Ireland
- [15] Lima R, R F Díaz, A Castro, S Hoedtke and V Fievez 2011 *Anim. Feed Sci. Technol.* 164
- [16] Schroeder J W 2013 *Silage Fermentation and Preservation*. Extension Dairy Speciaslist. AS-1254 <http://www.ext.nodak.edu/>
- [17] Saricicek B Z and Kilic U 2011 *Asian J. Anim. Vet. Advances* 6