

# Optimization of *Spirulina platensis* Culture Media as an Effort for Utilization of *Pangasius* Farming Waste Water

*By Dade Jubaedah*

## Optimization of *Spirulina platensis* Culture Media as an Effort for Utilization of Pangasius Farming Waste Water

Marini Wijayanti<sup>1\*</sup>, Dade Jubaedah<sup>1</sup>, Nuni Gofar<sup>2</sup>, Devi Anjastari<sup>3</sup>

<sup>1</sup>Aquaculture Program, Department of Fisheries, Faculty of Agriculture, Universitas Sriwijaya

<sup>2</sup>Department of Soil Science, Faculty of Agriculture, Universitas Sriwijaya

<sup>3</sup>Student of Aquaculture, Department of Fisheries, Faculty of Agriculture, Universitas Sriwijaya  
Jl. Raya Palembang-Prabumulih KM.32 Indralaya, Ogan Ilir, South Sumatera, Indonesia

\*Corresponding Author: e-mail address : mariniwijayanti@fp.unsri.ac.id

### Article history

| Received         | Received from revised | Accepted        | Available online |
|------------------|-----------------------|-----------------|------------------|
| 28 November 2018 | 07 January 2019       | 10 January 2019 | 11 January 2019  |

**Abstract:** *Pangasius* is a fast-growing fish species that has great potential for production and export growth in Indonesia. Their farming produces a lot of organic material and ammonia which potentially make pollution in freshwater body. The wastewater can be used for high value microalgal cultivation media. The microalgae are used in various fields, one of those is *Spirulina platensis*, a spiral blue green algae. This aims of this study was to determine the best composition of the technical fertilizer in the pond waste of *Pangasius* farming pond to obtain maximum density and know the specific growth rate of *Spirulina platensis*. The research method used Completely Randomized Design (CRD) with 6 treatments (3 replication), P<sub>0</sub> using 100% technical fertilizer without using *Pangasius* waste water while P<sub>1</sub>-P<sub>5</sub> use 0% (P<sub>1</sub>), 25% (P<sub>2</sub>), 50% (P<sub>3</sub>), 75% (P<sub>4</sub>) and 100% (P<sub>5</sub>) technical fertilizer using *Pangasius* farming waste water. The most efficient treatment obtained 23.90 gL<sup>-1</sup> maximum density, 6.22%.d<sup>-1</sup> specific growth rate and 87.77% ammonia removal.

**Keywords:** *Spirulina*, *Pangasius* farming, waste water, culture media

**Abstrak (Indonesian):** *Pangasius* adalah spesies ikan cepat tumbuh yang memiliki potensi besar untuk produksi dan pertumbuhan ekspor di Indonesia. Pemeliharaannya menghasilkan banyak bahan organik dan amonia yang berpotensi membuat polusi dalam tubuh air tawar. Air limbah dapat digunakan untuk media budidaya mikroalga bernilai tinggi. Mikroalga digunakan di berbagai bidang, salah satunya adalah *Spirulina platensis*, ganggang hijau biru spiral. Tujuan penelitian ini adalah untuk menentukan komposisi pupuk teknis terbaik dalam kolam tambak budidaya *Pangasius* untuk mendapatkan kepadatan maksimum dan mengetahui tingkat pertumbuhan spesifik *Spirulina platensis*. Metode penelitian menggunakan Rancangan Acak Lengkap (RAL) dengan 6 perlakuan (3 ulangan), P<sub>0</sub> menggunakan pupuk teknis 100% tanpa menggunakan air limbah *Pangasius* sedangkan P<sub>1</sub>-P<sub>5</sub> menggunakan 0% (P<sub>1</sub>), 25% (P<sub>2</sub>), 50% (P<sub>3</sub>), 75% (P<sub>4</sub>) dan 100% (P<sub>5</sub>) pupuk teknis menggunakan air limbah pertanian *Pangasius*. Perlakuan yang paling efisien diperoleh 23,90 gL<sup>-1</sup> kepadatan maksimum, laju pertumbuhan spesifik 6,22% .h<sup>-1</sup> dan penghapusan amonia mencapai 87,77%.

**Katakunci:** *Spirulina*, pembesaran patin pangasius, air limbah, media kultur

### 1. Introduction

*Pangasius* is a fast-growing fish species that has great potential for production and export growth in Indonesia. Henriksson *et al.* (2017) [1] showed that 1000 kg *Pangasius* production in 195 m<sup>2</sup> pond system will produce 0.43 kg Ammonia, 0.79 kg Nitrous oxide, 34.2 kg Ammonium, 14.7 kg Nitrate, 1.93 kg Nitrogen, and 1.54 kg Phosphorus as environmental output. Their farming produces a lot of organic material and ammonia which potentially make pollution around freshwater. The wastewater can be spend for high value microalgal cultivation media before releasing to environment. Cyanobacterias have potential for growth using solar energy and large quantities of nutrients effluents rich in inorganic

pollutants, such as nitrogen and phosphate. They are an alternative for decreasing cost [16] large scale production. This production became treatment of the wastewaters occur through removal of the pollutants [2].

The high value microalgae like a cyanobacteria, *S.platensis*. It has several characteristics and nutrients that are suitable as functional foods, feed, and nutraceutical [3]. *Spirulina* have known for its high protein content and therapeutic properties which has been studied as a potential source of bioactive peptides for their ability to offer specific health benefits, such as antimicrobial, antiallergic, antihypertensive, antitumor, and immunomodulatory properties [4]. *Spirulina platensis* was successfully

cultured using organic nutrients from liquid waste, such as molasses [5], tofu and latex liquid waste [6], Palm Oil Mill Effluent/ POME [7] and vinasse from anaerobic processing [8].

Therefore, this study was carried out to examine the effect of *Pangasius* catfish culture waste which was added to fertilizer for *Spirulina* media. The growth of microalgae can remediate fish farm wastewater because aquaculture wastewater contains good organic material and ammonia to increase the density of *S. platensis*. Production of *Spirulina platensis* is good for removing ammonia and nitrate found in fish pond waste water [9].

*Spirulina platensis* can remove the nutrients dissolved on the fish Nile tilapia culture effluent with density of *S. platensis* resulted in the production of 0.22 g L<sup>-1</sup> of dry biomass and maximum productivity of 0.03 g L<sup>-1</sup> day<sup>-1</sup>. The *S. platensis* were capable of reducing the nitrite, nitrate and phosphate levels by 100; 98.7 and 94.8%, respectively [10]. The process of *Pangasius* catfish cultivation produces waste that can pollute the environment. This study is expected to reduce the impact of environmental pollution by utilizing *Pangasius* catfish pond wastewater for the growth media of *S. platensis* and to determine the best composition of technical fertilizers in the utilization.

## 2. Material and Methods

### 2.1. Acquiring the *Spirulina platensis*

The strain with 1000 mL of *S. platensis* was cultured from a aquaculture laboratory at Aquaculture Study Program, Fisheries Department, Faculty of Agriculture, Universitas Sriwijaya. Its volume was increased on scaling up to a maximum of 5 L by the addition of the modified technical fertilizer medium [6] with salinity of 1 ‰, remaining under constant aeration and artificial lighting of 1000-1200 lux, termed stock culture.

### 2.2. Optical density and biomass density

The monitoring of the growth of *S. platensis* cultures was performed daily by measuring the absorbance (optical density - OD) of a sample with 0,005 L of water from the culture in a spectrophotometer Spectronic G-20, with a wavelength of 560 nm (Abs560). The biomass density (g.L<sup>-1</sup>) was approached with :

$$\text{Biomassa density} = 5.9594 \cdot (\text{Abs560}) + 1.1837$$

### 2.3. Determining the yield and productivity of the *S. platensis* culture

To determine the yield and productivity of the *S. platensis* cultures it was necessary to separate the biomass from the culture medium. In order to do this, five liters of the culture, obtained from phases of the reduced growth, were filtrated with calico cloth. Subsequently, the biomass was dried in a laboratory oven for 48 h with air renewal at 60 °C.

### 2.4. Experimental design

The experimental design followed six treatments in triplicates (Table1). *Pangasius* pond water waste that has a size of 3 x 4 x 1 m<sup>3</sup> with a stocking density of 7 ind. m<sup>-3</sup> size 40-45 cm are maintained for 6 months by feeding floating pellets size 5-6 mm (36-38% protein, 5-6% fat, 4% crude fiber, 10% ash and 11% moisture content) given twice a day (morning and evening) containing a total nitrogen content of 1.9 mg.L<sup>-1</sup>, total organic carbon 11.4 mg.L<sup>-1</sup>, phosphorus 2.6 mg.L<sup>-1</sup> and ammonia 0.81 mg.L<sup>-1</sup>. Meanwhile the nutrient content of the technical fertilizer for *S. platensis* cultivation contained total nitrogen content of 31.94 mg.L<sup>-1</sup>, total organic carbon 24.05 mg.L<sup>-1</sup> and phosphorus 0.07 mg.L<sup>-1</sup>.

Table 1. Chemical composition of the modified technical fertilizer medium

| Material<br>of technical fertilizer                               | P <sub>0</sub><br>(100% fertilizer) | Treatment   |                       |                       |                       |                        |
|---|-------------------------------------|---|-----------------------|-----------------------|-----------------------|------------------------|
|   |                                     | % fertilizer in waster water <i>Pangasius</i> farming |                       |                       |                       |                        |
|   |                                     | P <sub>1</sub><br>0%                                  | P <sub>2</sub><br>25% | P <sub>3</sub><br>50% | P <sub>4</sub><br>75% | P <sub>5</sub><br>100% |
| MgSO <sub>4</sub> (g)   | 0.6                                 | -   | 0.15                  | 0.3                   | 0.45                  | 0.6                    |
| CaCl <sub>2</sub> (g)   | 0.12                                | -   | 0.03                  | 0.06                  | 0.09                  | 0.12                   |
| C <sub>10</sub> H <sub>16</sub> N <sub>2</sub> O <sub>8</sub> (g) | 0.24                                | -   | 0.06                  | 0.12                  | 0.18                  | 0.24                   |
| Ca(H <sub>2</sub> PO <sub>4</sub> ) (g)                           | 1.5                                 | -   | 0.375                 | 0.75                  | 1.125                 | 1.5                    |
| (NH <sub>2</sub> ) <sub>2</sub> CO (g)                            | 0.9                                 | -   | 0.225                 | 0.45                  | 0.675                 | 0.9                    |
| NH <sub>4</sub> SO <sub>4</sub> (g)                               | 3.96                                | -   | 0.99                  | 1.98                  | 2.97                  | 3.96                   |
| NaHCO <sub>3</sub> (g)  | 25.5                                | -   | 6.375                 | 12.75                 | 19.125                | 25.5                   |
| Larutan AB <sub>mix</sub> * (ml)                                  | 3                                   | -   | 0.75                  | 1.5                   | 6.75                  | 3                      |

Note : \* = modified technical fertilizer medium in A<sub>5</sub>.

The growth of cyanobacteria and the nutrients levels in the waters were monitored every day. The calculation of the specific growth rate *Spirulina platensis* can be performed using a modified formula [11] :

$$\mu = \frac{\ln N_{t_i} - \ln N_0}{t_i} \times 100\%$$

$\mu$  = daily growth rate (%. days<sup>-1</sup>)

$t_i$  = time (days) from  $N_0$  to  $N_t$

$N_0$  = Initial density (g.L<sup>-1</sup>)

$N_t$  = density at the time  $t_i$  (g.L<sup>-1</sup>)

## 2.5. Water Quality Parameters

Monitoring of water quality was done as supporting parameters for the research carried out. Salinity (refractometer, MW) and pH (pH meter, MW) were measured daily for experimental period. Value of pH maintained between 8 to 8.5 and salinity  $20 \pm 1$  ppt during the study. The ammonia measurements (American Public Health Association (APHA), 2005) carried out at the beginning and end of the study. The percentages of Ammonia removal presented in the effluents treated by *Spirulina platensis* were calculated using the following equation:

$$\text{Ammonia removal} = \frac{([\text{NH}_3]_0 - [\text{NH}_3]_t) / [\text{NH}_3]_0 \times 100\%}{}$$

## 2.6. Statistical Analysis

The results were submitted to simple analysis of variance tests (ANOVA) ( $p < 0.05$ ) and in the case of significant differences, the means were compared by the Least Significant Differences test ( $p < 0.05$ ).

## 3. Results and Discussion

### 3.1. Biomass density and specific growth rate

The highest peak density of *S. platensis* in each treatment was different time. Maximum density in treatment P0 is 16.31 gL<sup>-1</sup> on the 10<sup>th</sup> day of culture, in treatment P1 reached 23.90 gL<sup>-1</sup> on the 10<sup>th</sup> day of culture, in treatment P2 reached 25.09 gL<sup>-1</sup> on day 10 the culture period, at treatment P3 reached 19.68 gL<sup>-1</sup> on 10<sup>th</sup> 9<sup>th</sup> day of culture, P4 treatment reached 18.73 gL<sup>-1</sup> on the 10<sup>th</sup> day of culture and P5 treatment reached 20.78 gL<sup>-1</sup> on the 10<sup>th</sup> day culture. The maximum density of *S. platensis* in each treatment can be seen in Figure 1.

The highest density of *S. platensis* was found in P2 treatment which had the highest biomass density with an average of 25.09 g.L<sup>-1</sup>, while the lowest density was found in treatment P0 which had biomass density 16.31 g.L<sup>-1</sup>. The high and low density of *S. platensis* is influenced by nutrients found in *S. platensis* cultivation media. The difference of nutrient content in culture media, influences cell quality and density. Technically, microalgae absorbs the content of organic compounds and nutrients in waste, and produces oxygen which can reduce the levels of COD

and BOD in waste through the help of organic matter decomposing bacteria [12].




Figure 1. Growth of *S. platensis* in the treatment cultivation medium

The results of the variance analysis showed that the provision of technical fertilizer in *Pangasius* culture wastewater had a significant effect on the maximum density and specific growth rate of *S. platensis*. The maximum density and specific growth rate of *S. platensis* for each treatment are shown in Table 2.

Table 2. Maximum density of *S. platensis* (g.L<sup>-1</sup>) and their specific growth (%.d<sup>-1</sup>)

| Repetition | Mean ± SD<br>$\alpha = 2,06$ | Mean ± SD<br>$\alpha = 0,98$ |
|------------|------------------------------|------------------------------|
| P0         | 16,47 ± 1,05 <sup>a</sup>    | 3,54 ± 0,67 <sup>a</sup>     |
| P1         | 23,90 ± 1,09 <sup>c</sup>    | 6,22 ± 0,51 <sup>c</sup>     |
| P2         | 25,09 ± 1,50 <sup>c</sup>    | 6,85 ± 0,65 <sup>c</sup>     |
| P3         | 20,00 ± 0,16 <sup>b</sup>    | 5,01 ± 0,33 <sup>b</sup>     |
| P4         | 18,73 ± 1,73 <sup>b</sup>    | 4,25 ± 0,66 <sup>ab</sup>    |
| P5         | 20,78 ± 0,69 <sup>b</sup>    | 5,05 ± 0,47 <sup>b</sup>     |

Note: The numbers followed by different small superscript letters show significant differences ( $p < 0.05$ ).

Based on the LSD test presented in Table 2, it shows that the treatment of P2 was significantly different than the treatment P0, P3, P4 and P5 but not significantly different from the treatment P1. Treatment P2 obtained the best maximum density value with a value of 25.09%. However, a more efficient treatment was P1, because treatment P1 used waste media without additional technical fertilizer. While the lowest density was treatment P0 with a value of 16.47%.

Microalgae can grow well if the nutrients and environment are suitable, if not optimal then microalgae growth will be low. The use of technical fertilizers with high concentrations can reduce water quality and decrease the quality of *S. platensis* growth. It will also cause discoloration (turbidity) in the cultivation media so that the light penetration decreases which will interfere photosynthesis and production of *S. platensis* [13]. The population density

at P0, P3, P4 and P5 is lower than P1 and P2 which shows a relatively high growth pattern.

P2 treatment obtained the best specific growth rate with growth of 6.85% day<sup>-1</sup>, while treatment P0 obtains in a relatively low growth rate of 3.54% day<sup>-1</sup>. P2 treatment was significantly different than the treatment P0, P3, P4 and P5 but was not significantly different from treatment P1, because it was not significantly different between P1 and P2 (0% and 25%), the most efficient treatment was P1 with waste media without fertilizer for *S. platensis* culture. The difference in specific growth rate in each treatment is caused by the ability of cells to absorb nutrients contained in culture media, not all materials can be directly absorbed and used by cells [14]. In addition, differences in specific growth rates are also influenced by nutritional factors contained in culture media.

The specific growth rate in P2 treatment due to the administration of 25% fertilizer mixed with *Pangasius* cultivation wastewater was in accordance with the nutritional needs of *S. platensis*. Nitrogen is a nutrient that is needed in the formation of chlorophyll, where chlorophyll is needed for photosynthesis and cell growth [15]. When nitrogen concentrations in optimal culture media, cell metabolic activity also goes well, including chlorophyll synthesis, because high chlorophyll content will cause photosynthesis to work well and *S. platensis* growth will be more optimal [16].

The regression relationship between population density (y) and the percentage of technical fertilizer in *Pangasius* farming pond waste water was shown in Figure 2. The polynomial regression analysis results from the average daily population density indicated by the equation  $y = 51,237x^3 - 70,549x^2 + 16,191x + 24,048$  with coefficient of determination  $R^2 = 0.9443$ .




Figure 2. Regression of *S. platensis* biomass density to percentage of technical fertilizer

Based on the regression equation in Figure 2, it was found that the optimum technical fertilizer concentration was in the range of 13.44% which resulted in the highest maximum density of *S. platensis* of 25.07 g.L<sup>-1</sup>. The level of effectiveness of *S. platensis* growth is indicated by the increase in biomass

production. Addition of 13.44% of technical fertilizer into *Pangasius* farming wastewater is thought to be most suitable for the nutritional needs of *S. platensis*.

### 3.2. Ammonia removal

Ammonia is a toxic substance in the waters that is harmful to the health of fish. Ammonia microalgae can be used as nutrients so as to reduce pollution of waters. The percentage of ammonia absorption during the study can be seen in Table 3. Based on the LSD test presented in Table 3, the treatment P2 shows the best ammonia absorption results compared to other treatments, but the treatment of P2 was not significantly different from the treatments P1 and P3, but it was significantly different in the treatments P0, P4 and P5.

Table 3. Ammonia removal in cultivation medium during *S. platensis* cultivation (%)

| Treatment | Ammonium Removal<br>$\alpha = 2,03$ |
|-----------|-------------------------------------|
| P0        | 81,95 ± 1,02 <sup>a</sup>           |
| P1        | 87,77 ± 1,01 <sup>c</sup>           |
| P2        | 88,07 ± 1,03 <sup>c</sup>           |
| P3        | 87,07 ± 1,45 <sup>bc</sup>          |
| P4        | 80,47 ± 1,00 <sup>a</sup>           |
| P5        | 85,57 ± 1,24 <sup>b</sup>           |

Note: The numbers followed by different small superscript letters show very significant differences ( $p < 0,05$ )

The ability of *S. platensis* to absorb ammonia in wastewater from *Pangasius* cultivation with the addition of different fertilizer doses has the highest value of 88.07%. During the 13 days of maintenance the ammonia level in the cultivation media diminished. Ammonia was used by microalgae as a nutrient for its growth, which was to help in protein synthesis. Algae requires ammonium for growth. Amanatin and Nurhidayati (2013) [16] showed that ammonium is produced through the dissociation process of ammonium hydroxide. Ammonium hydroxide is ammonia which dissolves in water. Algae must convert nitrate (the main form of nitrogen in water) to ammonium before they can use it. *S. platensis* utilizes 88.07% ammonia from *Pangasius* farming wastewater and from technical fertilizers, so that the density of *S. platensis* has increased. *S. platensis* is able to absorb ammonia from culture media optimally.

High ammonia can be dangerous for aquatic biota because it is toxic but is different from *S. platensis*. Yuan *et al.* (2011) [17] explained that cyanobacteria algae is able to tolerate high ammonia for its growth, but if excessive ammonia will inhibit the growth of *S. platensis*. Effendi (2003) [18] states that an increase in ammonia is influenced by pH. As the pH increases above 7 in the water, the ammonia percentage will increase as well. *S. platensis* was cultivated at alkaline medium at pH 8-8.5 so that ammonium was still more

dominant than ammonia. Control of pH in a culture medium is very important to maintain the balance of *S. platensis* cell growth.

#### 4. Conclusion

Utilization of *Pangasius* farming wastewater as the most efficient medium for *Spirulina platensis* growth was in treatment P1 with waste media without the addition of technical fertilizer which produces a maximum density of 23.90 gL<sup>-1</sup> with a growth rate of 6.22% day<sup>-1</sup> and the ammonia removal 87.77% was the highest.

#### Acknowledgement

We thank to Sriwijaya University for funding this project through the "Penelitian Unggulan Profesi Dosen" with the research contract of No. 0109.26/UN9/SB3.LP2M.PT/2018.

#### References

- [1] P.J.G. Henriksson, N. Tran, C.V. Mohan, C.Y. Chan, U-P. Rodriguez, S. Suri, L.D. Mateos, N.B.P. Utomo, S. Hall, M.J. Phillips, Indonesian aquaculture futures e Evaluating environmental and socioeconomic potentials and limitations, *Journal of Cleaner Production* 162, 1482-1490, 2017.
- [2] Markou, G. and Georgakakis, D. Cultivation of filamentous cyanobacteria (blue-green algae) in agro-industrial wastes and wastewaters: A review. *Applied Energy* 88, 3389–3401, 2011.
- [3] L.M. Andrade, C.J. Andrade, M. Dias, C.A.O. Nascimento, M.A. Mendes, *Chlorella* and *Spirulina* Microalgae as Sources of Functional Foods, Nutraceuticals, and Food Supplements; an Overview. *MOJ Food Process Technol.* 6(2), 2018.
- [4] C.A. Ovando, J.C. de Carvalho, G.V. de Melo Pereira, P. Jacques, V.T. Soccol & C.R. Soccol. Functional properties and health benefits of bioactive peptides derived from *Spirulina*: A review, *Food Reviews International*, DOI: 10.1080/87559129.2016.1210632, 2016.
- [5] M.R. Andrade dan J.A.V. Costa. Mixotrophic cultivation of microalga *Spirulina platensis* using molasses as organic substrate. *Elsevier B.V. Biochemical Engineering Laboratory, Department of Chemistry, Federal University Foundation of Rio Grande*, 2007.
- [6] M. Wijayanti, D. Juabedah, R. Puspitasari, 2010. *Spirulina* production in fertilizer medium combined by tofu and latex liquid wastes. *Proceeding International Seminar On Food And Agricultural Sciences-Isfas*, 16-17 February 2010, pp 44-50.
- [7] Suharyanto, Tri-Panji, S.Permatasari, & K. Syamsu. Production of *Spirulina platensis* in continous photobioreactor using palm oil mill effluent media. *Menara Perkebunan*, 82(1), 1-9, 2014.
- [8] Budiyono, I., Syaichurrozi, S. Sumardiono, dan S.B. Sasongko. Production of *Spirulina platensis* Biomass Using Digested Vinasse as Cultivation Medium. *Trends in Applied Sciences Research*. 9 (2), 93-102, 2014
- [9] S.C. Wuang, M.C. Khin, P.G.D. Chua, dan Y.D. Luo. Use of *Spirulina* biomass produced from treatment of aquaculture wastewater as agricultural fertilizers. *Algal Research*. 15, 59-64, 2016.
- [10] S.M.S. Nogueira, J.S. Junior, H.D.Maia, J.P.S. Saboya, and W.R.L. Farias. Use of *Spirulina platensis* in treatment of fish farming wastewater. *Revista Ciência Agronômica*, 49(4), 599-606, 2018.
- [11] A. Vonshak. *Spirulina: Growth, physiology and biochemistry*. In : A. Vonshak (Ed.) *platensis (Arthrospira) Physiology, cell-biology and biotechnology. (textbooks)*. Ben-Gurion University of the Negev. Israel. 1, 1-15, 1997.
- [12] Hadiyanto, M.M.A. Nur, dan G.D. Hartanto, Cultivation of *Chlorella* sp. as Biofuel Sources in Palm Oil Mill Effluent (POME). *Int. Journal of Renewable Energy Development*. 1 (2), 45-49, 2012.
- [13] N. Rahmawati, Yuliani dan E. Ratnasari. Pengaruh Pupuk Kompos Berbahan Campuran Limbah Cair Tahu, Daun Lamtoro dan Rumen Sapi sebagai Media Kultur terhadap Kepadatan Populasi *Spirulina* sp. *Jurnal Lentera Bio*. 1 (1), 17-23, 2012.
- [14] A.W. Leksono, D. Mutiara, dan I.A. Yusanti. Penggunaan Pupuk Organik Cair Hasil Fermentasi dari *Azolla pinnata* Terhadap Kepadatan Sel *Spirulina* sp. *Jurnal Ilmu-ilmu Perikanan dan Budidaya Perairan*. 12 (1), 2017.
- [15] T. Chrismadha, P. Lily, dan M. Yayah. Pengaruh Konsentrasi Nitrogen dan Fosfor terhadap Pertumbuhan, Kandungan Protein, Karbohidrat dan Fikosianin pada Kultur *Spirulina fusiformis*. *Berita Biologi*. 8 (3), 2006.
- [16] D.R. Amanatin dan T. Nurhidayati, Pengaruh Kombinasi Konsentrasi Media Ekstrak Tauge (MET) dengan Pupuk Urea terhadap Kadar Protein *Spirulina* sp. *Jurnal Sains dan Seni Pomits*. 2 (2), 2337-3520, 2013.
- [17] X. Yuan, A. Kumar, A.K Sahu., dan S.J.Ergas. Impact of ammonia concentration on *Spirulina platensis* growth in an airlift photobioreactor. *Bioresource Technology*. 102, 3234–3239, 20
- [18] H. Effendi. *Telaah Kualitas Air*, Kanisius, Yogyakarta, 2003.

# Optimization of Spirulina platensis Culture Media as an Effort for Utilization of Pangasius Farming Waste Water

---

ORIGINALITY REPORT

---

13%

SIMILARITY INDEX

---

PRIMARY SOURCES

---

- 1 E Setyaningsih, Kismiyati, S Subekti. " The effect of noni fruits ( ) with different ripeness stages against the total erythrocytes and leukocytes of comet goldfish ( ) infested by ", IOP Conference Series: Earth and Environmental Science, 2019  
Crossref 44 words — 1%
- 2 [academicjournals.org](http://academicjournals.org)  
Internet 39 words — 1%
- 3 [doaj.org](http://doaj.org)  
Internet 38 words — 1%
- 4 Claudia Anahite Ovando, Julio Cesar de Carvalho, Gilberto Vinícius de Melo Pereira, Philippe Jacques et al. " Functional properties and health benefits of bioactive peptides derived from : A review ", Food Reviews International, 2016  
Crossref 35 words — 1%
- 5 [mail.scialert.net](http://mail.scialert.net)  
Internet 24 words — 1%
- 6 Yelmira Zalfiatri, Fajar Restuhadi, Angga Pramana, Fajar Yuliandri. "Decreasing Levels of BOD, COD and Oil in Palm Oil Mill Effluent (Pome) Using Scale Up Experiment of Symbiosis Mutualism Technology Between Microalgae 22 words — 1%

Chorella Sp and Agrobost", Journal of Applied Agricultural Science and Technology, 2020

Crossref

---

7 Pan, Shanglei. "Bioremediation of Landfill Leachate Using an Algal-Based System: An Application of Red Alga Galdieria sulphuraria", Lamar University - Beaumont, 2021 17 words — 1%  
ProQuest

---

8 [www.coursehero.com](http://www.coursehero.com) 16 words — 1%  
Internet

---

9 Jun An, Fuli Gao, Qiuyue Ma, Yang Xiang, Difeng Ren, Jun Lu. "Screening for enhanced astaxanthin accumulation among Spirulina platensis mutants generated by atmospheric and room temperature plasmas", Algal Research, 2017 15 words — < 1%  
Crossref

---

10 Van Chung Mai, Kinga Drzewiecka, Henryk Jeleń, Dorota Narożna et al. "Differential induction of Pisum sativum defense signaling molecules in response to pea aphid infestation", Plant Science, 2014 15 words — < 1%  
Crossref

---

11 [bayjournal.com](http://bayjournal.com) 15 words — < 1%  
Internet

---

12 M Ulumiah, M Lamid, K T Pursetyo. "Substitution of commercial feed with maggot meal (Hermetia illucens) to the growth rate, feed conversion ratio and feed efficiency catfish (Pangasius pangasius)", IOP Conference Series: Earth and Environmental Science, 2021 14 words — < 1%  
Crossref

---

13 Suvidha Gupta, Sanjay B. Pawar, R.A. Pandey. "Current practices and challenges in using 13 words — < 1%



microalgae for treatment of nutrient rich wastewater from agro-based industries", *Science of The Total Environment*, 2019

Crossref

14 Sang-Hyo Lee, Ju Eun Lee, Yoori Kim, Seung-Yop Lee. "The Production of High Purity Phycocyanin by *Spirulina platensis* Using Light-Emitting Diodes Based Two-Stage Cultivation", *Applied Biochemistry and Biotechnology*, 2015

12 words — < 1%

Crossref

15 Shinta Elystia, Sri Rezeki Muria, Herta Furaida Erlangga. "Cultivation of *Chlorella pyrenoidosa* as a raw material for the production of biofuels in palm oil mill effluent medium with the addition of urea and triple super phosphate", *Environmental Health Engineering and Management*, 2020

11 words — < 1%

Crossref

16 Zhang, Xin, Hui Chen, Weixian Chen, Yaqin Qiao, Chenliu He, and Qiang Wang. "Evaluation of an Oil-Producing Green Alga *Chlorella* sp. C2 for Biological DeNO<sub>x</sub> of Industrial Flue Gases", *Environmental Science & Technology*

10 words — < 1%

Crossref

17 [core.ac.uk](http://core.ac.uk)

Internet

9 words — < 1%

18 Kargin Yilmaz Hilal, Sezgin zlem. "Production of *Spirulina platensis* by adding sodium bicarbonate and urea into chicken manure medium", *African Journal of Biotechnology*, 2014

8 words — < 1%

Crossref

19 Li, Z.Y.. "Effects of electromagnetic field on the batch cultivation and nutritional composition of

8 words — < 1%

Spirulina platensis in an air-lift photobioreactor", Bioresource Technology, 200702

Crossref

20 Markou, G.. "Cultivation of filamentous cyanobacteria (blue-green algae) in agro-industrial wastes and wastewaters: A review", Applied Energy, 201110 8 words — < 1%

Crossref

21 [jalgalbiomass.com](http://jalgalbiomass.com) 8 words — < 1%

Internet

22 Fei Luo Jin, Ping Jiang Lu. "Production of aquatic feed grade algal powder from turtle breeding wastewater using a locally isolated Spirulina sp. JXSC-S1", African Journal of Microbiology Research, 2015 7 words — < 1%

Crossref

23 Fernando Martínez-Jerónimo, Dante Ivo Flores-Hernández, Juvencio Galindez-Mayer. "Formulation of an Alternative Culture Medium for ( ) Based on "Tequesquite," a Traditional Mineral Resource ", Journal of the World Aquaculture Society, 2017 7 words — < 1%

Crossref

24 H Kurnia, R R Rifadi, Agustono, M N G Amin, S A Sudjarwo, M A Alamsjah. " The potential of seaweed waste ( sp. and ) as a medium density fiberboard (mdf)-based pot material for better water use efficiency in tomato plants ", IOP Conference Series: Earth and Environmental Science, 2019 7 words — < 1%

Crossref

25 Beas, Rosa Elena Yaya. "Bio-Filtration of Helminth Eggs and Coliforms from Municipal Sewage for Agricultural Reuse in Peru.", Wageningen University and Research, 2021 6 words — < 1%

ProQuest

---

26 Budiyo ., Iqbal Syaichurro, Siswo Sumardiono, Setia Budi Sasongko. "Production of Spirulina platensis Biomass Using Digested Vinasse as Cultivation Medium", Trends in Applied Sciences Research, 2014

6 words — < 1%

Crossref

---

EXCLUDE QUOTES ON

EXCLUDE MATCHES OFF

EXCLUDE BIBLIOGRAPHY ON