



ISSN: 1412-033X
E-ISSN: 2085-4722

BIODIVERSITAS

Journal of Biological Diversity

Volume 23 - Number 1 - January 2022

Front cover: Panthera tigris sumatrae (Pocock, 1929)
(PHOTO: SMUDGE 9000)

Published monthly

PRINTED IN INDONESIA

ISSN: 1412-033X

E-ISSN: 2085-4722



9 771412 033849



9 772085 472843

BIODIVERSITAS

Journal of Biological Diversity
Volume 23 – Number 1 – January 2022

ISSN/E-ISSN:

1412-033X (printed edition), 2085-4722 (electronic)

EDITORIAL BOARD:

Abdel Fattah N.A. Rabou (Palestine), **Agnieszka B. Najda** (Poland), **Ajay Kumar Gautam** (India), **Alan J. Lymbery** (Australia), **Annisa** (Indonesia), **Bambang H. Saharjo** (Indonesia), **Daiane H. Nunes** (Brazil), **Darlina Md. Naim** (Malaysia), **Ghulam Hassan Dar** (India), **Hassan Pourbabaie** (Iran), **Joko R. Witono** (Indonesia), **Kartika Dewi** (Indonesia), **Katsuhiko Kondo** (Japan), **Kusumadewi Sri Yulita** (Indonesia), **Livia Wanntorp** (Sweden), **M. Jayakara Bhandary** (India), **Mahdi Reyahi-Khoram** (Iran), **Mahendra K. Rai** (India), **Mahesh K. Adhikari** (Nepal), **Maria Panitsa** (Greece), **Mochamad A. Soendjoto** (Indonesia), **Mohamed M.M. Najim** (Srilanka), **Mohib Shah** (Pakistan), **Nurhasanah** (Indonesia), **Praptiwi** (Indonesia), **Rasool B. Tareen** (Pakistan), **Sayed Aliakbar Hedayati** (Iran), **Sayed Mehdi Talebi** (Iran), **Shahabuddin** (Indonesia), **Shahir Shamsir** (Malaysia), **Shri Kant Tripathi** (India), **Subhash C. Santra** (India), **Sugeng Budiharta** (Indonesia), **Sugiyarto** (Indonesia), **Taufiq Purna Nugraha** (Indonesia), **Yosep S. Mau** (Indonesia)

EDITOR-IN-CHIEF:

S u t a r n o

EDITORIAL MEMBERS:

English Editors: **Graham Eagleton** (grahameagleton@gmail.com), **Suranto** (surantouns@gmail.com); Technical Editor: **Solichatun** (solichatun_s@yahoo.com), **Artini Pangastuti** (pangastuti_tutut@yahoo.co.id); Distribution & Marketing: **Rita Rakhmawati** (oktia@yahoo.com); Webmaster: **Ari Pitoyo** (aripitoyo@yahoo.com)

MANAGING EDITORS:

Ahmad Dwi Setyawan (unsjournals@gmail.com)

PUBLISHER:

The Society for Indonesian Biodiversity

CO-PUBLISHER:

Department of Biology, Faculty of Mathematics and Natural Sciences, Sebelas Maret University, Surakarta

ADDRESS:

Jl. Ir. Sutami 36A Surakarta 57126. Tel. +62-271-7994097, Tel. & Fax.: +62-271-663375, email: editors@smujo.id

ONLINE:

biodiversitas.mipa.uns.ac.id; smujo.id/biodiv

.....



**Society for Indonesia
Biodiversity**



**Sebelas Maret University
Surakarta**

GUIDANCE FOR AUTHORS

Aims and Scope *Biodiversitas*, *Journal of Biological Diversity* or *Biodiversitas* encourages submission of manuscripts dealing with all aspects of biodiversity including plants, animals and microbes at the level of the gene, species, and ecosystem. Ethnobiology papers are also considered.

Article types The journal seeks original full-length: (1) **Research papers**, (2) **Reviews**, and (3) **Short communications**. Original research manuscripts are limited to 8,000 words (including tables and picture), or proportional with articles in this publication number. Review articles are also limited to 8,000 words, while Short communications should be less than 2,000 words, except for pre-study.

Submission The journal only accepts online submission, through open journal system (<https://smujo.id/biodiv/about/submissions>) or email to the editors at unsjournals@gmail.com. Submitted manuscripts should be the original works of the author(s). Please ensure that the manuscript is submitted using the *Biodiversitas* template, which can be found at (<https://biodiversitas.mipa.uns.ac.id/D/guidance.htm>). The manuscript must be accompanied by a cover letter containing the article title, the first name and last name of all the authors, a paragraph describing the claimed novelty of the findings versus current knowledge. Please also provide a list of five potential reviewers in your cover letter. Submission of a manuscript implies that the submitted work has not been published before (except as part of a thesis or report, or abstract); and is not being considered for publication elsewhere. When a manuscript written by a group, all authors should read and approve the final version of the submitted manuscript and its revision; and agree the submission of manuscripts for this journal. All authors should have made substantial contributions to the concept and design of the research, acquisition of the data and its analysis; drafting of the manuscript and correcting of the revision. All authors must be responsible for the quality, accuracy, and ethics of the work.

Ethics Author(s) must obedient to the law and/or ethics in treating the object of research and pay attention to the legality of material sources and intellectual property rights.

Copyright If the manuscript is accepted for publication, the author(s) still hold the copyright and retain publishing rights without restrictions. Authors are allowed to reproduce articles as long as they are not used for commercial purposes. For the new invention, authors are suggested to manage its patent before published.

Open access The journal is committed to free-open access that does not charge readers or their institutions for access. Readers are entitled to read, download, copy, distribute, print, search, or link to the full texts of articles, as long as not for commercial purposes. The license type is CC-BY-NC-SA.

Acceptance Only articles written in U.S. English are accepted for publication. Manuscripts will be reviewed by editors and invited reviewers (double blind review) according to their disciplines. Authors will generally be notified of acceptance, rejection, or need for revision within 1 to 2 months of receipt. Manuscripts will be rejected if the content does not in line with the journal scope, does not meet the standard quality, is in an inappropriate format, contains complicated grammar, dishonesty (i.e. plagiarism, duplicate publications, fabrication of data, citations manipulation, etc.), or ignoring correspondence in three months. The primary criteria for publication are scientific quality and biological significance. **Uncorrected proofs** will be sent to the corresponding author by email as .doc or .docx files for checking and correcting of typographical errors. To avoid delay in publication, corrected proofs should be returned in 7 days. The accepted papers will be published online in a chronological order at any time, but printed in January, April, July and October.

A charge Starting on January 1, 2017, publishing costs waiver is granted to foreign (non-Indonesian) authors who first publish the manuscript in this journal, especially for graduate students from developing countries. However, other authors are charged USD 250 (IDR 3,500,000).

Reprints The sample journal reprint is only available by special request. Additional copies may be purchased when ordering by sending back the uncorrected proofs by email. Manuscript preparation Manuscript is typed on A4 (210x297 mm²) paper size, in a single column, single space, 10-point (10 pt) Times New Roman font. The margin text is 3 cm from the top, 2 cm from the bottom, and 1.8 cm from the left and right. Smaller lettering size can be applied in presenting table and figure (9 pt). Word processing program or additional software can be used, however, it must be PC compatible, use the *Biodiversitas* template, and Microsoft Word based (.doc or .rtf; not .docx). **Scientific names** of species (incl. subspecies, variety, etc.) should be written in italics, except in italicised sentences. Scientific names (Genus, species, author), and cultivar or strain should be mentioned completely for the first time mentioning it in the body text, especially for taxonomic manuscripts. The Genus name can be shortened after first mention, except where this may generate confusion. Name of the author can be eliminated after first mentioning. For example, *Rhizopus oryzae* L. UICC 524, hereinafter can be written as *R. oryzae* UICC 524. Using trivial names should be avoided. **Biochemical and chemical nomenclature** should follow the order of the IUPAC - IUB. For DNA sequence, it is better used Courier New font. Standard chemical abbreviations can be applied for common and clear used, for example, completely written butilic hydroxyl toluene (BHT) to be BHT hereinafter. **Metric measurements** should use IS denominations, and other system should use equivalent values with the denomination of IS mentioned first. Abbreviations like g, mg, mL, etc. should not be followed by a dot.

Minus index (m-2, L-1, h-1) suggested to be used, except in things like "per-plant" or "per-plot". **Mathematical equations** can be written down in one column with text, in that case can be written separately. **Numbers** one to ten are written in words, except if it relates to measurement, while values above them written in number, except in early sentence. The fraction should be expressed in decimal. In the text, it should be used "%" rather than "percent". Avoid expressing ideas with complicated sentence and verbiage, and used efficient and effective sentence.

The **Title** of the article should be written in compact, clear, and informative sentence, preferably not more than 20 words. Author name(s) should be completely written. Name and institution address should also be completely written with street name and number (location), postal code, telephone number, facsimile number, and email address. Manuscripts written by a group, author for correspondence along with address is required. First page of the manuscript is used for writing above information.

The **Abstract** should not be more than 200 words. Include between five and eight **Keywords**, using both scientific and local names (if any), research themes, and special methods which used; and sorted from A to Z. All important **abbreviations** must be defined at their first mention. Running title is about five words. The **Introduction** is about 400-600 words, covering the background and aims of the research. **Materials and Methods** should emphasize on the procedures and data analysis. **Results and Discussion** should be written as a series of connecting sentences, however, for manuscript with long discussion should be divided into subtitles. Thorough discussion represents the causal effect mainly explains for why and how the results of the research were taken place, and do not only re-express the mentioned results in the form of sentences. A **Conclusion** should be given at the end of the discussion. **Acknowledgments** are expressed in brief; all sources of institutional, private and corporate financial support for the work must be fully acknowledged, and any potential conflicts of interest must be noted.

Figures and Tables of three pages maximum should be clearly presented. Include a label below each figure, and a label above each table (see example). Colored figures can only be accepted if the information in the manuscript can lose without those images; chart is preferred to use black and white images. Author could consign any picture or photo for the front cover, although it does not print in the manuscript. All images property of others should be mentioned source. There is no **Appendix**, all data or data analysis are incorporated into Results and Discussions. For broad data, supplementary information can be provided on the website.

References In the text give the author names followed by the year of publication and arrange from oldest to newest and from A to Z. In citing an article written by two authors, both of them should be mentioned, however, for three and more authors only the first author is mentioned followed by et al., for example: Saharjo and Nurhayati (2006) or (Boonkerd 2003a, b, c; Sugiyarto 2004; El-Bana and Nijs 2005; Balagadde et al. 2008; Webb et al. 2008). Extent citation as shown with word "cit" should be avoided. Reference to unpublished data and personal communication should not appear in the list but should be cited in the text only (e.g., Rifai MA 2007, pers. com. (personal communication); Setyawan AD 2007, unpublished data). In the reference list, the references should be listed in an alphabetical order. Names of journals should be abbreviated. Always use the standard abbreviation of a journal's name according to the **ISSN List of Title Word Abbreviations** (www.issn.org/2-22661-LTWA-online.php). Please include DOI links for journal papers. The following examples are for guidance.

Journal:

Saharjo BH, Nurhayati AD. 2006. Domination and composition structure change at hemic peat natural regeneration following burning; a case study in Pelalawan, Riau Province. *Biodiversitas* 7: 154-158. DOI: 10.13057/biodiv/d070213

Book:

Rai MK, Carpinella C. 2006. Naturally Occurring Bioactive Compounds. Elsevier, Amsterdam.

Chapter in book:

Webb CO, Cannon CH, Davies SJ. 2008. Ecological organization, biogeography, and the phylogenetic structure of rainforest tree communities. In: Carson W, Schnitzer S (eds) *Tropical Forest Community Ecology*. Wiley-Blackwell, New York.

Abstract:

Assaeed AM. 2007. Seed production and dispersal of *Rhazya stricta*. 50th annual symposium of the International Association for Vegetation Science, Swansea, UK, 23-27 July 2007.

Proceeding:

Alikodra HS. 2000. Biodiversity for development of local autonomous government. In: Setyawan AD, Sutarno (eds.) *Toward Mount Lawu National Park; Proceeding of National Seminary and Workshop on Biodiversity Conservation to Protect and Save Germplasm in Java Island*. Universitas Sebelas Maret, Surakarta, 17-20 July 2000. [Indonesian]

Thesis, Dissertation:

Sugiyarto. 2004. Soil Macro-invertebrates Diversity and Inter-Cropping Plants Productivity in Agroforestry System based on Sengon. [Dissertation]. Universitas Brawijaya, Malang. [Indonesian]

Information from internet: Balagadde FK, Song H, Ozaki J, Collins CH, Barnett M, Arnold FH, Quake SR, You L. 2008. A synthetic *Escherichia coli* predator-prey ecosystem. *Mol Syst Biol* 4:187. www.molecularsystembiology.com

Antibacterial activity of endophytic fungi isolated from the stem bark of jambu mawar (*Syzygium jambos*)

KURRATUL 'AINI^{1,2*}, ELFITA^{3,4*}, HARY WIDJAJANTI⁴, ARUM SETIAWAN⁴, ALFIA R. KURNIAWATI²

¹Graduate School of Sciences, Faculty of Mathematics and Natural Sciences, Universitas Sriwijaya. Jl. Padang Selasa No. 524, Palembang 30139, South Sumatra, Indonesia. *email: kurratulaini_uin@radenfatah.ac.id

²Universitas Islam Negeri Raden Fatah Palembang. Jl. Prof. K.H. Zainal Abidin Fikri Km. 3, Palembang 30126, South Sumatra, Indonesia

³Department of Chemistry, Faculty of Mathematics and Natural Sciences, Universitas Sriwijaya. Jl. Raya Palembang-Prabumulih Km 32, Indralaya, Ogan Ilir 30662, South Sumatra, Indonesia. Tel.: +62-711-580056, *email: elfita.elfita.69@gmail.com

⁴Department of Biology, Faculty of Mathematics and Natural Sciences, Universitas Sriwijaya. Jl. Raya Palembang-Prabumulih Km 32, Indralaya, Ogan Ilir 30662, South Sumatra, Indonesia

Manuscript received: 3 September 2021. Revision accepted: 30 December 2021.

Abstract. 'Aini K, Elfita, Widjajanti H, Setiawan A, Kurniawati AR. 2021. Antibacterial activity of endophytic fungi isolated from the stem bark of jambu mawar (*Syzygium jambos*). *Biodiversitas* 23: 521-532. *Syzygium jambos* (jambu mawar) is widely used in South Sumatra as a traditional medicine to treat various diseases, including pathogenic bacterial infections. Literature studies report that parts of *S. jambos* have been used worldwide for treating diarrhea accompanied by fever, dysentery, sore throat, diabetes, and other infectious diseases. Endophytic fungi isolated from medicinal plants have high diversity, and the biological activity of their secondary metabolites is associated with their host. Therefore, this study aimed to determine the diversity of endophytic fungi from the stem bark of *S. jambos* and the antibacterial activity of secondary metabolites of endophytic fungi. Fungal isolates with high antibacterial activity were identified to obtain fungal species, and the structure of their compounds was determined. The disc diffusion method carried the antibacterial activity test against Gram-positive (*Staphylococcus aureus* and *Bacillus subtilis*) and Gram-negative (*Escherichia coli* and *Salmonella typhi*) bacteria. Characteristics of endophytic fungi were identified microscopically and macroscopically. Active endophytic fungi were identified by molecular analysis of the internal transcribed region (ITS) of the ribosomal DNA. Isolation of antibacterial compounds using chromatographic techniques and determination of chemical structures using spectroscopy was performed. Eight endophytic fungi were obtained from the stem bark of *S. jambos*, namely SJ1–SJ8. The endophytic fungus SJ6 showed the highest activity and based on phylogenetic analysis, was identified as *Fusarium verticillioides*. Pure compounds isolated as yellowish-white solids showed good antibacterial activity against *S. aureus* and *S. typhi* with minimum inhibitory concentration values of 64 g/mL each. Based on 1D and 2D nuclear magnetic resonance spectroscopic analysis, compound 1 was identified as 3-hydroxy-4-(hydroxy(4-hydroxyphenyl)methyl) dihydrofuran-2-one.

Keywords: Antibacterial compound, endophytic fungi, *Fusarium verticillioides*, *Syzygium jambos*

INTRODUCTION

Most of the diseases that arise today are caused by microbial infections. Microbes that cause infection have become increasingly resistant to drugs over time, posing a major challenge to human health. New bioactive compounds from plants and microorganisms can provide the best alternative as a potential drug source to combat these infectious diseases (Sudha et al. 2016). Plants are the main source of bioactive compounds as antibacterial agents such as alkaloids, phenolics, polyphenols, flavonoids, tannins, terpenes (Katz and Baltz 2016; Anand et al. 2019). Endophytic fungi play an important role in discovering these new bioactive compounds. They live in healthy plant tissues without disease symptoms, protecting plants from herbivores and pathogens by producing bioactive secondary metabolites in host tissues (Rai et al. 2012, Martinez-Klimova et al. 2017; Abdel-Azeem et al. 2019).

In India, the fruit and leaves of *Syzygium jambos* are used as a traditional treatment for diuretics, toothaches, eye pain, rheumatism and as an expectorant. In Indo-China, its flowers are used to reduce fever, and its seeds are used to

treat diabetes, diarrhea, and dysentery (Bonfanti et al. 2013, 2014; Sobeh et al. 2018; Rajkumari et al. 2018). Also, its leaf extract shows anti-herpes simplex virus activity and has antibacterial, antitoxic, antioxidant, and hepatoprotective properties (Athikomkulchai et al. 2008; Mohanty and Cock 2010; Ghareeb et al. 2016; Devakumar and Sudha 2017). A decoction of rose apple leaves and bark is used to treat diarrhea, rheumatism, asthma, bronchitis, and hoarseness and as an expectorant. Its leaf juice is used as a febrifuge. In some countries, such as Suriname, China, India, and Cambodia, its leaves, seeds, and fruit are used to remedy diarrhea, diarrhea with fever, dysentery, sore throat, diabetes, and other infectious diseases (Morton 1987; Lim 2012).

Endophytic fungi are widespread in many plant species. Many strains of endophytic fungi can produce various natural products that are active as antimicrobials. Endophytic fungi are a source of novel bioactive compounds with diverse biological activities. The bioactive compounds of endophytic fungi isolated from medicinal plants are known as volatile derivatives, such as esters, ethers, alkaloids, and phenolic compounds (Nisa et al.

2020). In the literature, several studies have been made on the endophytic fungi of the genus *Syzygium*, including *S. jambos*. *Chrysosporthe deuterocubines*, *Microthia havanensis*, *Celoportha hauliensis* sp. nov., *C. hawaiiensis* sp. nov., *C. paradisiaca* sp. nov. found in the bark of *S. jambos* (Roux et al. 2020). However, there are no reports of its antibacterial activity and its secondary metabolites. Therefore, this study aimed to reveal the diversity of endophytic fungi from *S. jambos* from South Sumatra, Indonesia, their antibacterial activity, and their bioactive compounds.

MATERIALS AND METHODS

Plant material

The stem bark of *S. jambos* (L.) Alston. was collected at the Multipurpose Field, Jalan Sakura 3 Kompleks Kencana Damai, Kelurahan Sukamaju, Kecamatan Sako, and Kota Palembang. The samples were morphologically identified at the Biosystematics Laboratory, Department of Biology, Faculty of Mathematics and Natural Sciences, Sriwijaya University, Indonesia. ID number: 233/UN9.1.7/4/EP/2021.

Isolation of endophytic fungi

The stem bark of *S. jambos* was washed thoroughly under running tap water. The surface of the sample was sterilized by dipping it into 70% ethanol for 2 min and then soaking it in sodium hypochlorite (NaOCl) for 1 min. Then, the sample was rinsed with sterile distilled water three times (2 min each time) and dried using sterile filter paper. Each sample explant (without epidermis) was aseptically cut to a size of $2 \times 1 \text{ cm}^2$ and placed on potato dextrose agar (PDA) media (200 g/L potatoes, 20 g/L dextrose, and 20 g/L agar) to which 200 g of chloramphenicol was added to prevent bacterial growth. Samples were incubated in the dark at $\pm 30^\circ\text{C}$ for 1–2 weeks. The hyphal tips of several different colonies emerging from the segments were isolated and subcultured on new PDA media to obtain pure colonies (Elfita et al. 2019; Khalil et al. 2021).

Identification of endophytic fungi

Characteristics of endophytic fungal isolates were identified macroscopically and microscopically. Macroscopic characterization depends on the colony growth pattern, texture, color margin, and other features. Microscopic characterization was performed using the culture slide method in which one drop of lactophenol blue reagent was added. The slides were examined using a light microscope (Hirox MXB-2500REZ). The characteristic data are then compared with identification books for fungi (Habisukan et al. 2021), such as Larone's Medically Important Fungi (Walsh et al. 2018), Pictorial Atlas of Soil and Seed Fungi (Watanabe 2010) and other journals of fungal identification.

Cultivation and extraction

Endophytic fungi culture (5–7 days) on PDA medium was cut into small pieces ($\pm 5 \text{ mm}$ in diameter), and six pieces were transferred into each 1-L culture bottle containing 300 mL of potato dextrose broth medium (ingredients: 200 g potato, 20 g dextrose in 1 L of water). The number of culture bottles for each endophytic fungal isolate is five. After incubation at room temperature for 4 weeks, fungal liquid culture was vacuum filtered and partitioned in a separating funnel with ethyl acetate (1:1) solvent three times. The ethyl acetate extract was evaporated to obtain a concentrated extract (Habisukan et al. 2021). The chemical procedure for obtaining organic constituents from plant tissue uses solvent ethyl acetate for more polar compounds (Wan 2017).

Antibacterial activity test

The antibacterial test was carried out using the disc diffusion method. Each sample of the fungal extract was dissolved in dimethyl sulfoxide (DMSO, Merck, Germany). Antibacterial activity was tested on Gram-positive bacteria (*Staphylococcus aureus* InaCCB4 and *Bacillus subtilis* InaCCB4) and Gram-negative bacteria (*Escherichia coli* InaCCB5 and *Salmonella typhi* ATCC 1408) on nutrient agar media. The extract concentration tested was 400 $\mu\text{g}/\text{disc}$, using tetracycline antibiotics with a concentration of 30 $\mu\text{g}/\text{disc}$ as a positive control. The activity of the bacterial inhibition zone was measured after incubation at 37°C for 24 hours by measuring the clear zone formed around the paper disc. The clear zone formed indicates the bacteria's sensitivity to the test material's antibacterial material, which is expressed by the width of the diameter of the inhibition zone. The following equation determines the antibacterial activity of the test sample and the criteria for the inhibition zone diameter (Elfita et al. 2019).

The minimum inhibitory concentration (MIC) of pure compounds is determined by preparing a concentration series of 256 mg/mL to 1 g/mL. If the diameter of the inhibition zone is $> 9 \text{ mm}$ at a certain concentration series, the antibacterial activity at that concentration is positive. The lowest concentration of the concentration series positive for antibacterial activity is defined as the MIC value (Ding et al. 2019).

$$\text{Weak: } \frac{A}{B} \times 100\% < 50\%; \text{Moderate: } 50\% < \frac{A}{B} \times 100\% < 70\%; \text{Strong: } \frac{A}{B} \times 100\% > 70\%$$

Where:

A = zone of inhibition (mm) test sample

B = zone of inhibition (mm) standard antibiotic

Molecular analysis of active endophytic fungi

Molecular analysis were identified at Genetika Lab, PT Genetika Science Indonesia. Genomic DNA extraction method with Quick-DNA Fungal/Bacterial Miniprep Kit (Zymo Research, D6005). PCR amplification with (2x) MyTaq HS Red Mix (Bioline, BIO-25048) using universal primers ITS1 5'-TCCGTAGGTGAACCTGCGG-3' and ITS4 5'-TCCTCCGCTTATTGATATGC-3' (White et al.

1990). The sequence results were identified of species using the Basic Local Alignment Search Tool (BLAST) at the website address <http://blast.ncbi.nlm.nih.gov/Blast.cgi>. Furthermore, multiple alignments were performed using the Mega 11 program (Tamura et al. 2021) use the CLUSTAL W method and building phylogenetic tree using the Neighbor-joining tree method with a bootstrap value of 1000 (Katoch and Pull 2017; Kuswytasari et al. 2019; Saitou and Nei 1987). The percentage of replicate trees in which the associated taxa clustered together in the bootstrap test (1000 replicates) are shown next to the branches (Felsenstein 1985). The tree is drawn to scale, with branch lengths in the same units as the evolutionary distances used to infer the phylogenetic tree. The evolutionary distances were computed using the number of differences method (Nei and Kumar 2000) and are in the units of the number of base differences per sequence.

Isolation and identification of bioactive compound

The ethyl acetate extract, which showed strong antibacterial activity, was analyzed by thin-layer chromatography (TLC) of silica gel G-60 F 254, using solvents with various eluents to observe the staining pattern. The extract was subjected to column chromatography using a stationary phase of silica gel (70-230 mesh). The samples prepared by pre-absorption were eventually put into the chromatographic column and eluted using a gradient eluent. The eluate was collected in vials every 10 ml, and thin-layer chromatography was performed to group into column fractions based on the staining pattern. The fraction containing potential secondary metabolites was further chromatographed by column and rinsed with a suitable solvent to obtain a pure compound, namely compound 1. The structure of the pure compound (compound 1) was identified by spectroscopic methods, including ^1H -NMR, ^{13}C -NMR, HMQC, HMBC, and COSY (Fadhilah et al. 2019).

RESULTS AND DISCUSSION

Isolation of endophytic fungi

Endophytic fungi from the bark of *S. jambos* were obtained in eight isolates with codes SJ1 to SJ8 (Figure 1). Macroscopic characteristics were identified by the following criteria: color, texture, topography, pattern, exudate drops, radial line, and concentric circle. Microscopic characteristics

were identified by the following criteria: spore type, spore shape, hyphae, and other specific characteristics. The results of this identification were compared with the literature and the key to determining fungi (Watanabe 2010).

Endophytic fungi SJ1: brownish-white and dark brown, cottony, raised, zonate, erect conidiophores of variable height, apical conidia, transverse and longitudinal septa, narrowed setae near the septa, identified as *Spegazzinia tessartha* (Cole 1974; Watanabe 2010; Mena-Portales et al. 2017). Endophytic fungi SJ2: white and white-middle yellow, cottony, raised, pater arrowed, aerial mycelia abundant, hyphae hyaline, septate, branched, terminal or lateral arising hyaline conidiogenous cells, asci globose, identified as *Auxarthronopsis* sp. (Sharma et al. 2013a; Zhang et al. 2017).

Endophytic fungi SJ3: gray-green and yellowish-green, velvety, raised and flowered, ascospore, globose, septate, hyphae hyaline to brown, scattered, according to the characteristics of *Pectinotrichum* sp. (Varsavsky and Orr 1971; Zhang et al. 2017). Endophytic fungi SJ4: light brown, velvety, umbonate, radiated, concentric circle, ascospore, subglobose, septate, paraphyses hyaline, cylindrical, curved, thread-like, according to the characteristics *Phaeotrichosphaeria* (Sivanesan 1983; Watanabe 2010).

Endophytic fungi SJ5: white to gray with yellow edges and white and yellow around, cottony, raised, radiated, concentric circle, conidia, globose, septate, frequently intercalary in hyphae sole, identified as *Poaceascoma* (Phookamsak et al. 2015; Luo et al. 2016; Fors et al. 2020). Endophytic fungus SJ6: pink with a lighter periphery, cottony, zonate, conidiophores hyaline, erect, long, branched verticillately, rarely simple, conidia lunate shaped, long elliptical, 4-5 celled identified as *Fusarium verticillioides* (Walsh et al. 2018; Watanabe 2010).

Endophytic fungi SJ7: green-white around and dark green to gray, cottony, raised, spreading, conidia and globose spores, coenocytic hyphae, catenulate bearing conidia and chlamydospores apically, identified as *Thielaviopsis* sp. (Watanabe 2010; Álvarez et al. 2012; Majumdar and Mandal 2018). Endophytic fungi SJ8: black and white, cottony, umbonate, zonate, concentric circle, conidia and globose spores, conidiogenous cells giving rise to conidia, septate hyphae, these endophytes identified as *Nigrospora bambusae* (Wang et al. 2017). Macroscopic and microscopic identification of endophytic fungi isolated from the bark of *S. jambos* is shown in Tables 1 and 2, respectively.

Table 1. Macroscopic characteristics of endophytic fungi isolate SJ1 – SJ8

Isolate	Colony color	Reverse colony color	Texture	Topography	Pattern	Concentric circle
SJ1	Brownish white	Dark brown	Cottony	Raised	Zonate	-
SJ2	White	White-middle yellow yellow	Cottony	Raised	Arrowed	-
SJ3	Grey-green	Yellowish green	Velvety	Raised	Flowery	-
SJ4	Light brown	Light brown	Velvety	Umbonate	Radiated	√
SJ5	white to gray with yellow edges	White and yellow around	Cottony	Raised	Radiated	√
SJ6	Pink and white around	Pink and white around	Cottony	Raised	Zonate	-
SJ7	Green-white around	Dark green to gray	Cottony	Raised	Radiated	-
SJ8	Black and White around	Black and White around	Cottony	Umbonate	Zonate	√

Note: (-) = characteristic doesn't appear; (√) = characteristic appear




Figure 1. Morphological identification of endophytic fungi from the stem bark of *S. jambos* macroscopically and microscopically

Table 2. Microscopic characteristics of endophytic fungi isolate SJ1 – SJ8

Isolate	Type of spore	Shape of spore	Hyphae	Specific characteristic	Genus / species
SJ1	Conidia	Globose	Septate	Hyaline conidiophores, simple, cylindrical, apically narrowed.	<i>Spegazzinia tessarthra</i>
SJ2	Conidia	Globose	Septate	Hyphae hyaline to brown, branched, thin-walled aerial mycelia abundant.	<i>Auxarthronopsis</i> sp.
SJ3	Ascospore	Globose	Septate	Hyphae hyaline to brown, scattered	<i>Pectinotrichum</i> sp.
SJ4	Ascospore	Subglobose	Septate	Paraphyses hyaline, cylindrical, curved, thread-like	<i>Phaeotrichosphaeria</i> sp.
SJ5	Conidia	globose	Septate	frequently intercalary in hyphae sole	<i>Poaceascoma</i> sp.
SJ6	Conidia	Verticillate	Septate	Conidia lunate with a foot cell, long elliptical, 4-5 celled	<i>Fusarium verticillioides</i>
SJ7	Conidia	globose	Coenocytic	Bearing catenulate conidia and chlamydospores apically	<i>Thielaviopsis</i> sp.
SJ8	Conidia	Globose	Septate	conidiogenous cells giving rise to conidia Scale bars: d–g = 10 μ m.	<i>Nigrospora bambusae</i>

Table 3. Antibacterial activity of ethyl acetate extract of the endophytic fungus *Staphylococcus jambos* (SJ1-SJ8) compared to standard antibiotics (tetracyclines)

Sample (400 µg/disc)	The comparison of the inhibition zone (mm) of the sample to the inhibition zone of tetracycline antibiotics (mm) expressed in % antibacterial activity ^a			
	<i>Staphylococcus aureus</i>	<i>Salmonella typhi</i>	<i>Escherichia coli</i>	<i>Bacillus subtilis</i>
SJ1	13.3±1.5 56.9**	8.6±1.2 38.5*	9.5±0.8 49.0*	10.7±2.1 50.0**
SJ2	15.3±2.9 65.4**	13.5±2.5 60.3**	6.8±1.2 35.2*	10.1±1.8 47.2*
SJ3	14.1±1.4 60.3**	9.1±2.3 40.7*	6.7±1.4 34.8*	7.2±1.5 33.6*
SJ4	13.7±2.0 58.6**	7.0±1.0 31.1*	6.9±0.6 35.7*	8.3±1.5 38.9*
SJ5	13.3±3.0 56.8**	10.8±0.4 48.0*	7.4±1.3 38.1*	8.2±0.3 38.4*
SJ6	20.2±3.3 86.2***	18.0±2.2 80.4***	11.7±1.2 60.7**	10.9±1.4 51.3**
SJ7	15.3±1.7 65.3**	8.7±1.2 38.6*	11.5±2.0 59.7**	7.8±1.6 36.7*
SJ8	15.3±1.5 65.2**	15.7±1.6 70.1***	11.7±1.3 60.5**	11.5±1.2 53.9**
Tetracycline	23.4±1.4 100***	22.4±1.5 100***	19.3±1.3 100***	21.3±2.1 100***

Note: ^a: *** strong: inhibition ≥ 70%, ** moderate: inhibition 50-70%, * weak: inhibition < 50%, TM: no inhibition zone

Cultivation and extraction

Cultivation was carried out to increase endophytic fungi culture. A total of five bottles of endophytic fungi culture containing 300 mL, after an incubation period of 4 weeks, were extracted by partition in ethyl acetate solvent. The result of evaporation produces a concentrated extract of ethyl acetate, which weighs SJ1 6.5 g, SJ2 5.6 g, SJ3 5.8 g, SJ4 6.2 g, SJ5 5.9 g, SJ6 5.5 g, SJ7 5.3 g, and SJ8 5.0 g.

Antibacterial activity test

The activity of ethyl acetate extract of endophytic fungi from the bark of *S. jambos* was tested using the disc diffusion method by measuring the diameter of the growth inhibition zones at a concentration of 400 µg/disc. Endophytic fungi extract was tested on *Staphylococcus aureus*, *Bacillus subtilis*, *Escherichia coli*, and *Salmonella typhi* (Table 3). The extract of endophytic fungi with strong antibacterial activity was subjected to continuous isolation to obtain the active compound. Endophytic fungi SJ6 had strong activity against *S. aureus* and *S. typhi* bacteria, moderate activity against *E. coli* and *B. subtilis* bacteria. Endophytic fungi SJ8 had strong activity against *S. typhi* bacteria and moderate activity against three other bacteria. For endophytic fungi SJ1-SJ5 and SJ7, it has moderate and weak activity against all four bacteria. Endophytic fungi SJ6 has the strongest antibacterial activity, then proceed with molecular analysis and isolation of secondary metabolite.

Molecular analysis

Molecular characterization combined with morphological identification can help mold taxonomy in differentiation to the species level (Poursafar et al. 2018). The results of PCR products using a pair of universal primers of the ITS rDNA area varied by ±600bp, as shown in figure 2. Result of PCR amplification sequence assembly showed 543bp with CTTCCGTAGGTGAACCTGCGGAGGGATCATTACCG AGTTTACAACCTCCCAAACCCCTGTGAACATACCAA TTGTTGCCTCGGCGGATCAGCCCGCTCCCGGTAAA ACGGGACGGCCCGCCAGAGGACCCCTAAACTCTG TTTCTATATGTAACCTTCTGAGTAAACCATATAATA AATCAAAACTTTCAACAACGGATCTCTTGGTTCTG GCATCGATGAAGAACGCAGCAAAATGCGATAAGT AATGTGAATTGCAGAATTCAGTGAATCATCGAATC TTTGAACGCACATTGCGCCCGCCAGTATTCTGGCG GGCATGCCTGTTTCGAGCGTCATTTCAACCCTCAAG CCCAGCTTGTTGTTGGGACTCGCGAGTCAAATCGC GTTCCCCAAATTGATTGGCGGTACGTCGAGCTTC CATAGCGTAGTAGTAAAACCCCTCGTTACTGGTAAT CGTCGCGGCCACGCCGTTAAACCCCAACTTCTGAA TGTGACCTCGGATCAGGTAGGAATACCCGCTGAA CTTAAGCATATCAATAAGCC. GenBank accession numbers is OK426382.

The ITS sequence obtain from isolated will be compared with the sequence available in the GenBank database to analyze the phylogenetic affiliation. For samples of microorganisms, when using 16S rRNA markers, it is said to be identical (similar) at the species level if the "percentage identity" value is above 97.5% (Stackebrandt and Goebel 1994). The neighbor-Joining method is used to unrooted trees is focused only on relationships among taxa rather than evolutionary change. Outgroup rooted method assumes that are divergent for one or more of taxa, branch linking of ingroup and outgroup becomes the starting point to evolutionary of subsequent (Kinene et al. 2016). Sequence reading results from each of the forward and reverse primers of fungi endophyte SJ6 identified as *F. verticillioides*, shown in phylogenetic tree (Figure 3). The outgroup rooted in this phylogenetic tree is *Colletotrichum acutatum*. ITS sequence of endophytic fungal SJ6 isolated from *S. jambos* showed the highest level of similarity with percentage identity 99% is closest to *F. verticillioides*. Bootstrasp value in next brach shown >70, it can be concluded that the endophytic fungus SJ6 was identified as *F. verticillioides*.

Isolation of bioactive compound

The concentrated extract of ethyl acetate (1.0 g) was separated using gravity column chromatography with silica gel as a stationary phase and a mixture of *n*-hexane and ethyl acetate with increasing polarity (10:0–0:10) and a mixture of ethyl acetate with methane (9:1). The results of the separation were accommodated using vials every 10 mL, and as many as 55 vials were obtained. The eluate was then analyzed using thin-layer chromatography (TLC) with a mixed eluent of *n*-hexane and ethyl acetate (5:5). TLC with similar chromatogram patterns was combined into one fraction. Based on the results of the chromatogram pattern,

three fractions, namely F1–F3, were obtained. In the F2 fraction, a yellowish-white solid was formed, and after being purified with *n*-hexane, a pure compound (compound 1) with a weight of 18 mg was obtained. The antibacterial activity test of compound 1 showed good antibacterial activity against *S. aureus* and *S. typhi*, with an MIC value of 64 g/mL for both (Table 4).

Determination of chemical structure

The chemical structure of compound 1 was determined based on 1D and 2D NMR analysis. The ^1H -NMR spectrum of compound 1 (Figure 4) shows the presence of six proton signals, including two doublet signals, on the aromatic chemical shift, namely δ_{H} 7.57 (1H, d, $J = 8.5$ Hz) and 8.21 ppm (1H, d, $J = 8.5$ Hz). In the spectrum, each of these signals has an integration of two protons with the ortho-coupling constant ($J=8.5$ Hz). This indicates that compound 1 is a para-substituted aromatic compound, so it has two pairs of equivalent protons. Also, there are four signals in the chemical shift of δ_{H} 4.00–6.00 ppm, indicating the presence of four groups of sp^3 protons on oxygenated carbon and protons of methine. The four signals appear at δ_{H} 5.78 (1H, s), 5.33 (1H, d, $J = 2.0$ Hz), 4.16 (1H, m), and 4.01 ppm (2H, m). Based on the analysis of the ^1H -NMR spectrum, compound 1 is a para-substituted aromatic compound with nine protons bonded to a carbon atom. The solvent used in this measurement is CDCl_3 so that protons bound to heteroatoms do not appear in the spectrum.

The ^{13}C -NMR spectrum of compound 1 showed the presence of 11 signals. Four carbon signals appear in the aromatic region, namely at δ_{C} 164.4, 147.6, 126.8, and 123.8 ppm. A characteristic feature is the presence of two pairs of equivalent aromatic protons in the presence of two high-intensity signals at δ_{C} 126.8 and 123.8 ppm. It was also seen that the presence of aromatic oxyaryl carbon in the low field was C 164.4 ppm. Carbon in the lowest field appears at δ_{C} 175.6 ppm, which indicates that compound 1 has an ester group. Also, three oxygenated carbon signals appear in the δ_{C} 60.0–75.0 ppm area and one tertiary carbon signal at δ_{C} 55.6 ppm. The proton and carbon NMR spectra were confirmed by the data on the HMQC spectrum shown in Figure 5 and Table 5, namely the ^1H - ^{13}C correlation through one bond. The HMQC spectrum showed six correlations consisting of two correlations on the aromatic ring, three correlations on oxygenated ^1H - ^{13}C , and one correlation with proton methine.

The HMBC spectrum (Figure 6A) showed a ^1H - ^{13}C correlation through two or three bonds. The aromatic proton signal at δ_{H} 8.21 ppm is correlated through three bonds with its equivalent aromatic carbon (δ_{C} 123.8 ppm) and quaternary aromatic carbon (δ_{C} 147.6 ppm). The aromatic proton at δ_{H} 7.57 ppm is correlated through three bonds with its equivalent aromatic carbon (δ_{C} 126.6 ppm) and oxygenated carbon (73.5 ppm) and correlated through two bonds with quaternary aromatic carbon (δ_{C} 147.6 ppm). Furthermore, oxygenated methine protons at δ_{H} 5.33 ppm were correlated through three bonds with equivalent aromatic carbon (δ_{C} 126.6 ppm) and two bonds correlated with quaternary aromatic carbon (δ_{C} 147.6 ppm). The correlation indicates that the oxygenated methine group is

directly attached to the aromatic ring and is para-substituted with a hydroxyl group. In the spectrum, there is no correlation between the three protons (δ_{H} 4.01, 4.16, and 5.78 ppm). This may be due to the insufficient number of samples when measuring the spectrum. The proton hydroxyl signal does not appear on the spectrum because the pure compound is measured with the solvent CDCl_3 . The 1D and 2D NMR spectral data for compound 1 are shown in Table 5.




Figure 2. Electrophoresis of isolate SJ6. M (Marker 100 bp DNA ladder; 2 (sample isolate code SJ6 in ± 600 bp)




Figure 3. Phylogenetic position of endophytic fungal SJ6. The evolutionary history was inferred using the Neighbor-Joining method. The optimal tree with the sum of branch length = 3.00000000 is shown. The percentage of replicate trees in which the associated taxa clustered together in the bootstrap test (1000 replicates) are shown next to the branches. The tree is drawn to scale, with branch lengths in the same units as those of the evolutionary distances used to infer the phylogenetic tree. The analysis involved 20 nucleotide sequences. All positions containing gaps and missing data were eliminated. There were a total of 148 positions in the final dataset

Table 4. MIC value of pure compound and tetracycline against four test bacteria

Sample	MIC Value (µg/mL)			
	<i>Staphylococcus aureus</i>	<i>Salmonella typhi</i>	<i>Escherichia coli</i>	<i>Bacillus subtilis</i>
Compound 1	64	64	> 256	> 256
Tetracycline	1	1	1	1

Note: Inhibition zone >9mm indicates that the compound has antibacterial activity at the appropriate concentration.

Table 5. The NMR data of Compound 1

No. C	δ _C ppm	δ _H ppm (ΣH, multiplicity, J (Hz))	HMBC	COSY
2	175.6			
3	66.2	5.78 (1H, s)		
4	55.6	4.16 (1H, m)		5.33
5	64.0	4.01 (2H, m)		
6	73.5	5.33 (1H, d, J=2.0 Hz)	126.8; 147.6	4.16
1'	147.6			
2'	126.8	7.57 (1H, d, J=8.5 Hz)	147.6; 126.8; 73.5	8.21
3'	123.8	8.21 (1H, d, J=8.5 Hz)	147.6; 123.8	7.57
4'	164.4			
5'	123.8	8.21 (1H, d, J=8.5 Hz)	147.6; 123.8	7.57
6'	126.8	7.57 (1H, d, J=8.5 Hz)	147.6; 126.8; 73.5	8.21

Table 6. Diversity of endophytic fungi from the *Syzygium* genera

<i>Syzygium jambos</i>	<i>Syzygium jambos</i> (Roux et al. 2020)	<i>Syzygium aqueum</i> (Habisukan et al. 2021)	<i>Syzygium cordatum</i> (Marsberg et al. 2014)	<i>Syzygium cumini</i> (Hanin and Fitriyari 2019; Nurhaida et al. 2019)	<i>Syzygium samarangense</i> (Budiono et al. 2019)
<i>Spegazzinia tessarthra</i>	<i>Chrysoporthe deuterocubensis</i>	<i>Chaetomium</i>	<i>Mycosphaerella markii</i>	<i>Collectotrichum</i>	<i>Lasiodiplodia venezuelensis</i>
<i>Auxarthronopsis</i> sp.	<i>Microthia havanensis</i>	<i>Cochliobolus</i>	<i>M. vietnamensis</i>	<i>Phomopsis</i> sp.	
<i>Pectinotrichum</i> sp.	<i>Celoportha hauoliensis</i>	<i>Penicillium</i>		<i>Penicillium</i>	
<i>Phaeotrichosphaeria</i>	<i>Cel. hawaiiensis</i>	<i>Cylindrocladium</i>		<i>Pestalotiopsis</i>	
<i>Poaceascoma</i>	<i>Cel. paradisiaca</i>			<i>Fusarium</i>	
<i>Fusarium verticillioides</i>				<i>Aspergillus</i>	
<i>Thielaviopsis</i> sp.				<i>Myrothecium</i>	
<i>Nigrospora bambusae</i>				<i>Neofusicoccum parvum</i>	

The COSY spectrum in Figure 6B shows that the aromatic proton at δ_H 7.57 ppm correlates with the ¹H–¹H triple bonds with the aromatic proton at δ_H 8.21 ppm. Furthermore, there is a ¹H–¹H correlation of three hydrogenated protons of methine at δ_H 5.33 ppm with protons of methine at 4.16 ppm. This indicates that the two aromatic equivalent protons are in the ortho position. This reinforces the structural suggestion that compound 1 is a benzene ring directly bonded to the oxygenated methine carbon at the para position with a hydroxyl group.

Based on the spectral analysis of ¹H-NMR, ¹³C-NMR, HMQC, HMBC, and COSY, it can be explained that compound 1 has a para-substituted benzene ring between the hydroxyl group and the oxygenated methine group. This oxygenated methine group binds to the 3-hydroxydihydrofuran-2-one ring. The proton at δ_H 5.78 ppm appears singlet, probably due to the transposition with the proton at δ_H 4.16 ppm (Figure 7), is in the farthest geometric position so that cleavage does not occur. Thus, it

is proposed that the chemical structure of compound 1 is 3-hydroxy-4-(hydroxy(4-hydroxyphenyl)methyl)dihydrofuran-2-one, as shown in Figure 7.

Discussion

Eight endophytic fungi isolated from the stem bark of *S. jambos* have been found, namely *Spegazzinia*, *Auxarthronopsis*, *Pectinotrichum*, *Phaeotrichosphaeria*, *Poaceascoma*, *Fusarium*, *Thielaviopsis*, and *Nigrospora*. Seven endophytic fungi have not been found in other *Syzygium* genera, except *Fusarium* which is also found in *S. cumini* (Hanin and Fitriyari 2019). Based on the findings of endophytic fungi in this research and literature study from plants of the same genus, there are similarities in the diversity of endophytic fungi. The diversity of endophytic fungi from *S. jambos* and other species of the genus *Syzygium* is shown in Table 6.




Figure 4. The ^1H -NMR spectrum of compound 1




Figure 5. A Spectrum of ^{13}C -NMR (A) dan HMQC (B) of compound 1

The potential of *S. jambos* bark extract has been known to have antibacterial activity against *Staphylococcus aureus*, *Escherichia coli*, *Salmonella typhi*, *Bacillus subtilis* (Murugan et al. 2011; Begum et al. 2015; Wamba et al. 2018), and *Propionibacterium acnes* (Sharma et al. 2013b). It has been reported that chemical compounds from the bark extract of *S. jambos* contain polyphenols, anthraquinones, tannins, triterpenes, steroids, saponins (Wamba et al. 2018), pentacyclic triterpene, sitosterol, triterpene derivative 3-nor-2,3-fridelan (Haq et al. 2015),

jambosine alkaloids, pedunculagin ellagitannins, casuarinin, tellimagrandin I, strictinin, casuarictin, 2,3-HHDP glucose, and tellimagrandin II (Baliga et al. 2017). Some of these compounds are known to be antibacterial (Murugan et al. 2011). These antibacterial compounds are also produced by endophytic fungi, such as alkaloids, peptides, steroids, terpenoids, flavonoids, and quinines which can inhibit Gram-positive and Gram-negative bacteria (Manganyi and Ateba 2020).




Figure 6. Spectrum of Compound 1 (A. Analysis HMBC, B. Analysis COSY)

The results of the antibacterial test showed that the ethyl acetate fungal extract had antibacterial activity against the four sample bacteria. Endophytic fungi SJ6 (*Fusarium* sp.) had the strongest activity against *S. aureus* and *S. typhi*, while against *E. coli* and *B. subtilis* had moderate activity. It is known that *Fusarium* sp has the strongest antibacterial activity. Previous studies have reported that a secondary metabolite of *Fusarium* sp species, namely fusariumin D, has the potent activity against *S. aureus* (Wei and Wu 2020). The chemical compound of pyrrolo[1,2-a]pyrazine-1,4-dione, hexahydro-3-(2-methylpropyl) showed significant antibacterial activity against *B. subtilis*, *S. aureus*, *E. coli* (Putra and Karim 2020). Aliphatic carbonyl substituted compounds has antibacterial activity against *B. subtilis*, *E. coli*, *P. aeruginosa*, *S. aureus*, *S. mutans*, and *S. typhi* (Yuniati and Rollando 2018). Antibiotic Y was isolated from *F.*

avenaceum show very strong antibacterial activity against *B. subtilis* (Stępień et al. 2020).




Figure 7. The chemical structure of compound 1 is 3-hydroxy-4-(hydroxy(4-hydroxyphenyl)methyl) dihydrofuran-2-on

The biochemical diversity of secondary metabolites produced by *Fusarium* species, namely trichothecenes, fumonisins, zearalenone, enniatins and beauvericins, fusaric acid, fusarins, moniliformin (Stepień et al. 2020). The chemical compounds from the endophytic fungus *Fusarium* sp. have also been found whose structure was identified as fusaric acid (5-butylpicolinic acid, FA), alkaloid fungerin contains an imidazole, a-(N-formyl) carboxamide moiety, racemic imidazolidin-2,4-dione moiety, polyketide isoquinoline alkaloid fusarimine, fusaravenin, bisindole enantiomers, (±)-fusaspoid A, 7-desmethyloscorpinone and 7-desmethyl-6-methylbostrycoidin, apicidin derivatives: apicidin, apicidin A, apicidins B, C, D1, D2 and D3, apicidin E, apicidins F, J and K, apicidin G–I (Li et al. 2020). Wei and Wu (2020) isolated metabolite secondary, including γ -Pyrone derivative, α -pyrones, fupyrone A and B, anthraquinones and naphthoquinones, isocoumarins (karimunone B), macrolides (β -resorcylic macrolides, 7-hydroxy-14-de-Omethyl-lasiodiplodin), ethyl-3,5-dihydroxy-7-(6,8-dihydroxynonyl) benzoate, and 7-ethylbenzoate-7-heptanoic acid, γ -methylidene-spirobutanolide core, fusaspirols A-D, fusarielins. There are several groups of compounds produced by endophytic fungi of the *Fusarium* genus that have antibacterial abilities, including from the sesquiterpene group, such as 3 β -hydroxy- β -acorenol and fusariumins D, the terpene group (fusariumins C), naphthalenone derivative, pyrone derivative, namely pyrano[2,3-g]indole moieties and amoenamide C. There are also group of compounds derived from naphthoquinone, 6-hydroxy-astropaquinone B dan astropaquinone D. Other compounds, such as sesquiterpenoids derivative, emericellins A-B, phenolic bisabolane sesquiterpenoids, tremulane sesquiterpene, furan derivatives, poliketides also have antibacterial ability (Zheng et al. 2021).

Some endophytic fungi also produce the same bioactive compounds as their host, such as paclitaxel (taxol), baccatin III, podophyllotoxin, hypericin, camptothecin, vinblastine, huperzine a, diosgenin, a-irone, toosendanin produced by fungi isolated from *Taxus* sp (Zhao et al. 2010). Based on the Dictionary of Natural Products database on 9 Oktober 2021, 3-hydroxy-4-(hydroxy(4-hydroxyphenyl)methyl)dihydrofuran-2-on were isolated from the *Fusarium verticillioides* are new phenolic compounds. Phenolic compounds are the best source of antibiotic. Several phenolic compounds were antibacterial activity against *S. epidermidis* and *P. aeruginosa*, such as tannic acid, epigallocatechin gallate, etc. (Mandal et al. 2017). Phenolic compounds are known antioxidants and other bioactive agents that have benefits for human health and many diseases (Tungmunthum et al. 2018). However, based on the literature, there are no compounds similar to those found in this study. The compound produced by the endophytic fungus *F. verticillioides* is different from the compounds found in its host *S. jambos*, although it is still in the phenolic compound group.

ACKNOWLEDGEMENTS

The authors thank to the DRPM Kemenristek Republik Indonesia, which provided research funding through Hibah Disertasi Doktor 2021, with contract no. 0166.05/UN9/SB3.LP2M.PT/2021

REFERENCES

- Abdel-Azeem AM, Abdel-Azeem MA, Khalil WF. 2019. Endophytic Fungi as a New Source of Antirheumatoid Metabolites. In Bioactive Food as Dietary Interventions for Arthritis and Related Inflammatory Diseases. Academic Press. DOI: 10.1016/B978-0-12-813820-5.00021-0.
- Álvarez E, Llano GA, Loke JB, Chacon MI. 2012. Characterization of Thielaviopsis paradoxa Isolates from Oil Palms in Colombia, Ecuador and Brazil. J Phytopathol 160 (11-12): 690-700. DOI: 10.1111/jph.12012.
- Anand U, Jacobo-Herrera N, Altemimi A, Lakhssassi N. 2019. A comprehensive review on medicinal plants as antimicrobial therapeutics: potential avenues of biocompatible drug discovery. Metabolites 9 (11): 1-13. DOI: 10.3390/metabo9110258.
- Athikomkulchai S, Lipipun V, Leelawittayanont T, Khanboon A, Ruangrunsi N. 2008. Anti-Herpes simplex virus activity of *Syzygium jambos*. J Health Res 22 (1): 49-51.
- Baliga MS, Pai KSR, Saldanha E, Ratnu VS, Priya R, Adnan M, Naik TS. 2017. Rose apple (*Syzygium jambos* (L.) Alston). Fruit Vegetable Phytochem: Chem Human Health: 2nd Edition 2: 1235-1241. DOI: 10.1002/9781119158042.ch65.
- Begum MM, Haque M, Reyad-ul-Ferdous M, Hasan M, Tarek H, Noor-A-alam M, Hossain MB, Uddin MN, Begum T, Hasan CM. 2015. Screening of antioxidant and antimicrobial properties of the *Syzygium jambos* L. Am J BioSci 3 (2-1): 23-26. DOI: 10.11648/j.ajbio.s.2015030201.15.
- Bonfanti G, Bitencourt P, Bona K, Silva P, Jantsch L, Pigatto A, Boligon A, Athayde M, Gonçalves T, Moretto M. 2013. *Syzygium jambos* and *Solanum guaraniticum* show similar antioxidant properties but induce different enzymatic activities in the brain of rats. Molecules 18 (8): 9179-9194. DOI: 10.3390/molecules18089179.
- Bonfanti G, De Bona KS, de Lucca L, Jantsch L, Pigatto AS, Boligon AA, Athayde ML, Moretto MB, Gonçalves TD. 2014. Delta-ALA-D inhibitory potential and protective action of *Syzygium jambos* and *Solanum guaraniticum* leaf extracts on oxidatively stressed erythrocytes. Redox Rep 19 (5): 206-213. DOI: 10.1179/1351000214Y.0000000092.
- Budiono, Elfita, Muharni, Yohandini H, Widjajanti H. 2019. Antioxidant activity of *Syzygium samarangense* L. and their endophytic fungi. Molekul 14 (1): 48-55. DOI: 10.20884/1.jm.2019.14.1.503.
- Cole GT. 1974. Conidiophore and conidium ontogeny in *Spiegazzinia tessartha*. Can J Bot 52 (6): 1259-1264. DOI: 10.1139/b74-163.
- Devakumar J, Sudha SS. 2017. In vitro phytochemical, antioxidant and cytotoxic evaluation of *Syzygium jambos* L. (Alston). J Pharm Res 11 (3): 235-238.
- Ding Z, Tao T, Wang L, Zhao Y, Huang H, Zhang D, Liu M, Wang Z, Han J. 2019. Bioprospecting of novel and bioactive metabolites from endophytic fungi isolated from rubber tree *Ficus elastica* leaves. J Microbiol Biotechnol 29 (5): 731-738. DOI: 10.4014/jmb.1901.01015.
- Elfita, Mardiyanto, Fitrya, Eka LJ, Julinar, Widjajanti H, Muharni. 2019. Antibacterial activity of *Cordylone fruticosa* leaf extracts and its endophytic fungi extracts. Biodiversitas 20 (12): 3804-3812. DOI: 10.13057/biodiv/d201245.
- Fadhillah, Elfita, Muharni, Yohandini H, Widjajanti H. 2019. Chemical compound isolated from antioxidant active extract of endophytic fungus *Cladosporium tenuissimum* in *Swietenia mahagoni* leaf stalks. Biodiversitas 20 (9): 2645-2650. DOI: 10.13057/biodiv/d200929.

- Felsenstein J. 1985. Confidence limits on phylogenies: an approach using the bootstrap. *Evolution* 39 (4): 783-791. DOI: 10.1111/j.1558-5646.1985.tb00420.x.
- Fors RO, Patreze CM, Barbera RLL, Cameiro MAC, Saggin OJ. 2020. Dark septate endophytic fungi associated with sugarcane plants cultivated in Sao Paulo, Brazil. *Diversity* 12 (9): 1-21. DOI: 10.3390/D12090351.
- Ghareeb MA, Saad AM, Abdel-Aleem AAH, Abdel-Aziz MS, Hamed MM, Hadad AH. 2016. Antioxidant, antimicrobial, cytotoxic activities and biosynthesis of silver and gold nanoparticles using *Syzygium jambos* leaves growing in Egypt. *Der Pharma Chemica* 8 (18): 107-116.
- Habisukan UH, Elfita, Widjajanti H, Setiawan A, Kurniawati AR. 2021. Diversity of Endophytic Fungi in *Syzygium aqueum*. *Biodiversitas* 22 (3): 1129-1137. DOI: 10.13057/biodiv/d220307.
- Hanin NA, Fitriyanti PD. 2019. Identification of endophytic fungi from fruits and seeds of jambolana (*Syzygium cumini* L.) Skeels. *IOP Conf Ser: Earth Environ Sci* 276: 012060. DOI: 10.1088/1755-1315/276/1/012060.
- Haq M, Begum MM, Hasan M, Rahman AFMT, Hussain MI, Rahman MM, Ali MH, Islam MA, Sultan MZ, Reyad-ul-Ferdous M, Hasan CM. 2015. Investigation of the medicinal potentials of *Syzygium jambos* (L.) extract and characterization of the isolated compounds. *Am J BioSci* 3 (2): 12-18. DOI: 10.11648/j.ajbio.s.2015030201.13.
- Katoch M, Pull S. 2017. Endophytic fungi associated with *Monarda citriodora*, an aromatic and medicinal plant and their biocontrol potential. *Pharm Biol* 55 (1): 1528-1535. DOI: 10.1080/13880209.2017.1309054.
- Katz L, Baltz RH. 2016. Natural product discovery: past, present, and future. *J Ind Microbiol Biotechnol* 43 (2-3): 155-176. DOI: 10.1007/s10295-015-1723-5.
- Khalil AMA, Hassan SED, Alsharif SM, Eid AM, Eweis EED, Azab E, Gobouri AA, Elkesh A, Fouda A. 2021. Isolation and characterization of fungal endophytes isolated from medicinal plant *Ephedra pachyclada* as plant growth-promoting. *Biomolecules* 11 (2): 1-18. DOI: 10.3390/biom11020140.
- Kinene T, Wainaina J, Maina S, Boykin LM. 2016. Rooting Trees, Methods for. *Encyclopedia Evol Biol* 3: 489-493. DOI: 10.1016/B978-0-12-800049-6.00215-8.
- Kuswiyatari ND, Kurniawati AR, Alami, NH, Zulaika E, Shovitri M, Oh KM, Puspaningsih NNT, Ni'Matuzahroh. 2019. Plastic degradation by *Coriopsis byrsina*, an identified white-rot, soil-borne mangrove fungal isolate from Surabaya, East Java, Indonesia. *Biodiversitas* 20 (3): 867-871. DOI: 10.13057/biodiv/d200334.
- Li M, Yu R, Bai X, Wang H, Zhang H. 2020. *Fusarium*: A treasure trove of bioactive secondary metabolites. *Nat Prod Rep* 37 (12): 1568-1588. DOI: 10.1039/d0np00038h.
- Lim TK. 2012. *Syzygium jambos*. *Edible Medicinal and Non Medicinal Plants*. Springer Netherlands. DOI: 10.1007/978-94-007-2534-8_101.
- Luo ZL, Bahkali AH, Liu XY, Phookamsak R, Zhao YC, Zhou DQ, Su HY, Hyde KD. 2016. *Poaceascoma aquaticum* sp. nov. (Lentitheciaceae), a New Species from Submerged Bamboo in Freshwater. *Phytotaxa* 253 (1): 71-80. DOI: 10.11646/phytotaxa.253.1.5.
- Majumdar N, Mandal NC. 2018. Growth and sporulation physiology of postharvest pathogen *Thielaviopsis paradoxa* (De Seynes.) Höhn. *Intl J Curr Microbiol Appl Sci* 7 (7): 537-544. DOI: 10.20546/ijcmas.2018.707.066.
- Mandal SM, Dias RO, Franco OL. 2017. Phenolic compounds in antimicrobial therapy. *J Med Food* 20 (10): 1031-1038. DOI: 10.1089/jmf.2017.0017.
- Manganyi MC, Ateba CN. 2020. Untapped potentials of endophytic fungi: a review of novel bioactive compounds with biological applications. *Microorganisms* 8 (12): 1-25. DOI: 10.3390/microorganisms8121934.
- Marsberg A, Slippers B, Wingfield MJ, Gryzenhout M. 2014. Endophyte isolations from *Syzygium cordatum* and a *Eucalyptus* clone (Myrtaceae) reveal new host and geographical reports for the Mycosphaerellaceae and Teratosphaeriaceae. *Australas Plant Pathol* 43 (5): 503-512. DOI: 10.1007/s13313-014-0290-y.
- Martinez-Klimova E, Rodríguez-Peña K, Sánchez S. 2017. Endophytes as sources of antibiotics. *Biochem Pharmacol* 134: 1-17. DOI: 10.1016/j.bcp.2016.10.010.
- Mena-Portales J, Cantillo-Pérez T Minter DW. 2017. A new species of the Conidial fungal genus *Spegazzinia* (Pleosporales, Didymosphaeriaceae) collected on sugarcane in Cuba. *Phytotaxa* 331 (2): 295-298. DOI: 10.11646/phytotaxa.331.2.14.
- Mohanty S, Cock IE. 2010. Bioactivity of *Syzygium jambos* methanolic extracts: Antibacterial activity and toxicity. *Pharmacognosy Res* 2 (1): 4-9. DOI: 10.4103/0974-8490.60577.
- Morton JF. 1987. Fruits of warm climates. JF Morton.
- Murugan S, Devi PU, Kannika Parameswari N, Mani KR. 2011. Antimicrobial activity of *Syzygium jambos* against selected human pathogens. *Intl J Pharm Pharm Sci* 3 (2): 44-47.
- Nei M, Kumar S. 2000. *Molecular Evolution and Phylogenetics*. Oxford University Press, Inc. DOI: 10.1002/wilm.10463.
- Nisa S, Khan N, Shah W, Sabir M, Khan W, Bibi Y, Jahangir M, Haq IU, Alam S, Qayyum A. 2020. Identification and bioactivities of two endophytic fungi *Fusarium fujikuroi* and *Aspergillus tubingensis* from foliar parts of *Debregeasia salicifolia*. *Arab J Sci Eng* 45 (6): 4477-4487. DOI: 10.1007/s13369-020-04454-1.
- Nurhaida, Yenn TW, Ibrahim D. 2019. Endophytic fungi from *Syzygium cumini* (L.) Skeels leaves and its potential as antimicrobial agents. *IOP Conf Ser: Earth Environ Sci* 151: 01006. DOI: 10.1088/1755-1315/364/1/012023.
- Phookamsak R, Manamgod, DS, Li WJ, Dai DQ, Singtripop C, Hyde KD. 2015. *Poaceascoma helicoides* Gen et sp. nov, a New Genus with *Scolecospores* in Lentitheciaceae. *Cryptogam Mycol* 36 (2): 225-236. DOI: 10.7872/crym/v36.iss2.2015.225.
- Poursafar A, Ghosta Y, Orina AS, Gannibal PB, Javan-Nikkhah M, Lawrence DP. 2018. Taxonomic study on *Alternaria* sections infectoriae and *Pseudoalteraria* associated with black (Sooty) head mold of wheat and barley in Iran. *Mycol Prog* 17 (3): 343-356. DOI: 10.1007/s11557-017-1358-1.
- Putra MY, Karim F. 2020. Antibacterial and antioxidant activity-guided isolation studies on *Fusarium* sp. associated with the ascidian *Botryllus schlosseri*. *AIP Conf Proc* 2243. DOI: 10.1063/5.0001297.
- Rai M, Gade A, Rathod D, Dar M, Varma A. 2012. Review: Mycoendophytes in medicinal plants: Diversity and bioactivities. *Nusantara Biosci* 4: 86-96. DOI: 10.13057/nusbiosci/n040207.
- Rajkumari J, Borkotoky S, Murali A, Busi S. 2018. Anti-quorum sensing activity of *Syzygium jambos* (L.) Alston against *Pseudomonas aeruginosa* PAO1 and identification of its bioactive components. *S Afr J Bot* 118: 151-157. DOI: 10.1016/j.sajb.2018.07.004.
- Roux J, Nkuekam GK, Marincowitz S, van der Merwe NA, Uchida J, Wingfield MJ, Chen SF. 2020. Cryphonectriaceae associated with rust-infected *Syzygium jambos* in Hawaii. *Mycoskeys* 76: 49-79. DOI: 10.3897/MYCOKEYS.76.58406.
- Saitou N, Nei M. 1987. The neighbor-joining method: a new method for reconstructing phylogenetic trees. *Mol Biol Evol* 4 (4): 406-425. DOI: 10.1093/oxfordjournals.molbev.a040454.
- Sharma R, Gräser Y, Singh SK. 2013a. Auxarthronopsis, a new genus of onygenales isolated from the Vicinity of Bandhavgarh National Park, India. *IMA Fungus* 4 (1): 89-102. DOI: 10.5598/imafungus.2013.04.01.09.
- Sharma R, Kishore N, Hussein A, Lall N. 2013b. Antibacterial and anti-inflammatory effects of *Syzygium jambos* L. (Alston) and isolated compounds on *Acne vulgaris*. *BMC Complement Altern Med* 13 (292): 1-10. DOI: 10.1186/1472-6882-13-292.
- Sivanesan A. 1983. Studies on Ascomycetes. *Trans Brit Mycol Soc* 81 (2): 313-332. DOI: 10.1016/s0007-1536(83)80084-9.
- Sobeh M, Esmat A, Petruk G, Abdelfattah MAO, Dmirieh M, Monti DM, Abdel-Naim AB, Wink M. 2018. Phenolic compounds from *Syzygium jambos* (Myrtaceae) exhibit distinct antioxidant and hepatoprotective activities in vivo. *J Funct Foods* 41: 223-231. DOI: 10.1016/j.jff.2017.12.055.
- Stackebrandt E, Goebel BM. 1994. Taxonomic note: A place for DNA-DNA reassociation and 16s rRNA sequence analysis in the present species definition in bacteriology. *Intl J Syst Evol Microbiol* 44 (4): 846-849. DOI: 10.1016/S0140-6736(01)43317-4.
- Stępień Ł, Lalak-Kańczugowska J, Witaszak N, Urbaniak M. 2020. *Fusarium* Secondary Metabolism Biosynthetic Pathways: So Close but So Far Away. *Co-Evol Second Metabolites* 211-247. DOI: 10.1007/978-3-319-96397-6_28.
- Sudha V, Govindaraj R, Baskar K, Al-Dhabi A, Duraipandian V. 2016. Biological properties of endophytic fungi. *Braz Arch Biol Technol* 59: 16150436. DOI: 10.1590/1678-4324-2016150436.
- Tamura K, Stecher G, Kumar S. 2021. MEGA11: Molecular Evolutionary Genetics Analysis Version 11. *Mol Biol Evol* 38 (7): 3022-3027. DOI: 10.1093/molbev/msab120.
- Tungmunthitum D, Thongboonyou A, Pholboon A, Yangsabai A. 2018. Flavonoids and other phenolic compounds from medicinal plants for

- pharmaceutical and medical aspects: an overview. *Medicines* 5 (93): 1-16. DOI: 10.3390/medicines5030093.
- Varsavsky E, Orr GF. 1971. A New Genus of the Gymnoascaceae. *Mycopathologia et Mycologia Applicata* 43 (2): 229-234. DOI: 10.1007/BF02051726.
- Walsh TH, Hayden RT, Larone DH. 2018. *Larone's Medically Important Fungi: A Guide to Identification* (6th Editio). ASM Press, Washington DC. DOI: 10.1128/9781555819880.
- Wamba BEN, Nayim P, Mbaveng AT, Voukeng IK, Dzotam JK, Ngalani OJT, Kuete V. 2018. *Syzygium jambos* displayed antibacterial and antibiotic-modulating activities against resistant phenotypes. *Evidence-Based Complement Altern Med* 5124735. DOI: 10.1155/2018/5124735.
- Wan JP. 2017. The Applications of Acetone and Ethyl Acetate. *Solvents as Reagents in Organic Synthesis: Reactions and Applications*. DOI: 10.1002/9783527805624.ch5.
- Wang M, Liu F, Crous PW, Cai L. 2017. Phylogenetic reassessment of *Nigrospora*: ubiquitous endophytes, plant and human pathogens. *Persoonia: Mol Phyl Evol Fungi* 39: 118-142. DOI: 10.3767/persoonia.2017.39.06.
- Watanabe T. 2010. Pictorial atlas of soil and seed fungi: morfologies of cultural fungi and key to species. In *Pictorial Atlas of Soil and Seed Fungi* (Third Edit). CRC Press, Taylor & Francis Group. DOI: 10.1201/ebk1439804193.
- Wei J, Wu B. 2020. Chemistry and Bioactivities of Secondary Metabolites from The Genus *Fusarium*. *Fitoterapia* 146: 104638. DOI: 10.1016/j.fitote.2020.104638.
- White TJ, Bruns T, Lee S, Taylor J. 1990. Amplification and direct sequencing of fungal ribosomal Rna genes for phylogenetics. *PCR Protocols* 2015: 315-322. DOI: 10.1016/b978-0-12-372180-8.50042-1.
- Yuniati Y, Rollando R. 2018. Isolation of antibacterial compounds from endophyte fungal of *Fusarium* sp. in *Phyllanthus niruri* linn. leaves. *J Pharm Sci Res* 10 (2): 260-264.
- Zhang ZF, Zhou X, Liu F, Cai L, Liu SJ, Liu XZ. 2017. Culturable mycobiota from karst caves in China II, with Descriptions of 20 New Species. *Persoonia: Mol Phyl Evol Fungi* 39 (1): 1-31. DOI: 10.1007/s13225-020-00453-7.
- Zhao J, Zhou L, Wang J, Shan T, Zhong L, Liu X, Gao X. 2010. Endophytic fungi for producing bioactive compounds originally from their host plants. *Curr Res, Technol Educ Top Appl Microbiol Microb Biotechnol* 2015: 567-576.
- Zheng R, Li S, Zhang X, Zhao C. 2021. Biological activities of some new secondary metabolites isolated from endophytic fungi: a review study. *Intl J Mol Sci* 22 (959): 1-80. DOI: 10.3390/ijms22020959.



Ads by Google

[Stop seeing this ad](#)[Why this ad? ⓘ](#)Biodiversitas

COUNTRY

Indonesia

Universities and research
institutions in IndonesiaSUBJECT AREA AND
CATEGORYAgricultural and
Biological Sciences
Animal Science and
Zoology
Plant ScienceBiochemistry, Genetics
and Molecular Biology
Molecular Biology

PUBLISHER

Biology department,
Sebelas Maret
University SurakartaUniversitas Negeri
Sebelas Maret in
Scimago
Institutions
Rankings

H-INDEX

19

PUBLICATION TYPE

Journals

ISSN

1412033X, 20854722

COVERAGE

2014-2021

INFORMATION

[Homepage](#)[How to publish in this
journal](#)[grahameagleton@gmail.
com](mailto:grahameagleton@gmail.com)

Ads by Google

[Stop seeing this ad](#)

Ads by Google

[Stop seeing this ad](#)[Why this ad? ⓘ](#)

SCOPE

"Biodiversitas, Journal of Biological Diversity" or Biodiversitas encourages submission of manuscripts dealing with all biodiversity aspects of plants, animals and microbes at the level of gene, species, and ecosystem.

 Join the conversation about this journal

← Ads by Google

Stop seeing this ad

Why this ad? ⓘ

↶ Quartiles
—
☰

← Ads by Google

Stop seeing this ad

Why this ad? ⓘ

↵
←

Ads by Google

Stop seeing this ad

Why this ad? ⓘ

/ /

similarity

PeerJ
USA

similarity

Biologia (Poland)
DEU

similarity

Tropical Life Sci
MYS

similar





Ads by Google

Stop seeing this ad

Why this ad? ⓘ

1.6k

800




← <https://www.scimagojr.com>



SCImago Graphica

Explore, visually communicate and make sense of data with our [new data visualization tool](#).



Ads by Google

Stop seeing this ad

Why this ad? ⓘ

Metrics based on Scopus® data as of April 2022

W

Widi Sunaryo 5 months ago

Good Journal, fast response, good quality, good reviewing process, divers in diversity.....

reply



Melanie Ortiz 5 months ago

SCImago Team

Dear Widi, thanks for your participation! Best Regards, SCImago Team

▼


B

BOURAADA KHALID 11 months ago

Ads by Google

Stop seeing this ad

Why this ad? ⓘ



(<https://smujo.id/biodiv/issue/view/324>)

Vol. 23 No. 1 (2022)

Full Issue

Front Cover (<https://smujo.id/biodiv/issue/view/324/184>)

Articles

Assessing the sustainability status of mangrove forest ecosystem management by coastal community in Jaring Halus Village, North Sumatra, Indonesia (<https://smujo.id/biodiv/article/view/7409>)

EDY SAHPUTRA, R. HAMDANI HARAHAH, HESTI WAHYUNINGSIH, BUDI UTOMO

PDF (<https://smujo.id/biodiv/article/view/7409/5402>)

Biodiversity of *Escherichia coli* bacterial resistance to multidrug isolated on the Dumai coast of Indonesia (<https://smujo.id/biodiv/article/view/8854>)

FELI FELIATRA, MARDALISA MARDALISA, IRWAN EFFENDI, ADELINA ADELINA, VANESA ADIBA FELIATRA

PDF (<https://smujo.id/biodiv/article/view/8854/5395>)

Universal primer design for crustacean and bivalve-mollusc authenticity based on cytochrome-b gene (<https://smujo.id/biodiv/article/view/9545>)

DWIYITNO DWIYITNO, STEFAN HOFFMAN, KOEN PARMENTIER, CHRIS VAN KEER

PDF (<https://smujo.id/biodiv/article/view/9545/5396>)

Host range studies of *Fusarium oxysporum*, causal agent of seedling wilt disease of *Acacia mangium* (<https://smujo.id/biodiv/article/view/9450>)

SOLEHA SOLEHA, AHMAD MUSLIM, SUWANDI SUWANDI, SABARUDDIN KADIR, RAHMAT PRATAMA

PDF (<https://smujo.id/biodiv/article/view/9450/5397>)

Diversity, distribution and ecology of *Davallia* in Sumatra (Indonesia) and the nearby small islands (<https://smujo.id/biodiv/article/view/9868>)

MILDAWATI, SOBIR, SULISTIJORINI, TATIK CHIKMAWATI

PDF (<https://smujo.id/biodiv/article/view/9868/5394>)

Habitat preferences of wild orchids in Bantimurung Bulusaraung National Park to model their suitable habitat in South Sulawesi, Indonesia (<https://smujo.id/biodiv/article/view/9574>)

EKA MARTHA DELLA RAHAYU, SAFRAN YUSRI

PDF (<https://smujo.id/biodiv/article/view/9574/5403>)

Positive reinforcement conditioning as Sumatran tiger's (*Panthera tigris sumatrae*) social enrichment at Tambling Wildlife Nature Conservation Rescue Centre, Lampung, Indonesia (<https://smujo.id/biodiv/article/view/9652>)

GENOVEVA KIRANAPUTRI, LUTHFIRALDA SJAHFIRDI, LIGAYA ITA TUMBELAKA, ARI YANA, SADMOKO KUSUMO PRIYANTO, LESTY YULIETA ANGGARSARI, MARIZAL

PDF (<https://smujo.id/biodiv/article/view/9652/5404>)

Short Communication: Morphological diversity and the addition of golden snail protein: Its effect on flavonoid content on *Echinacea purpurea* (<https://smujo.id/biodiv/article/view/9774>)

WINA CHANDRA FERDYANA, YULI WIDIYASTUTI, BAMBANG PUJIASMANTO, AMALIA TETRANI SAKYA, AHMAD YUNUS

PDF (<https://smujo.id/biodiv/article/view/9774/5398>)

Monthly forage value of various plant organs of *Brocchia cinerea* (Vis.) from a Moroccan arid region (<https://smujo.id/biodiv/article/view/9076>)

NISRINE CHLIF, MOHAMMED DIOURI, MOHAMMED EL OMARI, AMAR BENTAYEB

PDF (<https://smujo.id/biodiv/article/view/9076/5405>)

Grasshopper diversity in several agricultural areas and savannas in Dompu, Sumbawa Island, Indonesia (<https://smujo.id/biodiv/article/view/9845>)

AMIN SETYO LEKSONO, BAGYO YANUWIADI, AINUL KHOTIMAH, ANISA ZAIRINA

PDF (<https://smujo.id/biodiv/article/view/9845/5406>)

Short Communication: Molecular study of bacteria isolated from meat and chicken frozen from Misan Governorate market in Iraq (<https://smujo.id/biodiv/article/view/9732>)

DUAA ALI HUSAIN, ZAHID SA'DOON AZIZ

PDF (<https://smujo.id/biodiv/article/view/9732/5407>)

Heteropneustes fuscus (Siluriformes: Heteropneustidae), a new catfish species from Kerala, India (<https://smujo.id/biodiv/article/view/9564>)

MATHEWS PLAMOOTTIL

PDF (<https://smujo.id/biodiv/article/view/9564/5417>)

Effects of harvest period, storage, and genotype on postharvest physiological deterioration responses in cassava (<https://smujo.id/biodiv/article/view/9663>)

RIKA SRI RAHMAWATI, NURUL KHUMAIDA, SINTHO WAHYUNING ARDIE, DEWI SUKMA, SUDARSONO

PDF (<https://smujo.id/biodiv/article/view/9663/5408>)

Novel microbial transformation of *Andrographis paniculata* by *Aspergillus oryzae* K1A (<https://smujo.id/biodiv/article/view/9740>)

RINI HANDAYANI, ACHMAD DINOTO, MARIA BINTANG

PDF (<https://smujo.id/biodiv/article/view/9740/5409>)

Diversity of phytoplankton in the whiteleg (*Litopenaeus vannamei*) shrimp ponds in the south coastal area of Pangandaran, Indonesia (<https://smujo.id/biodiv/article/view/9606>)

MUSTIKA PALUPI, REN FITRIADI, RUDI WIJAYA, PURWO RAHARJO, RISA NURWAHYUNI

PDF (<https://smujo.id/biodiv/article/view/9606/5418>)

Microplastic contamination in Indonesian anchovies from fourteen locations (<https://smujo.id/biodiv/article/view/9371>)

ENDAR W NINGRUM, MUFTI P PATRIA

PDF (<https://smujo.id/biodiv/article/view/9371/5419>)

An antibacterial compound purified from a tropical coastal plant, *Diospyros maritima* (<https://smujo.id/biodiv/article/view/10020>)

ALIM ISNANSETYO, DESY PUTRI HANDAYANI, INDAH ISTIQOMAH, ADITYA ARIF, TAKUSHI KANEKO

PDF (<https://smujo.id/biodiv/article/view/10020/5410>)

Exterior characteristics and body measurements of Bligon goat on the different agro-ecological zones in Bantul District, Yogyakarta, Indonesia (<https://smujo.id/biodiv/article/view/9838>)

RENY DWI RAHMAWATI, BAYU ANDRI ATMOKO, I GEDE SUPARTA BUDISATRIA, NONO NGADIYONO, PANJONO

PDF (<https://smujo.id/biodiv/article/view/9838/5420>)

Population dynamics and feeding habit of *Oreochromis niloticus* and *O. mossambicus* in Siombak Tropical Coastal Lake, North Sumatra, Indonesia (<https://smujo.id/biodiv/article/view/9577>)

AHMAD MUHTADI, MUHAMMAD NUR, HUSAIN LATUCONSINA, THOMAS HIDAYAT

PDF (<https://smujo.id/biodiv/article/view/9577/5431>)

Combined use of *Spirulina platensis* and *Saccharomyces cerevisiae*:
Implication on growth, blood profile and intestinal morphology and bacteria
of the Indonesian crossbred chickens
(<https://smujo.id/biodiv/article/view/9860>)

SUGIHARTO SUGIHARTO, BALUH MEDYABRATA ATMAJA, ENDANG WIDIASTUTI,
HADIYANTO HADIYANTO

PDF (<https://smujo.id/biodiv/article/view/9860/5421>)

Improving the larvae acceptance rate and morphometric characteristics of
queen *Apis cerana javana* by grafting larvae at different ages
(<https://smujo.id/biodiv/article/view/9533>)

USTADI, NUR IKHSAN, MOCH. JUNUS, SUYADI

PDF (<https://smujo.id/biodiv/article/view/9533/5432>)

Toxicity test of rose periwinkle (*Catharanthus roseus*) leaves endophytic
bacteria using Brine Shrimp Lethality Test (BSLT) method
(<https://smujo.id/biodiv/article/view/9820>)

FAUZIAH FAUZIAH, MAULINASARI, ESSY HARNELLY, YULIA SARI ISMAIL, LENNI FITRI

PDF (<https://smujo.id/biodiv/article/view/9820/5422>)

Diversity of *Gyrinops versteegii* from several agarwood plantation on
Lombok Island (Indonesia) as raw material of *Gyrinops* tea
(<https://smujo.id/biodiv/article/view/9870>)

I GDE ADI SURYAWAN WANGIYANA, SUPRIADI SUPRIADI, ALUH NIKMATULLAH, SUNARPI
SUNARPI, AGUNG AYU HARI TRIANDINI

PDF (<https://smujo.id/biodiv/article/view/9870/5456>)

Fish diversity and heavy metal accumulation of Pb, Cu and Zn after Mount
Sinabung Eruption in Benuken River, North Sumatra, Indonesia
(<https://smujo.id/biodiv/article/view/8739>)

MELDA AGNES PEBRIANI, TERNALA ALEXANDER BARUS, SYAFRUDDIN ILYAS

PDF (<https://smujo.id/biodiv/article/view/8739/5434>)

Identification of bioactive compounds of banana corm (*Musa paradisiaca*)
using GC-MS and its inhibitory effect against pathogenic bacteria
(<https://smujo.id/biodiv/article/view/9699>)

NURUL FAJRIH, KOMANG GEDE WIRYAWAN, SUMIATI, SURAYA KAFFI SYAHPURA, WIWIN WINARSIH

PDF (<https://smujo.id/biodiv/article/view/9699/5423>)

Genetic diversity analysis of Puan Kalianda kopyor coconuts (*Cocos nucifera*) from South Lampung, Indonesia based on SSR markers (<https://smujo.id/biodiv/article/view/9751>)

MEGAYANI SRI RAHAYU, ASEP SETIAWAN, ISMAIL MASKROMO, AGUS PURWITO, SUDARSONO

PDF (<https://smujo.id/biodiv/article/view/9751/5424>)

Diversity and abundance of plankton in different habitat zonation of Papan River, Lake Kenyir, Malaysia (<https://smujo.id/biodiv/article/view/9790>)

ALIFFIKRI RAMLEE, HIDAYU SUHAIMI, NADIAH W. RASDI

PDF (<https://smujo.id/biodiv/article/view/9790/5433>)

Genetically continuous populations of Striped Snakehead (*Channa striata*) in the Cingcingguling River fragmented by Sempor Reservoir, Central Java, Indonesia (<https://smujo.id/biodiv/article/view/10078>)

NUNING SETYANINGRUM, W. LESTARI, KRISMONO, AGUS NURYANTO

PDF (<https://smujo.id/biodiv/article/view/10078/5425>)

Population characteristics and habitat suitability of Khao Yai National Park, Thailand for Asian elephant and five ungulate species (<https://smujo.id/biodiv/article/view/10084>)

MANANYA PLA-ARD, NORASET KHIOESREE, BENCHARONG SUNKALAK, ANUTTARA NATHALANG, WARISARA THOMAS, SUWIMON UTHAIRATSAMEE, PAANWARIS PAANSRI, YUWALAK CHANACHAI, RONGLARP SUKMASUANG

PDF (<https://smujo.id/biodiv/article/view/10084/5426>)

Comparisons of the composition of spider assemblages in three vegetation habitats in Bogor, West Java, Indonesia (<https://smujo.id/biodiv/article/view/9190>)

MAYANDA LIA, AUNU RAUF, DADAN HINDAYANA

PDF (<https://smujo.id/biodiv/article/view/9190/5435>)

Genetic variation of *Austropuccinia psidii* in some species of Myrtaceae as host plants in Java, Indonesia based on simple sequence repeats (SSR) markers (<https://smujo.id/biodiv/article/view/9642>)

FARAH AULYA FARADILLA, ISTIANA PRIHATINI, SURANTO, ARI SUSILOWATI

PDF (<https://smujo.id/biodiv/article/view/9642/5427>)

Ethnobotanical knowledge of Marind-Anim Tribe in utilizing sago (*Metroxylon sagu*) in Merauke, Papua, Indonesia (<https://smujo.id/biodiv/article/view/9843>)

AKHMAD KADIR, SUHARNO, YOHANA REAWARUW, KOMARI, AGUSTINUS MAHUZE

PDF (<https://smujo.id/biodiv/article/view/9843/5428>)

Growth and reproductive biology of white-spotted rabbitfish (*Siganus canaliculatus*) on different seagrass habitats in Inner Ambon Bay, Indonesia (<https://smujo.id/biodiv/article/view/9342>)

HUSAIN LATUCONSINA, M. MUKHLIS KAMAL, RIDWAN AFFANDI, NURLISA A. BUTET

PDF (<https://smujo.id/biodiv/article/view/9342/5445>)

Cartography and spatial distribution of the genus *Pistacia* in Souk Ahras, Northeast Algeria (<https://smujo.id/biodiv/article/view/9649>)

ALI ABDELMOTALAB BAROUR, MOUNDJI TAOUARFIA

PDF (<https://smujo.id/biodiv/article/view/9649/5438>)

Agronomic characteristics and genetic relationship of putative transgenic rice lines of cv. Fatmawati with the Glu-1Dx5 transgene (<https://smujo.id/biodiv/article/view/9730>)

NONO CARSONO, GIGIH IBNU PRAYOGA, SANTIKA SARI, MEDDY RACHMADI

PDF (<https://smujo.id/biodiv/article/view/9730/5439>)

Bioremediation of batik wastewater by *Rhizobacteria* isolated from iron sand soils tolerant of Pb and Zn (<https://smujo.id/biodiv/article/view/9592>)

OEDJIJONO, SRI LESTARI, LESYA SURYANI SAMSUDIN, HERMILIA

PDF (<https://smujo.id/biodiv/article/view/9592/5446>)

Traditional medicinal plants and their utilization by local communities

around Lambung Mangkurat Education Forests, South Kalimantan, Indonesia (<https://smujo.id/biodiv/article/view/9756>)

YUSANTO NUGROHO, MOCHAMAD ARIEF SOENDJOTO, SUYANTO, JERIELS MATATULA, SYAMSU ALAM, PANDU YUDHA ADI PUTRA WIRABUANA

PDF (<https://smujo.id/biodiv/article/view/9756/5451>)

Short communication: Amphidromous goby postlarvae (penja) migration seasons and fisheries in West Sulawesi, Indonesia (<https://smujo.id/biodiv/article/view/9789>)

NURJIRANA, ANDI I. BURHANUDDIN, PHILIPPE KEITH, ABDUL HARIS, MUHAMMAD AFRISAL

PDF (<https://smujo.id/biodiv/article/view/9789/5441>)

Characteristics of physiology, phenology, and drought susceptibility index of two varieties of Job's tears under water deficit stress (<https://smujo.id/biodiv/article/view/9910>)

FIKY YULIANTO WICAKSONO, UMA RANI SINNIHA, RUMINTA, SUMADI, TATI NURMALA

PDF (<https://smujo.id/biodiv/article/view/9910/5440>)

Comparative diurnal and seasonal variations of ACTH, cortisol and aldosterone in Ouled Djellal and D'Man sheep breeds reared in arid lands (<https://smujo.id/biodiv/article/view/9654>)

ASSIA AMOKRANE-FERRAH, ASSIA ANANE, NOURIA BOUKENAOUI-FERROUK, MOUNIRA KHALDOUN, ZAINA AMIRAT, PIERRE MORMEDE, FARIDA KHAMMAR

PDF (<https://smujo.id/biodiv/article/view/9654/5442>)

Orangutan (*Pongo pygmaeus wurmbii*) human-oriented behavior at the Lamandau Wildlife Reserve, Indonesia (<https://smujo.id/biodiv/article/view/10120>)

NOVITA AMALIA, DYAH PERWITASARI FARAJALLAH, SRI SUCI UTAMI ATMOKO

PDF (<https://smujo.id/biodiv/article/view/10120/5443>)

Aboveground biomass and carbon stock of *Rhizophora apiculata* forest in Ca Mau, Vietnam (<https://smujo.id/biodiv/article/view/10092>)

TRAN QUANG BAO, NGUYEN THI HA, BUI THI MINH NGUYET, VO MINH HOAN, LE HONG VIET, DANG VIET HUNG

PDF (<https://smujo.id/biodiv/article/view/10092/5444>)

Short communication: Algal genotypes in White Syndrome infected coral *Acropora muricata* from the Karimunjawa Islands, Indonesia
(<https://smujo.id/biodiv/article/view/9458>)

DIAH PERMATA WIJAYANTI, MAULA NADIA, ELIS INDRAYANTI, DWI HARYANTI, MUGGI BACHTIAR

PDF (<https://smujo.id/biodiv/article/view/9458/5447>)

Indigenous Dayak Iban customary perspective on sustainable forest management, West Kalimantan, Indonesia
(<https://smujo.id/biodiv/article/view/9928>)

SANDY LEO, JATNA SUPRIATNA, KOSUKE MIZUNO, CHRIS MARGULES

PDF (<https://smujo.id/biodiv/article/view/9928/5452>)

Population dynamics of *Anadara antiquata* of East Coast of Aceh, Indonesia
(<https://smujo.id/biodiv/article/view/9912>)

FAUZIAH AZMI, ABDUL L. MAWARDI, MUH. SALEH NURDIN, SURI PURNAMA FEBRI, SORBAKTI SINAGA, TEUKU FADLON HASER

PDF (<https://smujo.id/biodiv/article/view/9912/5448>)

Aneuk Jamee traditional foods in the South Aceh District, Indonesia
(<https://smujo.id/biodiv/article/view/10003>)

SYAMSUARDI, NURAINAS, AHMAD TAUFIQ, TISNA HARMAWAN, ADI BEJO SUWARDI

PDF (<https://smujo.id/biodiv/article/view/10003/5454>)

Identification of poisonous plants and their solutions for traditional livestock in Bojonegoro District, East Java, Indonesia
(<https://smujo.id/biodiv/article/view/10124>)

ANGGI MUHTAR PRATAMA, OKTI HERAWATI, NURI RAHMA NURANISA, NURUL HANIFAH, AGUSTINA DWI WIJAYANTI, SATYAGUNA RAHMATULLAH, ELFIA NURAINI, AGUNG BUDIYANTO

PDF (<https://smujo.id/biodiv/article/view/10124/5449>)

Anti-bacteria and toxicity potential of a rare *Actinobacterium*

Pseudonocardia sp. SM1A, isolated from Mangrove Park, West Kalimantan, Indonesia (<https://smujo.id/biodiv/article/view/9713>)

RISA NOFIANI, RIZKY, RIDHO BRILANTORO, PUJI ARDININGSIH

PDF (<https://smujo.id/biodiv/article/view/9713/5457>)

Effect of soil properties on plant growth and diversity at various ages of coal mine reclamation in Indonesia (<https://smujo.id/biodiv/article/view/9729>)

TEDI YUNANTO, FARISATUL AMANAH, AYU RATNA WULANSARI, NABILA PUTRI WISNU

PDF (<https://smujo.id/biodiv/article/view/9729/5458>)

The enrichment process and morphological identification of anaerobic fungi isolated from buffalo rumen (<https://smujo.id/biodiv/article/view/9865>)

SINTA AGUSTINA, KOMANG GEDE WIRYAWAN, SRI SUHARTI, ANJA MERYANDINI

PDF (<https://smujo.id/biodiv/article/view/9865/5455>)

The role of arbuscular mycorrhizal fungi in phytoremediation of heavy metals and their effect on the growth of *Pennisetum purpureum* cv. Mott on gold mine tailings in Muara Bungo, Jambi, Indonesia (<https://smujo.id/biodiv/article/view/9933>)

BELA PUTRA, LILI WARLY, EVITAYANI, BOPALION PEDI UTAMA

PDF (<https://smujo.id/biodiv/article/view/9933/5459>)

Diversity of cellulolytic bacteria from *Macrotermes gilvus* gut isolated from Indralaya peatland region, Indonesia (<https://smujo.id/biodiv/article/view/9797>)

DWITA OKTIARNI, GETARI KASMIARTI, ERWIN NOFYAN, MIKSUSANTI, HASANUDIN, HERMANSYAH

PDF (<https://smujo.id/biodiv/article/view/9797/5465>)

Tree density impact on growth, roots length density, and yield in agroforestry based cocoa (<https://smujo.id/biodiv/article/view/9920>)

ABDUL RAHIM SALEH, SIKSTUS GUSLI, AMBO ALA, RISMA NESWATI, SRI SUDEWI

PDF (<https://smujo.id/biodiv/article/view/9920/5464>)

Growth and morpho-physiology of *Tectona philippinensis* under different

water stress and soil conditions (<https://smujo.id/biodiv/article/view/10048>)

JONATHAN O. HERNANDEZ, CZAHAINE A. TOLENTINO, LORRAINE ANNE E. QUIÑONES,
GERARDO B. VILLANCIO, LERMA SJ. MALDIA, MARILYN S. COMBALICER

PDF (<https://smujo.id/biodiv/article/view/10048/5462>)

Object based classification of benthic habitat using Sentinel 2 imagery by
applying with support vector machine and random forest algorithms in
shallow waters of Kepulauan Seribu, Indonesia
(<https://smujo.id/biodiv/article/view/10087>)

HARTONI, VINCENTIUS P. SIREGAR, SAM WOUTHUYZEN, SYAMSUL BAHRI AGUS

PDF (<https://smujo.id/biodiv/article/view/10087/5463>)

Antibacterial activity of endophytic fungi isolated from the stem bark of
jambu mawar (*Syzygium jambos*) (<https://smujo.id/biodiv/article/view/9353>)

KURRATUL 'AINI, ELFITA, HARY WIDJAJANTI, ARUM SETIAWAN, ALFIA R. KURNIAWATI

PDF (<https://smujo.id/biodiv/article/view/9353/5468>)

Behavioral, physiological, and blood biochemistry of Friesian Holstein dairy
cattle at different altitudes in West Java, Indonesia
(<https://smujo.id/biodiv/article/view/9615>)

UJANG HIDAYAT TANUWIRIA, IIN SUSILAWATI, DIDIN S. TASRIFIN, LIA BUDIMULYATI
SALMAN, ANDI MUSHAWWIR

PDF (<https://smujo.id/biodiv/article/view/9615/5469>)

Evaluation of the advanced yield trial on tropical wheat (*Triticum aestivum*)
mutant lines using selection index and multivariate analysis
(<https://smujo.id/biodiv/article/view/9621>)

MUH FADLI, MUH FARID, AMIR YASSI, NASARUDDIN, MUHAMMAD FUAD ANSHORI, AMIN
NUR, SURATMAN

PDF (<https://smujo.id/biodiv/article/view/9621/5470>)

Short Communication: Characterization and biological synthesis of zinc
oxide nanoparticles by new strain of *Bacillus foraminis*
(<https://smujo.id/biodiv/article/view/9772>)

DINA E. EL-GHWAS

PDF (<https://smujo.id/biodiv/article/view/9772/5471>)

Biodiversity of ethnomedicinal plants from the B'laan Tribe in Mount Matutum Protected Landscape, Southern Mindanao, Philippines (<https://smujo.id/biodiv/article/view/10051>)

MALONA V. ALINSUG, MARK HAROLD G. ESTANDARTE, EDEN MAY N. SOMODIO, MARIEL JADE J. SABARITA, CUSTER C. DEOCARIS

PDF (<https://smujo.id/biodiv/article/view/10051/5472>)

Potential of dadiah kapau from Agam District, West Sumatra, Indonesia as a source of probiotics for health (<https://smujo.id/biodiv/article/view/9676>)

ELLY ROZA, SALAM N. ARITONANG, YULIA YELLITA, HILDA SUSANTY, RIZQAN, YUDHA ENDRA PRATAMA

PDF (<https://smujo.id/biodiv/article/view/9676/5473>)

Cytotoxicity extract and fraction of knobweed (*Hyptis capitata*) and its effect on migration and apoptosis of T47D cells (<https://smujo.id/biodiv/article/view/9643>)

NELSIANI TO'BUNGAN, RARASTOETI PRATIWI, SITARINA WIDYARINI, LAURENTIUS HARTANTO NUGROHO

PDF (<https://smujo.id/biodiv/article/view/9643/5474>)

Community awareness and participation in biodiversity conservation at Phong Nha-Ke Bang National Park, Vietnam (<https://smujo.id/biodiv/article/view/9718>)

DINH DUC TRUONG

PDF (<https://smujo.id/biodiv/article/view/9718/5475>)

The growth of tiger shrimp (*Penaeus monodon*) and its dynamics of water quality in integrated culture (<https://smujo.id/biodiv/article/view/9765>)

ROSA AMALIA, SRI REJEKI, LESTARI LAKSHMI WIDOWATI, RESTIANA WISNU ARIYATI

PDF (<https://smujo.id/biodiv/article/view/9765/5476>)

Pollen morphology of some species in family Amaranthaceae from Thailand (<https://smujo.id/biodiv/article/view/10034>)

SURAPON SAENSOUK, PIYAPORN SAENSOUK

PDF (<https://smujo.id/biodiv/article/view/10034/5477>)

Information

For Readers (<https://smujo.id/biodiv/information/readers>)

For Authors (<https://smujo.id/biodiv/information/authors>)

For Librarians (<https://smujo.id/biodiv/information/librarians>)

Journals List

Biodiversitas Journal of Biological Diversity (<https://smujo.id/biodiv>)

Nusantara Bioscience (<https://smujo.id/nb>)

Prosiding Seminar Nasional Masyarakat Biodiversitas Indonesia (<https://smujo.id/psnmbi>)

Asian Journal of Agriculture (<https://smujo.id/aja>)

Asian Journal of Ethnobiology (<https://smujo.id/aje>)

Asian Journal of Forestry (<https://smujo.id/ajf>)

Asian Journal of Natural Product Biochemistry (<https://smujo.id/jnpb>)

Asian Journal of Tropical Biotechnology (<https://smujo.id/bbs>)

International Journal of Bonorowo Wetlands (<https://smujo.id/bw>)

Cell Biology and Development (<https://smujo.id/cbd>)

Indo-Pacific Journal of Ocean Life (<https://smujo.id/ol>)

International Journal of Tropical Drylands (<https://smujo.id/td>)

Reviewers List

Reviewers (<https://smujo.id/biodiv/reviewers/index>)

Visitor Statistics

Statistics (<https://smujo.id/info/stats>)



(<https://info.flagcounter.com/JKar>)



(<https://www.scopus.com/sourceid/21100332431>)



(<http://doaj.org/toc/3eda1e70aa014e1e8bd1fc367b4df956>)



(<http://scholar.google.co.id/citations?hl=id&user=rae-lrEAAAAJ>)



(<http://search.crossref.org/?q=biodiversitas&type=Journal>)



(<https://academic.microsoft.com/#/detail/2738269101>)