

# Freshwater fishes of Gegas Dam, South Sumatra Indonesia: composition and diversity

*By Arum Setiawan*

## Freshwater fishes of Gegas Dam, South Sumatra Indonesia: composition and diversity

Dian Samitra<sup>1,3\*</sup>, Harmoko<sup>1</sup>, Sepriyaningsih<sup>1</sup>, Zico Fakhurur Rozi<sup>1</sup>, Arum Setiawan<sup>2</sup> and Indra Yustian<sup>2</sup>

<sup>1</sup>STKIP PGRI Lubuklinggau, Jalan. Mayor Toha Kec. Lubuklinggau Timur 1, Lubuklinggau, South Sumatra, Indonesia

<sup>2</sup>Departement of Biology, Faculty of Science, Universitas Sriwijaya, Jalan Raya Palembang-Prabumulih km 32, Indralaya, South Sumatra, 30662, Indonesia.

<sup>3</sup>Zoology Division, Generasi Biologi Indonesia Fundation (Genbinesia), Jalan Swadaya Barat No.4, Gersik 61171, East Java, Indonesia

(Received 5 July, 2020; accepted 11 August, 2020)

### ABSTRACT

Fish studies can be essential information about the health of aquatic ecosystems as well as a basis for policy making management of fish resources. Study of fishes in the Gegas Dam has never been reported. This study aims to determine the diversity of fish in the Gegas Dam. Data collection was conducted during rainy season in March 2020 at the Gegas Dam, Musi Rawas Regency, South Sumatra. Fish samples were caught with fish bait, cast net, gill net, and hexagonal fish trap. The fish obtained are tabulated based on species, genus and family. Data analysis included the index of Shannon diversity, Pielou's evenness index and Simpson dominance index. Total fishes collected during the study were 414 individuals. The composition of the fish consists of 25 species representing 19 genera, and 12 families. *Desmopuntius gemellus* is the species most caught during the study. The results of diversity index analysis were 3.01, evenness index was 0.93 and dominance index was 0.05. Diversity in the Gegas Dam is of medium category with high species evenness and no species dominate.

**Key words :** Biodiversity, Ecology, Ecosystems, Fish, Population

### Introduction

Dam construction is a plays role in human development to control river flow and is used to generate electricity, and irrigation (Agostinho *et al.*, 2008; Jellyman & Harding, 2012). The dam threat to biodiversity is due to changing functional characteristics of biological communities, ecosystem processes and creating new ecosystems (Baxter, 1977; Cardinale *et al.*, 2012; Oliveira *et al.*, 2018). Dams change anthropogenic, including water pollution, overfishing, lake reclamation for agricultural land

(Hu *et al.*, 2015). Dam can also cause problems for fish i.e diversity, migration delay, non-native species (Agostinho *et al.*, 2008; Azami *et al.*, 2012; Cheng *et al.*, 2015; Jawad, 2003; Marschall *et al.*, 2011; Sá-Oliveira *et al.*, 2015).

Fish studies in South Sumatra has been widely carried out in rivers such as the Kelingi River (Samitra and Rozi, 2018a), Lakitan River (Samitra and Rozi, 2019b), Musi River (Prianto and Nurdawaty, 2013; Prianto and Suryanti, 2010, and Iqbal *et al.*, 2018) and the Banyuasin River (Prianto and Aprianti, 2012). The study focused on the river,

even though a complete study including the dam needs to be done. Fish studies can be essential information about the health of aquatic ecosystems and become the basis for policy making management of fish resources and conservation of an aquatic area (Hasan *et al.*, 2015; Mote, 2017; Sukmono *et al.*, 2013). In addition, the fish study of dam is helpful in problems like pollution control, the construction and renovation of dams and lakes, fish and aquatic life (Theurkar *et al.*, 2013). This study aims to determine the diversity of fish in the Gegas Dam.

## Materials and Methods

The field study was carried out in March 2020 at the Gegas Dam, Musi Rawas, South Sumatra, with coordinates 3°14'57.9"S 103°08'10.2" E (Figure 1). Fish sample were collected through fishing efforts using fish bait, cast net, gill nets, hexagonal fish trap and also help from the local Fisherman. Sampling is done at the selected point by purposive sampling based on potential spots of fish. Captured fish were identified refers to Kottelat *et al.* (1993); Iqbal (2011); Iqbal *et al.* (2018) and Sukmono and Margaretha (2017).



Fig. 1. Location of the Gegas Dam, Musi Rawas, South Sumatra

Fish data are tabulated by species and family. Data analysis includes species composition, Shannon diversity index, Pielou's evenness index and Simpson dominance index (Guo *et al.*, 2018; Hossain

*et al.*, 2017; Samitra and Rozi, 2019b).

## Results

### Fish composition

A total of 414 fish sample were caught during research. The fish composition consists of 25 species representing 19 genera, and 12 families (Table 1). Three families [Cyprinidae (n= 237 specimens, 9 species), Clariidae (n= 23 specimens, 3 species) and Osphronemidae (n=29 specimens, 3 species)] are the most species-rich families. The most common species found in Gegas Dam *Desmopuntius gemellus* (8.45%), while the least commonly found was *Pangio kuhlii* and *Chitala lopis* (0.24%) (Table 1).

Table 1. Fish Composition in Gegas Dam

Families	Species	Abundance Relative (%)
Anabantidae	<i>Anabas testudineus</i>	3.14
Aplocheilidae	<i>Aplocheilus panchax</i>	6.52
Bagridae	<i>Mystus nigriceps</i>	3.38
Channidae	<i>Channa striata</i>	2.66
	<i>Channa micropeltes</i>	3.14
Clariidae	<i>Clarias leiacanthus</i>	2.66
	<i>Clarias nieuhofii</i>	2.17
	<i>Clarias gariepinus</i>	0.72
Cobitidae	<i>Pangio kuhlii</i>	0.24
Cyprinidae	<i>Hampala macrolepidota</i>	6.28
	<i>Puntigrus tetrazona</i>	6.76
	<i>Cyclocheilichthys apogon</i>	5.07
	<i>Osteochilus vittatus</i>	3.86
	<i>Rasbora trilineata</i>	6.28
	<i>Labiobarbus leptocheilus</i>	6.28
	<i>Desmopuntius gemellus</i>	8.45
	<i>Rasbora nematotaenia</i>	7.49
	<i>Rasbora cephalotaenia</i>	6.76
Helostomatidae	<i>Helostoma temminckii</i>	5.56
Nandidae	<i>Nandus nebulosus</i>	1.21
Notopteridae	<i>Chitala lopis</i>	0.24
Osphronemidae	<i>Osphronemus goramy</i>	1.21
	<i>Trichopodus trichopterus</i>	3.62
	<i>Trichopodus pectoralis</i>	2.17
Pristolepididae	<i>Pristolepis fasciata</i>	4.11

### Ecological index

The results of the data analysis in the field revealed that the diversity of fish in the Gegas Dam was 3.01 with the medium category, while the evenness index was 0.93 with the evenly distributed category and the index of dominance was 0.05 with the low category (Table 2). Environmental parameter data

(Table 3) shows that water temperature is 27.5, visibility is 93.45 cm, pH is 6.87, and DO is 40 mg.L<sup>-1</sup>. Based on these data it can be said that the environmental parameters in the Gegas Dam are still within the tolerance limits for fish growth and development.

**Table 2.** Index of Diversity, Evenness and Dominance in Gegas Dam

Index	Value
Diversity (H')	3.01
Evenness (E)	0.93
Dominance (C)	0.05

**Table 3.** Data on Environmental Parameters at Gegas Dam

Parameter	Average
Water Temperature (!)	27.5
Visibility (cm)	93.45
pH	6.87
DO (mg/L)	42.3

## Discussion

The fish most common in Gegas Dam are different from the results of research in Lakitan Dam (*Barbonymus gonionotus*) and Perjaya Weir (*Mystacoleucus marginatus*) (Nizar *et al.*, 2014; Samitra and Rozi, 2018b) in Gegas Dam the most caught fish is *Desmopuntius gemellus*. This can affect the habitat in Dam Gegas suitable for the growth of *Desmopuntius gemellus*, habitat because it gives importance to the success of fish (Thomas *et al.*, 2003). The difference in species in each water due to differences in habitat characteristics, food composition as well as chemical and physical factors (Daga *et al.*, 2012; Tobes *et al.*, 2016).

Fish *Chitala Lopis* and *Pangio kuhlii* the fewest fish found in the Gegas Dam. Some studies explain that the population of belida fish (*Chitala sp*) this has decreased due to human activities that are not environmentally friendly, changes in environmental conditions (Nugroho *et al.*, 2019; Wibowo *et al.*, 2010). *Chitala sp* is much in demand by the community so that fishermen over fishing even though this fish has a low fecundity, so it can cause these fish to become scarce (Wibowo *et al.*, 2010). The Government of Indonesia in 2018 has determined that all types of belida fish in Indonesia are protected (Indo-

nesian Government, 2018)

We found *Pangio kuhlii* in puddles that arise after rain around the Gegas Dam. Though these fish live in forest canals and peat waters (Froese and Pauly, 2019), so the possibility of these fish carried by the current when it rains. This fish can be used as ornamental fish and has been traded in the domestic and export markets (Fahmi *et al.*, 2015).

The Cyprinidae family is the fish family most commonly found in the Gegas Dam, this is the same as several other studies such as in the Lakitan Dam, Musi Rawas, South Sumatra (Samitra and Rozi, 2019a, 2018b), Perjaya Weir, Komering River in South Sumatra (Nizar *et al.*, 2014) and Kelingi River, South Sumatra (Samitra and Rozi, 2018a) and the Luk Ulo River, Kebumen (Wahyuni and Zakaria, 2018). Cyprinidae is the largest family in the World except in Australia, Madagascar, New Zealand, and South America (although in some places it has been introduced) (Esmaeili *et al.*, 2010). They are easily adaptable, most commonly found in tropical waters and generally plays an important role in supporting human life (Kottelat and Whitten, 1996; Samitra and Rozi, 2019b; Thai *et al.*, 2007).

We refer to Froese & Pauly (2019) for species distribution and habitat. The native species were found as many as 24 species and non-native species as many as 1 species of fish. We did not find endemic fish in South Sumatra, but found the endemic fish of Sumatra, namely *Desmopuntius gemellus*. The non-native species found was *Clarias gariepinus*, this fish from Africa and was cultivated for consumption fish (Froese and Pauly, 2019; Muhammad and Andriyanto, 2013). We interview community around dam, that the Government of Musi Rawas Regency has ever released (introduced) fish such as *Clarias gariepinus*, *Pangasius sp*, and *Oreochromis niloticus*. The non-native species carried out by the government might aim fortaken restocking, this activity needs to beinto consideration because many studies have shown the non-native species can reduce native fish (Britton and Orsi, 2012; Latini and Jr, 2004; Nuryanto *et al.*, 2015; Umar and Sulaiman, 2013). The non-native species very dangerous for native fishes of the Gegas Dam, because it can be a competitor of native fish in obtaining habitat and food, predation, ecological changes (Gallardo *et al.*, 2016; Grabowska *et al.*, 2020; Ribeiro and Leunda, 2012). In fact, native fish can be cultivated for consumption and ornamental fish. *Osphronemus goramy*, *Hampala macrolepidota*, *Helostoma temminckii*,

*Pristolepis fasciata* and *Chitala lopis* for consumption. *Pangio kuhlii* and fish of the genus *Rasbora* can be cultivated into ornamental fish.

The diversity of fish in the Gegas Dam is of medium category, with evenly distributed species and low dominance. Fish diversity in the Gegas Dam is lower than in the Perjaya Dam (Nizar *et al.*, 2014), and higher than the diversity of fish in the Simpung Dam (Gustomi *et al.*, 2013). Diversity index depends on variation in the number of individuals of each species (Hossain *et al.*, 2013; Sihombing *et al.*, 2017). The greater the number of fish species and the variation in the number of individuals of each species, the greater the diversity of fish in an aquatic ecosystem (Grenouillet, *et al.*, 2002; Sriwidodo *et al.*, 2013).

The category diversity is showing the diversity of fish in the Gegas Dam in good condition because it is supported by a stable ecosystem and there is no disturbance. The dominance of fish in the Gegas Dam can be said to be relatively nonexistent as the dominance index value reaches almost 0. The absence of high dominance in the Gegas Dam signifies competition for balanced food and habitat as well as a balance between predators and prey (Zulfikri *et al.*, 2016; Sinclair *et al.*, 2006). The government has banned the use of electrofishing and poison, helping to maintain the diversity of fish in dams. Data on environmental parameters in the Gegas Dam still (Table 3) support fish life, where the optimum temperature is 20-30 °C, pH 6-9, DO > 4 mg.L<sup>-1</sup> (Wahyuni and Zakaria, 2018).

## Conclusion

Diversity in the Gegas Dam is of medium category with high species evenness and no species dominate.

## Acknowledgements

This research was funded by the Indonesian Ministry of Education and Culture. We would like to thank the Lubuklinggau Kehati Team for helping us gather data.

## References

Agostinho, A., Pelicice, F. and Gomes, L. 2008. Dams and the fish fauna of the Neotropical region : impacts and management related to diversity and fisheries. *Braz. J. Biol.* 68(4) : 1119-1132.

- Azami, K., Takemoto, M., Otsuka, Y., Yamahishi, S. and Nakazawa, S. 2012. Meteorology and species composition of plant communities, birds and fishes before and after initial impoundment of Miharu Dam Reservoir, Japan. *Landscape Ecol Eng.* 2012 : 81-105.
- Baxter, R.M. 1977. Environmental effects of dams and impoundments. *Annu. Rev. Ecol. Syst.* 8 : 255-283.
- Britton, J. R. and Orsi, M. L. 2012. Non-native fish in aquaculture and sport fishing in Brazil/: economic benefits versus risks to fish diversity in the upper Paraná River Parana. *Rev Fish Biol Fisheries.* 22 : 555-565.
- Cardinale, B. J., Duffy, J. E., Gonzalez, A., Hooper, D. U., Perrings, C. and Venail, 2012. Biodiversity loss and its impact on humanity. *Nature.* 486 : 59-65.
- Cheng, F., Li, W., Castello, L., Murphy, B. R. and Xie, S. 2015. Potential effects of dam cascade on fish/: lessons from the Yangtze River. *Reviews in Fish Biology and Fisheries.*
- Daga, V. S., Gubiani, É. A. and Cunico, A. M. 2012. Effects of abiotic variables on the distribution of fish assemblages in streams with different anthropogenic activities in southern Brazil. *Neotropical Ichthyology.* 10(3) : 643-652.
- Esmaili, H.R., Zareian, H., Gholamhosseini, A., Ebrahimi, M., Gholami, Z., Teimori, A. and Ansari, T.H. 2010. Karyotype Analysis of the King Nase Fish, *Chondrostoma regium* (Heckel, 1843) (Actinopterygii: Cyprinidae) from Iran. *Turkish Journal of Fisheries and Aquatic Sciences.* 10 : 477-481.
- Fahmi, M. R., Ginanjar, R. and Kusumah, R. V. 2015. *Keragaman Ikan Hias di Lahan Gambut Cagar Biosfer Bukit-Batu, Propinsi Riau.* Seminar Nasional Masyarakat Biodiversitas Indonesia, June, 51-58.
- Froese, R. and Pauly, D. 2019. <https://www.fishbase.de/> (Accessed 10 May 2020)
- Gallardo, B., Clavero, M., I, S. M. and Vila, M. 2016. Global ecological impacts of invasive species in aquatic ecosystems. *Global Change Biology.* 22 : 151-163.
- Grabowska, J., Jan, K. and Witkowski, A. 2020. Alien invasive fish species in Polish waters/: an overview Alien invasive fish species in Polish waters/: an overview. *Folia Zoologica.* 59 (1) : 73-85.
- Grenouillet, G., Pont, D. and Seip, K. L. 2002. Abundance and species richness as a function of food resources and vegetation structure: juvenile fish assemblages in rivers. *Ecography.* 25(6) : 641-650.
- Guo, Q., Liu, X., Ao, X., Qin, J., Id, X. W. and Ouyang, S. 2018. Fish diversity in the middle and lower reaches of the Ganjiang River of China: Threats and conservation. *PLOS ONE.* 13(11) : 1-17.
- Gustomi, A., Sulistiono, and Vitner, Y. 2013. *Keanekaragaman Sumber Daya Ikan di Kolong - Bendungan Simpung Kabupaten Bangka Provinsi Bangka Belitung.* Seminar Nasional Ikan Ke 8, 10 (1) : 33-39.
- Hasan, Z., Khan, M. ., Ali, Z., Zia, Q., Masood, Z. and

- Khan, W. 2015. Fish Diversity of Sharki Dam, District Karak, Khyber Pakhtunkhwa, Pakistan. *Sindh University Research Journal*. 47(1) : 167–170.
- Hossain, M. A., Akter, M. and Iqbal, M. M. 2017. Diversity of Fish Fauna in Kusiara River (Fenchungonj Upazilla), Northeast Bangladesh. *Journal of Aquaculture in the Tropics*. 32 : 1–13.
- Hossain, M. S., Das, N. G., Sarker, S. and Rahaman, M. Z. 2013. Fish diversity and habitat relationship with environmental variables at Meghna river estuary, Bangladesh. *The Egyptian Journal of Aquatic Research*. 38(3) : 213–226.
- Hu, M., Hua, Q., Zhou, H. and Wu, Z. 2015. The effect of dams on the larval abundance and composition of four carp species in key river systems in China. *Environ Biol Fish*. 98 : 1201–1205.
- Indonesian Government. 2018. [http://ksdae.menlhk.go.id/assets/news/peraturan/P.20\\_Jenis\\_TSL\\_.pdf](http://ksdae.menlhk.go.id/assets/news/peraturan/P.20_Jenis_TSL_.pdf) (Accessed 10 May 2020)
- Jawad, L. A. 2003. Impact of Environmental Change on The Freshwater Fish Fauna of Iraq. *Intern. J. Environ. Studies*. 60(6) : 581–593.
- Jellyman, P. G. and Harding, J. S. 2012. The role of dams in altering freshwater fish communities in New Zealand. *New Zealand Journal of Marine and Freshwater Research*. 46 (4) : 475–489.
- Iqbal, M. 2011. *Ikan-ikan di Hutan Rawa Gambut Merang Kepayang dan sekitarnya*. Merang REDD Pilot Project, Sumatra Selatan, 92 pp.
- Iqbal, M., Yustian, I., Setiawan, A. and Setiawan, D. 2018. *Ikan-Ikan di Sungai Musi dan Pesisir Timur Sumatera Selatan*. Kelompok Pengamat Burung Spirit of South Sumatra bekerjasama dengan Jurusan Biologi FMIPA Universitas Sriwijaya and Zoological Society for the Conservation of Species and Populations, Palembang, 294 pp
- Kottelat, M., Whitten, J.A., Kartikasari, N. and Wiryatmojo, S. 1993. *Freshwater Fishes of Western Indonesia and Sulawesi*. Periplus Edition, Jakarta, 293 pp
- Kottelat, M. and Whitten, A. J. 1996. [https://www.researchgate.net/publication/270508914\\_Freshwater\\_fishes\\_of\\_western\\_Indonesia\\_and\\_Sulawesi\\_additions\\_and\\_corrections](https://www.researchgate.net/publication/270508914_Freshwater_fishes_of_western_Indonesia_and_Sulawesi_additions_and_corrections) (Accessed 10 April 2020)
- Latini, A. O. and Jr, M. P. 2004. Reduction of a native fish fauna by alien species: an example from Brazilian freshwater tropical lakes. *Fisheries Management and Ecology*. 11 : 71–79.
- Marschall, E. A. M., Mather, M. E. M., Parrish, D. L. P., Llison, G. W. and McMenemy, J. R. 2011. Migration delays caused by anthropogenic barriers : modeling dams, temperature, and success of migrating salmon smolts. *Ecological Applications*. 21 (8) : 3014–3031.
- Mote, N. 2017. Biodiversitas Iktiofauna Di Muara Sungai Kumbe Kabupaten Merauke. *Al-Kauniyah: Jurnal Biologi*. 10(1) : 26–34.
- Muhammad, W. N. and Andriyanto, S. 2013. Manajemen Budidaya Ikan Lele Dumbo (*Clarias Gariepinus*) Di Kampung Lele, Kabupaten Boyolali, Jawa Tengah. *Media Akuakultur*. 8(1) : 63–71.
- Nizar, M., Mukhlis Kamal, M. and M Adiwilaga, E. 2014. Komposisi jenis dan struktur komunitas ikan yang bermigrasi melewati tangga ikan pada Bendung Perjaya, Sungai Komerang, Sumatera Selatan. *Depik*. 3(1) : 27–35.
- Nugroho, E., Roro, R., Pudji, S., Dewi, S., Subagja, J. and Priono, B. 2019. Keragaman Genetik Dan Karakter Biometrik Ikan Belida (*Chitala Lopis*, Bleeker 1851) Budidaya Asal Sungai Kampar, Riau. *Jurnal Riset Akuakultur*. 14(1) : 1–8.
- Nuryanto, A., Bhagawati, D. and Abulias, M. N. 2015. Fauna ikan di Sungai Cikawung Kabupaten Cilacap Jawa Tengah. *Jurnal Iktiologi Indonesia*. 15(1) : 25–37.
- Oliveira, A. G., Baumgartner, M. T., Gomes, L. C., Dias, R. M. and Agostinho, A. A. 2018. Long-term effects of flow regulation by dams simplify fish functional diversity. *Freshwater Biology*. 63(3) : 293–305.
- Prianto, E. and Aprianti, S. 2012. Komposisi jenis dan biomasa stok ikan di Sungai Banyuasin. *Jurnal Penelitian Perikanan Indonesia*. 18(1): 1–8.
- Prianto, E. and Nurdawaty, S. 2013. Distribusi, Kelimpahan Dan Variasi Ukuran Larva Ikan Di Estuaria Sungai Musi. *Bawal*. 5(2) : 73–79.
- Prianto, E. and Suryanti, N. K. 2010. Komposisi jenis dan potensi sumber daya ikan di Muara Sungai Musi. *Jurnal Penelitian Perikanan Indonesia*. 16(1): 1–8.
- Ribeiro, F. and Leunda, P. M. 2012. Non-native fish impacts on Mediterranean freshwater ecosystems : current knowledge and research needs. *Fisheries Management and Ecology*. 19 : 142–156.
- Sá-Oliveira, J. C., Hawes, J. E., Isaac-Nahum, V. J. and Peres, C. A. 2015. Upstream and downstream responses of fish assemblages to an eastern Amazonian hydroelectric dam. *Freshwater Biology*. 60(10): 2037–2050.
- Samitra, D. and Rozi, Z. F. 2018a. Keanekaragaman Ikan di Sungai Kelingi Kota Lubuklinggau. *Jurnal Biot*. 4(1) : 1–6.
- Samitra, D. and Rozi, Z. F. 2018b. *Keanekaragaman Ikan Air Tawar Di Bendungan Lakitan Kabupaten Musi Rawas Provinsi Sumatera Selatan*. Seminar Nasional Sains Dan Teknologi Terapan, Palembang, Indonesia
- Samitra, D. and Rozi, Z. F. 2019a. Potensi dan Status Konservasi Ikan Di Bendungan Lakitan Kabupaten Musi Rawas, Provinsi Sumatera Selatan. *Pro-Life*. 6(1) : 13.
- Samitra, D. and Rozi, Z. F. 2019b. The Fish Fauna in Lakitan River, Musi Rawas Regency, South Sumatra. *Jurnal Biodjati*. 4 (1) : 11–20.
- Sihombing, V. S., Gunawan, H. and Sawitri, R. 2017. Diversity and community structure of fish, plankton and benthos in Karangsang Mangrove Conservation

- Areas, Indramayu, West Java, Indonesia. *Biodiversitas*. 18(2) : 601–608.
- Sinclair, A.R.E., Fryxell, J.M. and Caughley, G. 2006. *Wild-life Ecology, Conservation, And Management 2<sup>nd</sup> Edition*. Blackwell Publishing, United Kingdom. 469 pp
- Sriwidodo, D. E., Budiharjo, A. and Sugiyarto, 2013. Keanekaragaman Jenis Ikan di Kawasan Inlet dan Outlet Waduk Gajah Mungkur Wonogiri. *Bioteknologi*. 10(2) : 43–50.
- Sukmono, T., Solihin, D. D., Rahardjo, M. F. and Affandi, R. 2013. Iktiofauna di perairan hutan tropis dataran rendah, Hutan Harapan Jambi. *Jurnal Iktiologi Indonesia*. 13(2) : 161–174.
- Sukmono, T. and Margaretha, M. 2017. Ikan Air Tawar di Ekosistem Bukit Tigapuluh. Yayasan Konservasi Ekosistem Hutan Sumatera & Frankfurt Zoological Society, Jambi. 104 pp
- Tobes, L., Gaspar, S., Oscoz, J. and Miranda, R. 2016. Diagnosing stream ecosystem integrity in the Ordesa-Vinamala Biosphere Reserve, central Spanish Pyrenees. *Journal Applied Ichthyology*. 32 : 229–239.
- Thai, B. T., Si, V. N., Phan, P. D. and Austin, C. M. 2007. Phylogenetic evaluation of subfamily classification of the Cyprinidae focusing on Vietnamese species
- Phylogenetic evaluation of subfamily classification of the Cyprinidae focusing on Vietnamese species. *Aquatic Living Resources*. 20 : 143–153.
- Theurkar, S. V, Takalakar, D. L., Jadhav, S. S. and Pawar, R. M. 2013. Diversity and Composition of Fishes of Chaskaman Dam, Rajgurunagar, part of Northern Western Ghats, Pune, MS, India. *Research Journal of Animal, Veterinary and Fishery Sciences*. 1(1) : 7–10.
- Thomas, J., S. Venu and Kurup, B.M. 2003. Length- weight relationship of some deep-sea fish inhabiting the continental slope beyond 250m depth along the West Coast of India. *NAGA, World Fish Center Quaterly*, 26(2) : 17-21.
- Umar, C. and Sulaiman, P. S. 2013. Status Introduksi Ikan dan Strategi Pelaksanaan Secara Berkelanjutan Di Perairan Umum Daratan Di Indonesia. *Jurnal Kebijakan Perikanan Indonesia*. 5(2) : 113–120.
- Wahyuni, T. T. and Zakaria, A. 2018. Keanekaragaman Ikan di Sungai Luk Ulo Kabupaten Kebumen. *Biosfera*. 35(1) : 23–28.
- Wibowo, A., Affandi, R., Soewardi, K. and Sudarto. 2010. Pengelolaan Sumber Daya Ikan Belida (*Chitala lopis*) Di Sungai Kampar, Provinsi Riau. *Jurnal Kebijakan Perikanan Indonesia*. 2(2) : 79–89.
- Zulfikri, A., Umroh, and Utami, E. 2016. Pengaruh Aktivitas Tambang Apung terhadap Keanekaragaman Ikan di Perairan Sungai Pakil, Bangka. *Akuatik, Jurnal Sumberdaya Perairan*. 10(1) : 42–46.
-

# Freshwater fishes of Gegas Dam, South Sumatra Indonesia: composition and diversity

---

ORIGINALITY REPORT

---

4%

SIMILARITY INDEX

---

MATCH ALL SOURCES (ONLY SELECTED SOURCE PRINTED)

---

★www.isca.me

Internet

1%

---

EXCLUDE QUOTES ON

EXCLUDE SOURCES < 1%

EXCLUDE BIBLIOGRAPHY ON

EXCLUDE MATCHES < 10 WORDS