

THE ANALYSIS OF COVID-19 PREVENTION PRACTICES ON HEALTH CARE WORKERS IN THREE HOSPITALS IN PALEMBANG

By Rico Sitorus

THE ANALYSIS OF COVID-19 PREVENTION PRACTICES ON HEALTH CARE WORKERS IN THREE HOSPITALS IN PALEMBANG

Tamarakha Yumna¹, Nur Alam Fajar^{2*}, Rico Januar³

^{1,2,3} Faculty of Public Health, Universitas Sriwijaya, Jl. Palembang-Prabumulih KM 32 Ogan Ilir, 30662

*Correspondence Author: nuralamfajar@fkm.unsri.ac.id; +62 811-7888-060

ABSTRACT

The COVID-19 pandemic is a global concern. In this case, Health Care Workers (HCW) are at the front lines in dealing with the pandemic as they provide diagnosis, care, and treatment. However, their duties have directly placed them at the most risk of infection. Mitigation and risk reduction are crucial for safeguarding HCWs' health as it prevents the spread of COVID-19. This research aimed to find out the practice of COVID-19 prevention and the correlation among practice and knowledge, attitudes, and personality traits of HCW in Palembang. A cross-sectional study was conducted between May and July 2021 on 255 HCW who have direct contact with patients. The number of respondents per hospital is taken proportionally according to the portion of HCW in each selected hospital. The instrument consisted of demographic variables, knowledge, attitudes, and practices. It proves that it is reliable with Cronbach's alpha value of 0.7. The researchers used descriptive statistics, chi-square, and logistic regression. Overall, 65.2% of participants had good knowledge, 60% had a positive attitude, and 53.3% had good practice. Education (p-value 0,022) and attitude (p-value 0,000) of HCW were all influencing factor in practice. HCW who have a negative attitude are 5,413 times more to have bad practices than those who have a positive attitude (p-value=0.000;OR=5,413). However, training for HCW and further research are recommended to analyze the availability of personal protective equipment, adequate facilities, hospital policies, or other factors that can influence and motivate the practice of HCW.

Keywords: COVID-19, Practice, Health Care Workers

ABSTRAK

Pandemi COVID-19 merupakan ancaman global, tenaga kesehatan berada pada garis terdepan dalam melakukan diagnosis, perawatan, dan pengobatan, sehingga tenaga kesehatan berada pada risiko terinfeksi tertinggi. Mitigasi dan pengurangan risiko sangat penting untuk melindungi kesejahteraan tenaga kesehatan dan mengurangi penyebaran COVID-19. Penelitian ini bertujuan untuk menganalisis praktik terhadap pencegahan COVID-19 dan hubungan antara praktik dengan pengetahuan, sikap serta karakteristik tenaga kesehatan di Palembang. Penelitian cross-sectional dilakukan antara Mei dan Juli 2021, di antara 255 tenaga kesehatan yang melakukan kontak langsung dengan pasien. Jumlah responden per rumah sakit diambil secara proporsional sesuai porsi tenaga kesehatan pada masing-masing rumah sakit terpilih. Instrumen yang terdiri dari karakteristik demografi, pengetahuan, sikap, dan praktik, terbukti reliabel dengan nilai cronbach's alpha 0,7. Analisis deskriptif, chi-square, dan regresi logistik dilakukan. Secara keseluruhan, 65,2% memiliki pengetahuan yang baik, 60% memiliki sikap positif, dan 53,3% memiliki praktik yang baik. Pendidikan (nilai-P 0,022) dan sikap tenaga kesehatan (nilai P 0,000) memengaruhi faktor perilaku. Tenaga Kesehatan yang memiliki sikap negatif berisiko 5,413 kali lebih besar untuk memiliki praktik yang buruk dibandingkan dengan sikap positif (nilai P 0,000; OR 5,413). Namun, pelatihan bagi petugas kesehatan dan penelitian lebih lanjut disarankan untuk menganalisis ketersediaan alat pelindung diri, fasilitas yang memadai, kebijakan rumah sakit, atau faktor lain yang dapat mempengaruhi dan memotivasi praktik petugas kesehatan.

Kata Kunci: COVID-19, Perilaku, Tenaga Kesehatan.

Introduction

COVID-19 is an acute infectious disease caused by the severe acute respiratory syndrome Coronavirus 2 (SARSCoV2). This virus infects the respiratory tract, causing respiratory failure or even human death.¹⁻³ Starting from February 10th, 2021, more than 106 million cases and 2,325,000 deaths have been reported worldwide. The number is increasing day by day. There are more than 1.17 million confirmed

cases and 31,976 deaths in Indonesia. South Sumatra Province (marked as a red zone) is the province with the seventh-highest number of confirmed cases in Indonesia, with 14,857 cases and 719 deaths.⁴

In order to understand about how, when, and in what types of settings SARS-CoV-2 transmission is essential to developing the effectivity public health and infection prevention and control measures to stop the spread of this virus. The spread of SARSCoV2 occurs through contact and droplets (through infected secretions such as saliva and respiratory secretions or droplets from an infected person that come out due to coughing, sneezing, talking, or singing), air transmission (infectious substances such as z over long distances and time in the Air floating) and the spread of pollutants (through contaminated surfaces).⁵⁻⁷

OSHA (Occupational Safety and Health Administration) divided work tasks into four risk exposure levels were HCW representing the highest exposure risk.⁸ The number of infections and deaths of health workers has not been announced and remains unclear. Meanwhile, Bandyopadhyay's systematic review showed that infected health workers reached 14,058 (9.2%) cases. In addition, 343 of the 1,413 cases (24.3%) were cases of health workers who died caused by COVID-19.⁹ A survey of 37 countries showing data between July 22nd and August 15th, 2020, reported that the virus has infected 299,155 health workers with 2,736 deaths.¹⁰ The Indonesian Doctors Mitigation Team revealed that the ongoing tragedy could continue and increasing the death rate. It was around 101 doctors have died due to COVID-19 reported on August 31st, 2020.¹¹ Thus, the total of health workers who died because of COVID-19 reached 228 people.¹²

Mitigation and risk reduction are critical to protect the welfare of health workers and prevent the spread of COVID-19. It can be shows by strengthening the COVID-19 prevention, health, and safety protocols to maintain the safety of health workers further. WHO recommends preventing spread through the use of personal protective equipment (PPE), hand hygiene measures, implementation guidelines in health care facilities, and infection prevention and control education and training.¹³

Health workers are at the forefront of dealing with the COVID-19 pandemic. At this point, it puts them at the highest risk of exposure to pathogens, long working hours, psychological stress, fatigue, professional burnout and stigma, and physical violence. The lack of understanding of the disease among the HCWs can lead to an identification and treatment delay leading to the rapid spread of infections. The guidelines for health care professionals and online update courses have been developed by OMS, CDC, The Ministry of Health RI, and several government organizations in different countries to promote knowledge and prevention strategies.¹⁴

The study by Yupartini L et al. in Serang stated that most health care workers experience stress during work. Daily stress, compared to traumatic events, is a predictor factor that can have a more significant impact on mental health. The physiological stress response prolonged duration is a risk factor that can cause the disease.¹⁵ In addition, the study in Yogyakarta revealed that some of the respondents did not consume additional food and multivitamins to increase endurance. Not only about consuming food, but also there are

still quite a few who have not complied with the use of PPE, hand-washing, or aseptic procedures before/after the process or examining patients both when dealing with COVID-19 patients, non-COVID-19 and when performing aerosol-generating actions.¹⁶

It is necessary to pay attention to knowledge and attitudes about COVID-19.¹⁷ Study by Shariff FO et al. in Banyuasin showed a reasonably good level of knowledge and practice. It also offers a good level of attitude towards doctors about the COVID-19 pandemic. Their study suggested that the years of experience is another factor that influences practice besides knowledge and attitude.¹⁸ Furthermore, studies conducted by Giao et al. found a sufficient knowledge and a positive attitude toward COVID-19 in health workers.¹⁹ On the other hand, Bhagavathula, in her study, revealed that most of the health care workers had a lack of knowledge about COVID-19 globally.²⁰ Sufficient knowledge will support an optimistic attitude and proper practice during work so that the risk of infection will be reduced.¹⁷ It is important to know the correlation between knowledge, attitudes, and characteristics of health workers with type of work, age, gender, education, work experience, and training, as well as their impacts on health worker compliance with COVID-19 prevention practices. Therefore, the purpose of this study was to analyze the prevention practices of COVID-19 by health workers in Palembang.

Method

Study Design and Site

The cross-sectional study was conducted in the fourth week of May 2021 to July 2021 at Siti Fatimah Hospital of South Sumatra (Provincial Hospital), Gandus Palembang Hospital (Regency Hospital), and Sriwijaya Hospital Palembang (Private Hospital). The sampling technique that used in this study is purposive sampling.

Study Population

Health care workers who directly contacted with patients (doctors, dentists, midwives, nurses, and analysts), and also worked in the mentioned hospitals were taken as the study population. The number of respondents per hospital was taken proportionally according to the portion of health care workers in each hospital until the minimum number of samples was sufficient. Respondents were adjusted by inclusion and exclusion criteria. A total of 255 health-workers agreed to fill out the questionnaire used in this study.

Questionnaire Design

In this study, the IDI guidelines from the WHO, the Decree of the Minister of Health of the Republic of Indonesia, and the Mitigation Doctors Team, have been reviewed as well as the literature on COVID-19. On the other hand, we also revised the previous studies that tested effectiveness. We have developed a

questionnaire consisting of four parts: (I) demographics, (II) knowledge and (III) attitudes about COVID-19, and (IV) practices of health workers. The validation was using a pilot study on a small sample of 30 health care workers who were not included in this study. Accordingly, the revised questionnaire proved reliable, with a Cronbach's alpha value of 0.7.

We created 12 questions about ethicology, the period of incubation, transmission, symptoms, prevention, and diagnosis of COVID-19 to facilitate an exhaustive assessment of the health-workers' knowledge. HCW attitudes were assessed with reference to 11 articles. Individually, one by one of them were asked about their felt as HCW, meanwhile the answers are divided into categories based on the Likert scale: (1) strongly agree, (2) agree, (3) disagree, and (4) strongly disagree with the favorable and unfavorable question. Our study has evaluated HCW practices through 11 questions with four options: (1) never, (2) occasional, (3) always with a favorable and unfavorable question.

Data Analyses

This study evaluated the KAP of health care workers based on a set of specific questions. The knowledge was classified in good and poor level. Knowledge, attitude, and practice were divided by the median. If the total is equal or higher than the median, then it will be classified as a positive attitude and vice versa. Practice can be good or bad, as well as attitudes that also can be good or bad. In this research, univariate, bivariate, and multivariate analysis data were used. The univariate analysis aimed to describe each variable. Bivariate analysis, in this case, used chi-square and analysis of variance, $p < 0.05$. Multivariate analysis was performed by using logistic regression analysis. This aims to determine the most important characteristic factors for the prevention of COVID-19 in health care workers.

The ethics committee of Sriwijaya University School of Public Health and Health Research approved this study (code: 153/UN9.FKM/TU.KKE/2021). We explained the purpose of this research at first to the participants before they deliberately participated in it.

Results

Based on the univariate analysis approach, the results obtained from data analysis are shown in Table 1. Meanwhile, based on bivariate and multivariate analysis the results are shown in Table 2.

Table 1. Frequency Distribution Based on Health Care Workers' Characteristics, knowledge, attitude, and practice (N= 255)

Variable	Frequency	Percentage (%)
Managing COVID-19 patients		
Yes	149	58,4
No	106	41,6

Occupation		
Doctor	54	21,2
Dentist	10	3,9
Midwife	39	15,3
Nurse	122	47,8
Medical analyst	30	11,8
Age-group (years)		
≤35	215	84,3
>35	40	15,7
Gender		
Male	58	22,7
Female	197	77,3
Education		
≤ Bachelors	228	89,4
> Bachelors	27	10,6
Year of experience		
≤5	181	71
>5	74	29
Source of information on COVID-19*		
WHO	196	76,6
Indonesian Government Policy	176	69,2
Social media e.g., WhatsApp, Facebook	186	73
News media e.g., TV, radio, newspaper	159	62,4
Journals	87	34,6
Others (Medicine Magazine)	1	0,4
Knowledge		
Poor	89	34,9
Good	166	65,1
Attitude		
Negative	102	40
Positive	153	60
Practices		
Poor	119	46,7
Good	136	53,3

*Multi-select questions

9
 T Table 1 shows that the characteristics of health care workers. Most respondents were managing COVID-19 patients (58.4%). The highest percentage of profession were nurses (47.8%). Furthermore, the proportion of respondents aged ≤35 years old were 84,3% and most of them were female (77,3%). More than half of the respondents (89.4%) had bachelor's or lower studies. Many of them have equal or less than 5 years of experience (71%). The main sources of COVID-19 information were WHO (76.6%). Most of respondents have good knowledge (65.1%), positive attitude (60%), and good practice (53.3%).
 45

Table 2. Bivariate Analysis of Factors associated with Practices of Preventing COVID-19 among Health Care Workers

Variables	Practice toward COVID-19 prevention				p-value	PR	95% CI
	Poor		Good				
	n	%	n	%			
Age-group (years)							
≤35	105	48,8	110	51,2	0,107	1,395	0,895-2,175
>35	14	35	26	65			
Total	119	46,7	136	53,3			
Gender							
Male	29	50	29	50	0,563	1,094	0,812-1,476
Female	90	45,7	107	54,3			
Total	119	46,7	136	53,3			
Managing COVID-19 patients							
Yes	72	48,3	77	51,7	0,530	1,090	0,832-1,428
No	47	44,3	59	55,7			
Total	119	46,7	136	53,3			
Education							
≤ Bachelors	112	49,1	116	50,9	0,022*	1,895	0,988-3,634
> Bachelors	7	25,9	20	74,1			
Total	119	46,7	136	53,3			
Year of experience							
≤5	88	48,6	93	51,4	0,328	1,161	0,854-1,578
>5	31	41,9	43	58,1			
Total	119	46,7	136	53,3			
Knowledge							
Poor	38	42,7	51	57,3	0,352	0,875	0,657-1,166
Good	81	48,8	85	51,2			
Total	119	46,7	136	53,3			
Attitude							
Negative	72	70,6	30	29,4	0,000*	2,298	1,756-3,007
Positive	47	30,7	106	69,3			
Total	119	61,6	136	53,3			

*Significant

According to Table 2, it shows that no significant correlation between characteristics and practice toward COVID-19 prevention, except for education. Respondents with the same level of education as bachelor's degree had 49.1% higher bad practices, than those with postgraduate education. Meanwhile, 48.8% of respondents (aged less than or equal to 35 years) had poor COVID-19 prevention practices higher than respondents aged over 35 years. In addition, 50% of respondents are males who have bad practices. The same results were also found in 45.7% of women. In total of 48.3% respondents are treat for COVID-19 patients have bad practices. Furthermore, a total of 48.6% of respondents who have experience equal to or less than 5 years have higher bad practices than respondents who have experience of more than 5 years.

Furthermore, the bivariate analysis shows that 42,7% of respondents have poor knowledge and poor practices lower than respondents who have good knowledge. The chi-square test showed no significant

correlation between knowledge and practice toward COVID-19 prevention. While 70,6% of respondents have negative attitude and poor practice much higher than respondents who have positive attitude, there is a significant correlation between attitude and practice.

Table 3. Multivariate Analysis with Multiple Logistic Regression

Variabel	B	S.E	Sig.	Exp(B)	95% CI for Exp (B)
Sikap	1,689	0,279	0,000	5,413	3,132-9,355
Constant	-2,564	0,469	0,000	0,077	

Variables that can be used for multivariate analysis are age, education, and attitude. This is because the p-value found is below 0.25. The final multivariate analysis model shows in Table 3 that there is only one independent variable that affects the dependent variable, namely the attitude variable. The Prevalence Ratio (PR) value shows that health workers who have a negative attitude have 5,413 times bad practices compared to health workers who have a positive attitude. The confidence interval in this case is 3.132-9.355. Researchers believe that negative attitudes are a risk factor for COVID-19 prevention practices in health care workers in 95% of the general population.

Discussion

OSHA (Management of Manpower and Occupational Safety) stated that the occupations with the most risk of being exposed to COVID-19 are health workers.⁸ Thus, it is necessary to notice the prevention of COVID-19 as early as possible.¹³ The practices of prevention could be affected by knowledge and attitudes about COVID-19. In-depth knowledge supports optimistic attitudes and correct work practices might help to avoid the risk of infection.¹⁷ This study showed a correlation between education, attitude, and practice. However, it was found that the most dominant factor affecting to COVID-19 prevention among health care workers was attitude.

The current study shows that most of the participants get the COVID-19 information from WHO. Olum R et.al suggested, that international health organizations WHO could be the source information on COVID-19.¹⁴ Yet, this study has not analyzed the knowledge based on the source of information. More than half of health workers had good knowledge (65,1%). This study has results that are no significant different from the research in Uganda, which showed that 69% of respondents were considered to have good knowledge.¹⁴ According to a study by Haghghi K et al, it was found about 72.8% of health care workers had sufficient knowledge²¹, while a study by Sidras shows 77.5% of health workers had good knowledge.²² Furthermore, the study by Maleki S et.al. shows 99% of respondents had excellent knowledge levels on the disease modes of transmission.²³ In line with the study conducted by Rizki,²⁴ it was stated that most Indonesian health care workers have good knowledge. In contrast, a study by Bhagavathula AS et.al. revealed about 61% of health care workers had poor knowledge of the virus transmission.²⁰

Overall, 60% of health care workers had a positive attitude in handling the COVID-19. This result is similar to another study conducted in Nepal, it was found that around 53.4% of health care workers had a positive attitude in handling COVID-19. This is similar because only half of the health workers believe that Nepal can win the fight against COVID-19. Besides the support from family members, they are still lack self-confidence as anxiety and anxiety in the workplace increase.¹⁷ This is in contradict with a study conducted in Vietnam where more than 90% of people responded positively to COVID-19 and only three-quarters of respondents have no idea if they might be infected with the disease. In addition, they are also think that one of their family members might also be infected and quarantined for suspected infection.¹⁹ This study found that the attitudes and practices of health workers were significantly lower than those who cared for COVID-19 patients. This may be because 44.3% of respondents are afraid to treat suspected as COVID-19 patients. It was confirmed by a study in Jordan which showed that anxiety, depression, and stress are increase among health care workers during the COVID-19 pandemic.²⁵ Men are more likely to have higher levels of attitudes than women, but Sahar J found women have significantly higher levels of attitudes.²⁶ However, other studies found no significant difference between attitude and gender.^{14,17,27} The experience of 1-5 years shows that the higher attitude occurs because of the longer experience in dealing with emergencies, it will increase the confidence and optimism in health workers.¹⁷

More than half of the respondents in this study had good practices (53,3%). The percentage is lower than the results of study conducted by Olum R et.al. and Limbu et.al. which showed that 74% and 81,5 % of the respondents had good practices, respectively.^{14,17} It is consistent with an Iranian study where 67.72 wrote best practices.²¹ A study in China showed that 85% of health care workers are afraid of being infected in the workplace. Health care workers must also deal with various emergencies and are very likely to be infected.²⁸

The correlation between education or attitudes towards COVID-19 prevention practices was found to be significant. Specifically, attitude is an essential factor that correlates with COVID-19 prevention practices. The prevalence ratio of the attitude variable is 5.413 and indicates that health care workers that have negative attitudes also have 5.413 times less practices than health care workers with positive attitude. Researchers believe that negative attitudes are the most risk factor for COVID-19 prevention practices in health care workers in 95% of the general population. In line with another study by Limbu et.al which shows that practices were significantly correlated with attitude.¹⁷ In contrast to other studies, based on this study, the factors related to the practice of preventing COVID-19 were age 40 years or older with D3 as the last education.¹⁴ Study by Zhang et.al. shows that work experience and job category were some risk factors that influences health care workers' practice.²⁸

There are many factors may cause differences in results between studies, such as time, country, health workforce composition, government policies, and information sources. However, this is the first

study that evaluates the COVID-19 prevention on Palembang health workers'. According to the respondents, it is found that 26.3% of the respondents never participate in the regular COVID-19 training organized by the hospital and 40% rarely participate. Thus, the health system needs to recognize the reason of an intervention plan to increase knowledge alone is not enough. Immediate action should be taken to promote correct practices regarding infectious diseases, especially during the COVID-19 pandemic. On the other hand, the worst-hit areas always have the necessary equipment is also needed to be improved, such as: the provision of PPE, a good working environment, adequate changing rooms, and the role of the government in coordinating efforts to ensure.^{24,29} It is recommended to standardize the data upload to regional and national authorities and creates a common basis for scientific comparisons of the effects of a pandemic on which new strategies should also be improved to prevent the spread of COVID-19.³⁰

The limitation of this study is a larger population needed. Therefore, it could be compared between hospitals that neither treat nor treat COVID-19 patients. Several environmental factors that influenced the practice of health care workers were not discussed in this study. The examination in this study was only shows based on respondents' answers and did not observe the practice of health workers directly. This research is also did not focus on health care facilities and policies that might support the practice of health care workers. Thus, further research with larger samples is suggested by using mixed methods and direct observations, on the practices carried out by health workers. Training and periodic evaluations can also be conducted to improve knowledge, attitudes, and practices in COVID-19 prevention. The training that can be carried out is such as socializing the steps for using PPE properly, the right way to use PPE, the transmission of COVID-19 prevention, procedures for handling patients, and COVID-19 suspects.

Conclusion

The practices by health care workers in Palembang for the prevention of COVID-19 were largely associated with their knowledge, attitude, and some characteristics (age, years of experience, and involvement of dealing with COVID-19). Older health care workers are 3.2 times more likely to apply best practices than younger health care workers. But it is recommended that further training and research on health care workers be conducted. It aims to analyze the availability of PPE, adequate facilities, hospital policies, or other factors that might influence health care workers, and motivate them to practice.

Acknowledgment:

We would like to thank the staff and health workers of Siti Fatimah Hospital, South Sumatra, Gandus Hospital Palembang, and Sriwijaya Hospital Palembang as well as all respondents who have provided assistance and support for this research.

Funding:

15

The authors should state that they have no funding for the research.

Conflict of Interest:

The authors declare that they have no conflict of interest.

References

1. Ather A, Dds BDS, Bds BP, Ruparel NB, Dds MS, Diogenes A, et al. Coronavirus Disease 19 (COVID-19): Implications for Clinical Dental Care. *J Endod.* 2020;19:1–12.
2. Israfil, Wiliyanarti PF, Selasa P. Literature Review : Risk of Death in COVID-19 Patients. *Unnes J Public Heal.* 2020;9(2):141–7.
3. Zhu N, Zhang D, Wang W, Li X, Yang B, Song J, et al. A Novel Coronavirus from Patients with Pneumonia in China, 2019. *N engl j med.* 2020;382:727–33.
4. WHO. WHO Coronavirus Disease (COVID-19) Dashboard. 23 January 2021. 2021.
5. WHO. Transmission of SARS-CoV-2: implications for infection prevention precautions. 2020.
6. Li Q, Guan X, Wu P, Wang X, Zhou L, Tong Y, et al. Early Transmission Dynamics in Wuhan, China, of Novel Coronavirus–Infected Pneumonia. *N Engl J Med.* 2020;382(13):1199–207.
7. Sahin AR, Erdogan A, PM A, Dineri Y, AY C. 2019 Novel Coronavirus (COVID-19) Outbreak: A Review of the Current Literature. *Eurasian J Med Oncol.* 2020;4(1):1–7.
8. OSHA. Worker Exposure Risk to COVID-19. 2020.
9. Bandyopadhyay S, Baticulon RE, Kadhum M, Alser M, Ojuka DK, Badereddin Y, et al. Infection and mortality of healthcare workers worldwide from COVID-19 : a systematic review. *BMJ Glob Heal.* 2020;1–11.
10. Erdem H, Lucey DR. Healthcare worker infections and deaths due to COVID-19 : A survey from 37 nations and a call for WHO to post national data on their website. *Int J Infect Dis.* 2021;102:239–41.
11. Tim Mitigasi Dokter IDI. Pedoman standar perlindungan dokter di era COVID-19. 2020. 1–91 p.
12. Halimatu H, Amelia T. HEALTHCARE WORKERS SECURITY : Jaminan, Regulasi, dan Sanksi. *Khatulistiwa Law Rev.* 2020;1(2):38–9.
13. WHO. Prevention , identification and management of health worker infection in the context of COVID-19. 2020.
14. Olum R, Bongomin F. Coronavirus Disease-2019 : Knowledge , Attitude , and Practices of Health Care Workers at Makerere University Teaching Hospitals, Uganda. *Front Public Heal.*

2020;8(181):1–9.

15. Yupartini L, Rustiawati E, Sulastris T. Stres Kerja dengan Perilaku Caring Petugas Kesehatan dalam Penanganan Pasien pada Masa Pandemi Covid-19. *J Keperawatan Silampari*. 2021;4(2):683–9.
16. Murdiyanto J, Suryadi H, Nuryati R, Wijaya T. Survei Mitigasi Risiko Covid-19 Pada Tenaga Kesehatan. *J Kesehat Kusuma Husada*. 2021;12(2):155–63.
17. Limbu DK, Piryani RM, Id AKS. Healthcare workers' knowledge, attitude and practices during the COVID-19 pandemic response in a tertiary care hospital of Nepal. *PLUS ONE*. 2020;12:7–13.
18. Shariff FO, Ladyani F, Priliansyah AL. Gambaran Pengetahuan Sikap Dan Perilaku Dokter Ptt Terhadap Pandemi Covid-19 Di Puskesmas Kabupaten Banyuwangi Tahun 2020. *MAHESA Malahayati Heal Student J*. 2021;1(2):113–21.
19. Giao H, Thi N, Han N, Khanh T Van, Ngan VK, Tam V Van, et al. Knowledge and attitude toward COVID-19 among healthcare workers at District 2 Hospital, Ho Chi Minh City. *Asian Pacific J Trop Med*. 2020;13(6):260–5.
20. Bhagavathula AS, Aldhaleei WA, Rahmani J, Ain A, Dhabhi A. Knowledge and Perceptions of COVID-19 Among Health Care Workers: Cross-Sectional Study. *JMIR Public Heal Surveill*. 2020;6(2):1–9.
21. Haghghi FK, Kouhi P, Amini M, Mohammadkarimi V, Hosseini SA, Moosavi M, et al. Knowledge, Attitude, and Practice Toward COVID-19 Among Healthcare Workers in Shiraz, Iran. *Shiraz E-Med J*. 2020;21(12):1–8.
22. Fadli F, Ahmad AS, Safruddin S, Sumbara S, Baharuddin R. Anxiety of Health Workers in the Prevention and Management of Covid-19 in Sidrap Regency. *Unnes J Public Heal*. 2020;9(2):91–7.
23. Maleki S, Najafi F, Farhadi K, Fakhri M, Hosseini F, Naderi M. Knowledge, Attitude and Behavior of Health Care Workers in the Prevention of COVID-19. *Res Sq*. 2020;1–17.
24. Rizki SA, Kurniawan J, Budimulia P, Sylvanus P, Alexandra A, Sinaga TD, et al. Knowledge, Attitude, and Practice in Indonesian Health Care Workers Regarding COVID-19. *Asia-Pacific J Public Heal*. 2021;33(5):662–4.
25. Alnazly E, Khraisat OM, Al-Bashairah AM, Bryant CL. Anxiety, depression, stress, fear and social support during COVID-19 pandemic among Jordanian healthcare workers. *PLoS One*. 2021;16(3 March).
26. Sahar J, Kiik SM, Wiarsih W, Rachmawati U. Coronavirus disease-19: Public health nurses' knowledge, attitude, practices, and perceived barriers in Indonesia. *Open Access Maced J Med Sci*. 2020;8(T1):422–8.
27. Saqlain M, Munir MM, Rehman SU, Gulzar A, Naz S, Ahmed Z, et al. Knowledge, attitude, practice and perceived barriers among healthcare workers regarding COVID-19: a cross-sectional survey

- from Pakistan. *J Hosp Infect.* 2020;105(3):419–23.
28. Zhang M, ZHOU M, TANG F, WANG Y, NIE H, Zhang L, et al. Knowledge, attitude, and practice regarding COVID-19 among healthcare worker in Henan, China. *J Hosp Infect.* 2020;(January):19–21.
 29. Ranney M, Griffeth V, Jha A. Critical Supply Shortages — The Need for Ventilators and Personal Protective Equipment during the Covid-19 Pandemic. *TheNew Engl J Med.* 2020;41(1):1969–73.
 30. Mendonça-Galaio L, Sacadura-Leite E, Raposo J, França D, Correia A, Lobo R, et al. The COVID-19 impact in hospital healthcare workers: Development of an occupational health risk management program. *Port J Public Heal.* 2021;38(1):26–31.

THE ANALYSIS OF COVID-19 PREVENTION PRACTICES ON HEALTH CARE WORKERS IN THREE HOSPITALS IN PALEMBANG

ORIGINALITY REPORT

18%

SIMILARITY INDEX

PRIMARY SOURCES

1	sites.kowsarpub.com Internet	75 words — 2%
2	Www.esp.org Internet	42 words — 1%
3	Zebader Walle, Gete Berihun, Awoke Keleb, Daniel Teshome, Leykun Berhanu. "COVID-19 Prevention Practices and Determinant Factors Among Healthcare Professionals Working in Hospitals of South Gondar Zone, Northwestern Ethiopia", Journal of Multidisciplinary Healthcare, 2021 Crossref	37 words — 1%
4	www.frontiersin.org Internet	37 words — 1%
5	www.researchsquare.com Internet	33 words — 1%
6	journals.plos.org Internet	32 words — 1%
7	garuda.ristekbrin.go.id Internet	27 words — 1%

8 Carl J. Heneghan, Elizabeth A. Spencer, Jon Brassey, Annette Plüddemann et al. "SARS-CoV-2 and the role of orofecal transmission: a systematic review", F1000Research, 2021 25 words — 1%
Crossref

9 DIPAK PRASAD UPADHYAYA, Rajan Paudel, Daniel J Bromberg, Dilaram Acharya et al. "Frontline healthcare workers' knowledge and perception of COVID-19 and willingness to work during the pandemic in Nepal: a nationwide cross-sectional web-based study", Cold Spring Harbor Laboratory, 2020 22 words — 1%
Crossref

10 storage.googleapis.com 19 words — < 1%
Internet

11 www.jmir.org 19 words — < 1%
Internet

12 Amina M. Al Marzouqi, Michael E. Otim, Heba H. Hijazi, Leena S. Kehail, Ramsha A. Kamal. "Knowledge, Attitudes, and Practice Related to COVID-19 Among Healthcare Workers in the United Arab Emirates: A Cross-sectional Study.", Research Square Platform LLC, 2021 18 words — < 1%
Crossref Posted Content

13 en.wikipedia.org 17 words — < 1%
Internet

14 Shiferaw Gelchu Adola, Girish Dagavi, Sarah Ezhil Kelna Edwin, Takala Utura, Udessa Gemedo, Pandiarajan Kasimayan. "Assessment of factors affecting practice towards COVID-19 among health care workers in health care facility of West Guji Zone, South Ethiopia, 2020", Pan African Medical Journal, 2021 16 words — < 1%
Crossref

15	www.e-mjm.org Internet	16 words — < 1%
16	downloads.hindawi.com Internet	15 words — < 1%
17	Osama Ahmed, Muhammad Sohaib Asghar, Muhammad Nadeem Khurshaidi, Farah Yasmin et al. "Provision of Surgical Services to COVID-19-Infected Patients at a Tertiary Care Center in Pakistan: A One-Year Clinical Review of the Year 2020 in General Surgery Department", Cureus, 2021 Crossref	13 words — < 1%
18	Giuliana Bontà, Guglielmo Giuseppe Campus, Maria Grazia Cagetti. "COVID-19 Pandemic and Dental Hygienists in Italy: A Questionnaire Survey.", Research Square, 2020 Crossref Posted Content	12 words — < 1%
19	www.karger.com Internet	12 words — < 1%
20	Sagar Pandey, Sujan Poudel, Ashok Gaire, Ritu Poudel, Prabin Subedi, Jyoti Gurung, Rituraj Sharma, Jeevan Thapa. "Knowledge, attitude and reported practice regarding donning and doffing of personal protective equipment among frontline healthcare workers against COVID-19 in Nepal: A cross-sectional study", PLOS Global Public Health, 2021 Crossref	11 words — < 1%
21	link.springer.com Internet	11 words — < 1%
22	medcraveonline.com Internet	11 words — < 1%

-
- 23 www.hindawi.com 11 words — < 1%
Internet
-
- 24 Alemu Degu Ayele, Gedefaye Nibret Mihretie, Habtamu Gebrehana Belay, Adanech Getie Teffera et al. "Knowledge and Practice to Prevent Against Corona Virus Disease (COVID-19) and Its Associated Factors Among Pregnant Women in Debre Tabor Town Northwest Ethiopia: a Community Based Cross-Sectional Study.", Research Square, 2020 10 words — < 1%
Crossref
-
- 25 Prasojo Pribadi, Lolita Lolita, Rayi Citra Ayu Pangestuti, Heni Lutfiyati, Silvia Mareti, Azis Ikhsanudin. "Knowledge and quality of life among Indonesian students during the COVID-19 pandemic", International Journal of Public Health Science (IJPHS), 2021 10 words — < 1%
Crossref
-
- 26 Wakgari Deressa, Alemayehu Worku, Workeabeba Abebe, Muluken Gizaw, Wondwosson Amogne. "Risk perceptions and preventive practices of COVID-19 among healthcare professionals in public hospitals in Ethiopia", Cold Spring Harbor Laboratory, 2020 10 words — < 1%
Crossref Posted Content
-
- 27 lib.atmajaya.ac.id 10 words — < 1%
Internet
-
- 28 oamjms.eu 10 words — < 1%
Internet
-
- 29 Benito Yáñez-Araque, Sagrario Gómez-Cantarino, Santiago Gutiérrez-Broncano, Víctor-Raúl López-Ruiz. "Examining the Determinants of Healthcare Workers' Performance: A Configurational Analysis during COVID-19" 9 words — < 1%

30 Jesús Muyor-Rodríguez, Francisco Caravaca-Sánchez, Juan Sebastián Fernández-Prados. "COVID-19 Fear, Resilience, Social Support, Anxiety, and Suicide among College Students in Spain", International Journal of Environmental Research and Public Health, 2021
9 words — < 1%
Crossref

31 Repository.umy.ac.id
Internet
9 words — < 1%

32 Yodang Yodang, Stefanus Mendes Kiik, Achmad Fauji, Hamka Hamka et al. "Knowledge, attitudes, and practices of Indonesian residents regarding COVID-19: A national cross-sectional survey", International Journal of Public Health Science (IJPHS), 2021
9 words — < 1%
Crossref

33 Zelalem Tilahun Tesfaye, Malede Berihun Yismaw, Zenebe Negash, Akeberegna Gorems Ayele. "COVID-19-Related Knowledge, Attitude and Practice Among Hospital and Community Pharmacists in Addis Ababa, Ethiopia", Integrated Pharmacy Research and Practice, 2020
9 words — < 1%
Crossref

34 dokumen.pub
Internet
9 words — < 1%

35 f1000research.com
Internet
9 words — < 1%

36 plus.mcmaster.ca
Internet
9 words — < 1%

37 Bedru Jemal, Zemedu Aweke, Simeneh Mola, Sileshi Hailu et al. "Knowledge, attitude, and practice of healthcare workers toward COVID-19 and its prevention in Ethiopia: A multicenter study", SAGE Open Medicine, 2021

8 words — < 1%

Crossref

38 Huiming Gao, Rujun Hu, Ling Yin, Xiaoli Yuan, Hao Tang, Lan Luo, Mei Chen, Di Huang, Ying Wang, Anyong Yu, Zhixia Jiang. "Knowledge, attitudes and practices of the Chinese public with respect to coronavirus disease (COVID-19): an online cross-sectional survey", BMC Public Health, 2020

8 words — < 1%

Crossref

39 Mansour Alsoghair, Mohammad Almazyad, Tariq Alburaykan, Abdulrhman Alsultan et al. "Medical Students and COVID-19: Knowledge, Preventive Behaviors, and Risk Perception", International Journal of Environmental Research and Public Health, 2021

8 words — < 1%

Crossref

40 Most. Zannatul Ferdous, Md. Saiful Islam, Md. Tajuddin Sikder, Abu Syed Md. Mosaddek, J. A. Zegarra-Valdivia. "Knowledge, attitude, and practice regarding COVID-19 outbreak in Bangladeshi people: An online-based cross-sectional study", Cold Spring Harbor Laboratory, 2020

8 words — < 1%

Crossref Posted Content

41 Syed Mohammed Basheeruddin Asdaq, Alshrari A.S, Mohd. Imran, Nagaraja Sreeharsha, Rokeya Sultana. "Knowledge, attitude and practices of healthcare professionals of Riyadh, Saudi Arabia towards covid-19: A cross-sectional study", Saudi Journal of Biological Sciences, 2021

8 words — < 1%

Crossref

42 Zelalem Desalegn, Negussie Deyessa, Brhanu Teka, Welelta Shiferaw et al. "Evaluation of COVID-

8 words — < 1%

19 related knowledge and preparedness in health professionals at selected health facilities in a resource-limited setting in Addis Ababa, Ethiopia", PLOS ONE, 2021

Crossref

-
- 43 apjtm.org Internet 8 words — < 1%
-
- 44 www.coursehero.com Internet 8 words — < 1%
-
- 45 www.dovepress.com Internet 8 words — < 1%
-
- 46 www.unicef.org Internet 8 words — < 1%
-
- 47 Robert Kaba Alhassan, Seth Owusu-Agyei, Evelyn Korkor Ansah, Margaret Gyapong. "COVID-19 vaccine uptake among health care workers in Ghana: a case for targeted vaccine deployment campaigns in the global south", Human Resources for Health, 2021
Crossref 7 words — < 1%
-
- 48 Astawus Alemayehu, Abebaw Demissie, Mohammed Yusuf, Dereje Feleke et al. "Covid-19 Prevention: Level and Determinants of Knowledge, Attitude and Practice toward WHO Recommended Covid-19 Prevention Measures Among Harar Population, Eastern Ethiopia: A Community-Based Cross-Sectional Study", SAGE Publications, 2021
Crossref Posted Content 6 words — < 1%
-
- 49 Richa Nepal, Kalyan Sapkota, Kalidas Adhikari, Pramod Paudel, Bhojraj Adhikari, Nabin Paudyal, Kiran Sapkota, Rhishikesh Nepal. "Knowledge, attitude and 6 words — < 1%

practice regarding COVID-19 among healthcare workers in Chitwan, Nepal", Research Square, 2020

Crossref Posted Content

EXCLUDE QUOTES OFF

EXCLUDE MATCHES OFF

EXCLUDE BIBLIOGRAPHY ON