

THE EFFECT OF MUNDU FRUIT (*Garcinia dulcis* (ROXB.) KURZ) METHANOL EXTRACT ON LYPHOPROTEIN PROFILE AND TRYGLISERIDE WHITE RAT (*Rattus norvegicus* L)

Arum Setiawan*, Laila Hanum* and Elvi Rusmiyanto Pancaning Wardoyo**

*Departement of Biology, Faculty of Mathematic and Natural Science, Sriwijaya University, Jl. Palembang-Prabumulih KM 32, Indralaya (OI) South Sumatera, Indonesia, 30662

**Departement of Biology, Faculty of Mathematic and Natural Science, Tanjungpura University

Jl. Prof. Hadari Nawawi, Pontianak, West Kalimantan, Indonesia 78124

Corresponding Author email: lailahanum@gmail.com

ABSTRACT

Garcinia dulcis (Roxb.) Kurz (Guttiferae) is a tropical fruit tree native to Indonesia and its known as an Asian medicinal plant used in folk medicines. In ethno pharmacology, *Garcinia dulcis* has been used in traditional medicine, such as anti-inflammatory, anti-tumor, anti-malaria and anti-bacterial. The scientific studies of biological activity of *Garcinia dulcis* on lipide profile and triglyceride are relatively measly. Therefore, the objectives of this research were to know the effect of methanol extract of *Garcinia dulcis* fruit on total cholesterol, low density lipoprotein-cholesterol, high density lipoprotein-cholesterol and triglyseride rat hyperlipidemic. These studies using *Garcinia dulcis* fruit from Banyuasin, South Sumatra. In this observation, 30 male rat from Laboratory and Research Institute for Integrated Testing, Gadjah Mada University, 2 months old were used and divided into five groups randomly. The first group was given high cholesterol and lipid diet, as a normal group. The second group was given high cholesterol, lipid diet and simvastatin 3.6 mg/kg bow/day dosage, as a simvastatin group. The third, fourth and fifth groups was given high cholesterol, lipid diet and methanol extract 1.8; 2.7 and 3.6 mg/kg bow/day dosage. Total cholesterol, low density lipoprotein-cholesterol, high density lipoprotein-cholesterol and triglyseride contains using spectrophotometry method, were observed 1 and 30 days. The differences among groups were analyzed using Anova followed by LSD to locate the differences. LSD analysis showed that *Garcinia dulcis* methanol extract treatment doses of 3.6 mg/kg bw and simvastatin lowers total cholesterol levels highest in the amount of 32.17 mg/dL and 24.1 mg/dL, significantly different compared with the treatment dose of 1.8 mg/kg bw. *Garcinia dulcis* fruit methanol extract treatment doses of 3.6 mg/kg bw and simvastatin lowers triglyceride levels by 7.22 mg/dL and 24.58 mg/dL, significantly different compared with the treatment dose of 1.8 mg/kg bw and controls. *Garcinia dulcis* fruit extract dose of 3.6 mg/dL lower low density lipoprotein-cholesterol highest of 16.45 mg/dL. *Garcinia dulcis* fruit methanol extract dose of 3.6 mg/kg bw was also able to increase the levels of high density lipoprotein-cholesterol by 5.62 mg/dL. The results showed that the fruit of *Garcinia dulcis* contain compounds that have potential as agents antihyperlipidemia and presumably through inhibition of the enzyme HMG-CoA reductase.

Key words: cholesterol, *Garcinia dulcis*, triglyseride

INTRODUCTION

Mangosteen is a plant that grow and spread in the lowland tropical forests of Southeast Asia, starting from the region of Thailand, Peninsular Malaysia and Indonesia and is known as the queen of fruits. In Indonesia, genus *Garcinia* are common in Sumatra, Java, Kalimantan and Maluku. The genus *Garcinia* (Familia: Guttiferae) is a tropical plant and has approximately 350 species, in-cluding *Garcinia dulcis*, *Garcinia bancana*, *Garcinia bor-neensis*, *Garcinia parvifolia*, *Garcinia rostrata*, and *Garcinia* spp (Sweeney, 2008). Besides eaten, this plant has been widely used by people for a variety of traditional tre-atments, such as limphatis, parotid, anti-inflammatory, anti-tumor and anti-bacterial activity on *Staphylococcus aureus*, *Shigella dysenteriae*, *Shigella flexneri*, *Escherichia coli*, *Vibrio cholerae* (Deachathai *et al.*, 2005).

The content of chemical compounds that have been found to vary widely in the genus *Garcinia* are flavonoids, xanton, triterpenoids, and quinone (Deachathai *et al.*, 2005). Flavonoid compounds that have been found in the genus *Garcinia* mostly biflavonoid types. *Garcinia scorte-chinii* fruit contains compounds biflavonoid (Sukpondma *et al.*, 2005) and kolaflavanon (Adaramoye & Muritala, 2013).

Research has been done on the genus *Garcinia* shows there biflavonoid other compounds, such as amantoflavan, agathisflavan, robustaflavan, hinokiflavan, volkensiflavan, rhusflavanon, succedaneiflavan of *Garcinia multiflora* which shows antidiabetic and antiviral activity and hepatoprotective (Lin *et al.*, 1997; Iwu *et al.*, 1990). Five compounds xanton i.e dihidroksixanton, dihidroxydes-D-garcigerrin, metilsimpoxanton, simpoxanton and

garcinixanton apparently showed antimalarial activity (Likhiwitayawuid *et al.*, 1998). Benzophenone derivative compounds, namely garcinol, isogarcinol, xantosimol, isoxantosimol and sikloxantosimol showed antibacterial activity against *Staphylococcus aureus* (Iinuma *et al.*, 1996).

Several compounds of *Garcinia xanthon* showed antioxidant activity, namely dihydroxy furan and 1,4,5-trihidroksixanton (Minami *et al.*, 1995). The compound 3-hydroxy- β -20,29,30-trinorlupan-9-on garciniellipton and Garsubellin A isolated from *Garcinia subelliptica* showed cytotoxic activity, antibiotics, and increase the activity of choline asetiltransferase (Lin *et al.*, 2012; Zhang *et al.* 2010; Asano *et al.*, 1996; Fukuyama *et al.*, 1997; Kosela *et al.*, 2000).

Until now, research on the activity of the antihyperlipidemia methanol extract of the fruit *Garcinia dulcis* from South Sumatra on lipoprotein profile and triglycerides are still rarely found. Antihyperlipidemia activity can be known through the inhibition of HMG-CoA reductase compounds. HMG-CoA reductase is an enzyme that catalyzes the conversion of 3-hydroxy-3-methylglutaryl coenzyme A (HMG-CoA) to mevalonic acid, one of the important stages in the pathway of cholesterol synthesis. Inhibition of the enzyme leads to increased density of LDL receptors in the liver cells resulting in decreased levels of cholesterol, LDL-cholesterol and triglycerides (Delgado & Remers, 1991).

Based on research conducted by Qudsiyah & Ward (2007), the methanol extract of *Garcinia dulcis* stem can reduce total cholesterol by 65% rats at levels of 3.6 mg/kg/ day. According Olaiya & Omolekan (2013), ethanol ex-tract contained in *Buchholzia coriacea* at dose

of 250 mg/ kg bw has anti-hyperlipidemia activity via decreased levels of total cholesterol, LDL-cholesterol and triglycerides hyperlipidemic mice at 45.9%, 73.5%, and 48.4 %. *Vernonia amygdalina* methanol extract dose of 200 mg/kg bw able to reduce levels of cholesterol, LDL-cholesterol and triglycerides hyperlipidemic mice by 29%, 49% and 28% (Ada-ramoye *et al.*, 2008). *Brucea javanica* seed methanol extract dose of 2.7 mg/kg bw could reduce levels of LDL-cholesterol and triglycerides were hyperlipidemic white mice by 43.8% and 32.2% (Wulandari *et al.*, 2011).

Therefore, this study aims to determine the activity of fruit extracts Mundu (*G. dulcis*) at different doses on levels of total cholesterol, triglycerides, HDL-cholesterol and LDL-cholesterol white rat (*Rattus norvegicus* L.).

MATERIAL AND METHOD

Extraction of Mundu (*Garcinia dulcis*)

Mundu (*G. dulcis*) fruit derived from Banyuasin Regency, South Sumatera Province, and has been determined by Herbarium Bogoriense LIPI, Bogor. Extraction is done by maceration using methanol pa. 250 gram samples that have been dried fruits Mundu, incorporated into maserator then add 500 mL methanol pa. Maceration is done for 3 x 24 hours. The extract obtained was concentrated by using a rotary evaporator for 3 hours.

Test Anti-hyperlipidemia

Test animals used were male Wistar rats (*Rattus norvegicus*), 2 months old with body weight of 150-250 g were obtained from preclinical Laboratory and Research Institute for Integrated Testing (LPPT) UGM. Thirty male white rats of Wistar male (*R. norvegicus*) are healthy and have normal activities were randomly divided into 5 groups with 6 replicates. The control group is the group fed a standard diet and feed high cholesterol and fat for 4 weeks; Statin group is a group fed a standard diet and feed high in fat and cholesterol and simvastatin at a dose of 3.6 mg/kg bw; Group M1, M2 and M3 are the group fed a standard diet and feed high in fat and cholesterol and methanol extract at a dose of 1.8, respectively; 2.7 and 3.6 mg/kg bw/day.

Before treatment, white male rats adapted for 1 week in the animal cages at the temperature and humidity of 50-60% 26-27°C. During the adaptation period, male rats were fed a standard 15 gr/150 gr BB mature rats and drinking water ad libitum (Gad and Chengelis, 1992). According to Anonim (1993), the composition of foods to increase levels of lipid and cholesterol of the rats per day are: egg yolks 5%, 10% beef tallow, palm oil and food standard 1% to 100%.

Determination of the dose refers to Dubey *et al.* (2005), the standard drug dose conversion for human consumption simvastatin dose to rats, at 0.018. Methanol extract dose used was 1.8; 2.7; 3.6 mg/kg rat. On days 1 and 30 rat blood drawn through orbital sinus. Before blood sampling rats were fasted for 16 hours with the aim of completing the process of lipid metabolism that comes from food (Felig & Frohman, 2001). Blood is drawn and then centrifuged at 3000 rpm for 15 minutes until the serum and blood cells separate. Blood serum obtained was used to determine total cholesterol, triglycerides, HDL-

cholesterol and LDL-cholesterol using analysis kit (Azima *et al.*, 2004).

Research Parameter

Analysis of cholesterol, HDL and triglycerides performed enzymatically using the cholesterol kit, kit triglycerides, HDL precipitant kit. Absorbance measurements performed using a spectrophotometer at a 500 nm wavelength.

Total cholesterol levels were measured enzymatically using CHOD-PAP method (cholesterol oxidase-p-amino-phenozone). A total of 5 mL and 5 mL of blood serum standard, each 500 mL mixed reagent. Then, serum, standards and blanks were incubated for 20 minutes at a 25-27°C temperature and the absorbance read at a 500 nm wavelength by using a spectrophotometer. Total cholesterol level was calculated using the formula:

$$\text{Total Cholesterol } \left(\frac{\text{mg}}{\text{dL}} \right) = \frac{\Delta A \text{ sampel}}{\Delta A \text{ Standar}} \times \text{Standard concentration } \left(\frac{\text{mg}}{\text{dL}} \right)$$

Triglyceride levels were measured by the GPO-PAP method (glycerol phosphate oxidase-p-aminophenozone). A total of 5 mL of blood serum and 5 mL of each standard 500mL mixed reagent. Then, serum, standards and blanks were incubated for 20 minutes at a temperature of 25-27°C and the absorbance read at a wavelength of 500 nm. Triglyceride levels calculated by the formula:

$$\text{Triglycerida } \left(\frac{\text{mg}}{\text{dL}} \right) = \frac{\Delta A \text{ sampel}}{\Delta A \text{ Standar}} \times \text{Standard concentration } \left(\frac{\text{mg}}{\text{dL}} \right)$$

HDL-cholesterol was measured by precipitation chylomicrons, VLDL and LDL plasma by adding a precipitant reagent composed by fosfotungstat acid and magnesium chloride. A total of 100 mL of blood serum plus 250 mL precipitant and incubated for 10 min at room temperature. After that, the mixture was centrifuged at a speed of 4000 rpm for 10 minutes so that chylomicrons, VLDL and LDL forms a precipitate. Clear supernatant was separated and tested levels of HDL-cholesterol CHOD-PAP method as in the measurement of total cholesterol. LDL-cholesterol levels obtained from the Friedewald equation as follows:

$$\text{LDL-cholesterol } \left(\frac{\text{mg}}{\text{dL}} \right) = (\text{Total Cholesterol} - \text{HDL cholesterol}) - \frac{1}{5} \text{ triglycerides}$$

Data measurements of total cholesterol, HDL-cholesterol, LDL-cholesterol and triglycerides were analyzed by ANOVA, treatment showed no significant difference followed by LSD test at 5% confidence level.

RESULT

Total cholesterol and triglyceride white rats at the beginning of treatment and end of treatment is shown in Table 1. Decrease in total cholesterol levels at most Mundu fruit methanol extract treatment doses of 3.6 mg/kg bw of (32.17 mg/dL). This decrease was greater than the addition of simvastatin (24.1 mg/dL). Decreased levels of triglycerides treatment doses of 2.7 and 3.6 mg / kg bw same relatively large (6.36 and 7.22 mg/dL), and is lower when compared with addition of simvastatin is able to reduce levels of triglycerides by 24.58 mg/dL.

HDL-cholesterol and LDL-cholesterol mice at the beginning of treatment and end of treatment can be seen in Table 2 HDL-cholesterol is known as good cholesterol. Increased levels of HDL-cholesterol was greatest in the methanol extract of the fruit Mundu treatment dose of 3.6 mg/kg bw, which amounted to 5.62 mg/dL, compared to doses of 1.8 and 2.7 mg/kg bw which only amounted to 1, 61 and 1.07 mg/dL. Increased levels of HDL-cholesterol in control is relatively small in the amount of 0.75 mg/dL. Increased levels of HDL-cholesterol following addition of

simvastatin (0.9 mg/dL) lower than the third methanol extract treatment. Neal (2005) suggest that simvastatin is a hypolipidemic drug that can increase levels of HDL-cholesterol by about 5%.

LDL-cholesterol is known as bad cholesterol. Addition of the methanol extract of the fruit Mundu treatment dose of 3.6 mg/kg bw able to reduce levels of LDL-cholesterol by 14.65 mg/dL, greater than the dose of 2.8 mg/kg bw (10.58 mg/dL) and simvastatin (8.33 mg/dL). Increased levels of LDL-cholesterol occurs in the control.

Table 1: Levels of total cholesterol and triglycerides (mg / dL) rats at the beginning of treatment and end of treatment

Treatment (mg / kg bw)	Total cholesterol (mg / dL)		Triglycerides (mg / dL)	
	Beginning treatment	End of treatment	Beginning treatment	End of treatment
Control	91,34±4,46 ^a	93,97±5,21 ^a	65,05±4,22 ^a	73,24±4,27 ^b
3,6 Statin	110,25±4,31 ^b	86,15±11,07 ^a	56,32±5,18 ^a	31,74±1,59 ^a
1,8 extract	99,22±4,67 ^a	112,38±8,99 ^b	58,36±4,28 ^a	68,71±4,76 ^b
2,7 extract	105,77±6,22 ^a	99,76±38,82 ^a	43,19±4,19 ^a	36,83±15,88 ^a
3,6 extract	112,54±5,45 ^b	80,37±29,69 ^a	39,28±4,28 ^a	32,06±16,79 ^a

Note: numbers followed by the same letter within a column indicate no significant difference ($\alpha = 5\%$)

Table 2. Levels of HDL-cholesterol and LDL-cholesterol (mg / dL) white rats at the beginning of treatment and end of treatment

Treatment (mg / kg bw)	HDL (mg/dL)		LDL (mg/dL)	
	Beginning treatment	End of treatment	Beginning treatment	End of treatment
Control	25,46±7,35 ^a	26,21±7,21 ^a	87,38±7,29 ^a	89,58±8,66 ^a
3,6 Statin	37,44±4,67 ^a	38,34±6,28 ^a	81,75±6,47 ^a	73,42±6,39 ^a
1,8 extract	38,21±6,53 ^a	39,82±6,24 ^a	78,56±8,28 ^a	81,24±4,67 ^a
2,7 extract	32,86±7,88 ^a	33,93±7,35 ^a	98,21±7,46 ^a	87,63±7,37 ^a
3,6 extract	36,29±5,78 ^a	41,91±7,43 ^a	82,47±6,43 ^a	67,82±6,43 ^a

Note: numbers followed by the same letter within a column indicate no significant difference ($\alpha = 5\%$)

DISCUSSION

Mundu fruit extraction is done by maceration using methanol solvent. The resulting methanol extract of purple yellow and is estimated to contain compounds that are polar. Based on the phytochemical analysis, the methanol extract containing flavonoids, polyphenols, quinones and alkaloids. Deachathai *et al.* (2005), the content of chemical compounds in the genus *Garcinia* i.e flavonoids, xanones, triterpenoids, and quinones.

The composition of chemical compounds contained in extracts of methanol is still very much. The methanol extract of the Mundu fruit contains many bioactive compounds, each compound has a different biological activity. Under these conditions, the bioactive compounds contained in the methanol extract of the fruit Mundu can interact hypolipidemic produce a complex mechanism.

According Deachathai *et al.*, 2005, the Mundu fruit contains flavonoids, xanones and quinones. This compound is expected to have an activity that can lower total cholesterol and triglycerides (Qudsiyah & Ward, 2007). Total cholesterol and triglycerides in the control treatment tended to be increased. This condition may occur due to the intake of carbohydrates that will induce the synthesis of fatty acids by fatty acid synthase enzyme into triglycerides.

Simvastatin (statin derivative) is hypolipidemic drug that inhibits cholesterol synthesis in the liver. Statins have active functional groups in the form of the lactone ring which is a dihydroxy acid as its active form. In the body of the lactone ring functional groups will be hydrolyzed to produce dihydroxy acid form of active compounds that

can inhibit the enzyme competitively with HMG-CoA reductase. According to Kawano *et al.* (2007), in addition to works by blocking the synthesis of endogenous cholesterol, statins can also increase the activity of the enzyme lipoprotein lipase (LPL). LPL enzyme is able to hydrolyze triglycerides into free fatty acids, and causes a decrease in blood triglycerides. The mechanism of reduction in triglyceride levels may be related to the inhibition of lipolysis in the liver and inhibition of endogenous cholesterol biosynthesis. Total cholesterol cannot give an exact picture of the effect of the methanol extract of the fruit Mundu.

Total cholesterol is cholesterol from food (chylomicrons), VLDL, LDL and HDL. Under these conditions, the mechanism of reduction in total cholesterol levels correlated with more specific lipoproteins, such as HDL-cholesterol and LDL-cholesterol.

Conditions are expected because of high cholesterol levels in the liver resulting in liver reduces the synthesis of LDL receptors remove excess cholesterol hepatocytes and hepatocytes in the form of VLDL into the blood stream. Catabolism of VLDL in the blood lead levels into LDL. This process causes the levels of LDL-cholesterol to rise (Felig and Frohman, 2001).

Simvastatin is a hypolipidemic drug that works by blocking an enzyme standard of HMG-CoA reductase. Inhibition of this enzyme causes a decrease in VLDL synthesis and increasing LDL receptors in the liver. Decrease in VLDL synthesis which brings triglycerides indirectly cause a decrease in blood triglyceride levels, whereas the increase in LDL receptors in the liver causes a decrease in blood levels of LDL-cholesterol (Neal, 2005). Another

consequence of the addition of simvastatin is synthesized VLDL cholesterol ester poor heart that this VLDL cholesterol esters from HDL to get. This process can lead to decreased levels of HDL-cholesterol following addition of simvastatin.

Methanol extract containing flavonoids, xanton, triterpenoids, and quinone groups are estimated to contain lactone ring. In the body, the active lactone groups will be hydrolyzed to form dihydroxy acid, the active compounds that can inhibit the enzyme competitively with HMG-CoA reductase. Barriers occur because dihydroxy acid form similar to the form of HMG-CoA reductase.

The mechanism of the biological activity of compounds of the research work by inhibiting the enzyme HMG-compete with CoA reductase. HMG-CoA reductase is an enzyme that catalyzes the conversion of 3-hydroxy-3-methyl-glutaril CoenzimA to mevalonic acid, one of the important stages in the pathway of cholesterol synthesis. Barriers of this enzyme causes an increase in the density of LDL receptors in the liver cells resulting in decreased levels of cholesterol, LDL cholesterol and increase HDL number.

From this study it can be concluded that the methanol extract of Mundu fruit (*Garcinia dulcis*) can reduce total cholesterol, triglycerides and LDL-cholesterol and the methanol extract of Mundu fruit (*Garcinia dulcis*) can increase levels of HDL-cholesterol.

ACKNOWLEDGMENTS

Acknowledgements submitted to DP2M the Directorate General of Higher Education for funding this research.

REFERENCES

- Adaramoye, O.A., Akintayo, O., Achem, J., Fafunso, M.A., 2008. Lipid-lowering effects of methanolic extract of *Vernonia amygdalina* leaves in rats fed on high cholesterol diet. *Vascular Health and Risk Management* 4: 235-241
- Adaramoye, O.A., and Arisekola, M., 2013. Kolaviron, a biflavonoid complex from *Garcinia kola* seeds, ameliorates ethanol-induced reproductive toxicity in male wistar rats. *Nigerian Journal of Physiological Science* 28: 009-015.
- Anonim, 1993. Penapisan Farmakologi, Pengujian Fitokimia dan Pengujian Klinik, Yayasan Pengembangan Obat Bahan Alam, *Phytomedica*. Jakarta.
- Asano, J., Chiba, K., and Tada, M., 1996. Cytotoxic xanthenes from *Garcinia hanburyi*. *Journal of Phytochemistry* 41: 815-820.
- Azima, F., Muchtadi, D., Zakaria, F.R., and Priosoeryanto, B.P., 2004. Potensi Anti-Hiperkolesterolemia Ekstrak Cassia Vera (*Cinnamomum burmanni* Nees ex Blume), *Jurnal Teknologi dan Industri Pangan* 15:145-152
- Deachathai, S.W., Mahabusarakam, Paichit, S.P., Taylor, W.C., Zhangarid, Y.J., and Yang, C.R., 2005. Phenolic Compound from The Flowers of *Garcinia dulcis*. *Journal of Phytochemistry* 66: 2368-2375.
- Delgado, J.N., and Remers, W.A., 1991. *Wilson and Gisvold's. Textbook of Organic Medicinal and Pharmaceutical Chemistry*. 9th Edition. JB Lippincott Co. Philadelphia. 8-9.
- Dubey, A.K., Devi, A., Kutty, G., and Shangkar, R.P., 2005. Hypolipidemic Activity of *Ginkgo biloba* Extract EGB 761 in Hypercholesterolemic Wistar Rat, *Iranian Journal of Pharmacology and Therapeutics* 4:9-12.
- Felig, P., and Frohman, L.A., 2001. *Endocrinology and Metabolism*, 4th edition, McGraw-Hill Inc.
- Fukuyama, Y., Minami, H., and Kuwayama, A., 1998. Garsubellins, poly-isoprenylated phloroglucinol derivatives from *Garcinia subelliptica*. *Journal of Phytochemistry* 49: 853-857.
- Gad, S.C., and Chengelis, C.P., 1992. *Animal Models In Toxicology*, Marcel Dekker Inc, New York.
- Iinuma, M., Tosa, H., Ito, T., Tanaka, T., and Riswan, S., 1996. Three new benzophenone-xanthenes dimers from the root of *Garcinia dulcis*. *Chemical and Pharmaceutical Bulletin* 44, 1744-1747.
- Iwu, M.M., 1986. *Biflavanones of Garcinia: Pharmacological and biological activities*. In Cody, V., Middleton, E., and Harbone, J.B., (Eds). *Plant Flavonoids in Biology and Medicine*, Alan R. Liss, New York, 485-488.
- Iwu, M.M., Igbo, O.A., Okunji, C.O., and Tempesta, M.S., 1990. Anti-diabetic and aldose reductase activities of biflavanones of *Garcinia kola*. *Journal of Pharmacology* 42: 290-292.
- Kawano, H., Nagata, T., Narahasa, M., Kanazawa, M. and Miyake, M., 2007. Triglycerides Accumulation by Peroxisome Proliferator In Rat, *Biological and Pharmaceutical Bulletin* 30:627-632.
- Kosela, S., Hu, L.H., Rachmatia, T., Hanafi, M., and Sim, K.Y., 2000. Dulxanthone F-H, Three New Pyranoxanthones from the *Garcinia dulcis*, *Journal of Natural Product* 63: 406-407.
- Likhitwitayawuid, K., Channamasathien, W., Ruangrunsi, N., and Krungkrai, J., 1998. Xanthone with Antimalarial Activity from *Garcinia dulcis*, *Planta Medica* 64: 281-282.
- Lin, Y.M., Anderson, H., Flavin, M.T., Pai, Y.H., Mata-Greenwood, E., Pengsupar, P.T., Pezzuto, J.M., Schinazi, R.F., Hughes, S.H., and Chen, F.C., 1997. In vitro anti-HIV activity of biflavanoids isolated from *Rhus succedanea* and *Garcinia multiflora*. *Journal of Natural Product* 60: 884-888.
- Lin, K.W., Huang, A.M., and Yang, S.C., 2012. Cytotoxic and antioxidant constituents from *Garcinia subelliptica*. *Journal of Food Chemistry* 135: 851-859.
- Minami, H., Takahashi, E., Fukuyama, Y., Kodama, M., Yoshizawa, T., Nakagawa. 1995. Novel xanthenes with superoxide scavenging activity from *Garcinia subelliptica*. *Chemical and Pharmaceutical Bulletin* 43: 347-349.
- Neal, M.J., 2005. *At a Glance Farmakologi Medis*, edisi ke-5. Penerbit Erlangga. Jakarta
- Olaiya, C.O., and Omolekan, T.O., 2013. Antihypercholesterolemic activity of ethanolic extract of in rats. *African Health Sciences* 13: 1084-1090
- Qudsiyah, H dan Wardoyo, E.R.P., 2007. *Pemanfaatan Batang Mundu (Garcinia dulcis (Roxb.) Kurz) Sebagai Antihiperkolesterolemia*, Laporan Penelitian, Universitas Tanjungpura Pontianak.
- Sukpondma, Y., Vatcharin, R., and Phongpaich, S., 2005. Xanthone and Sesquiterpene Derivatives from the Fruits of *Garcinia scor-techinii*. *Journal of Natural Product* 68: 1010-1017.
- Sweeney, P.W., 2008. Phylogeny and floral diversity in the genus *Garcinia* (Clusiaceae) and relatives. *International Journal of Plant Science* 169: 1288-1303.
- Wulandari, D., Suarsih, H., Ifadatin, S., Widiyantor, A., Kusmawati, A., dan Wardoyo, E.R.P., 2011. Efek Ekstrak Metanol Buah *Makassar (Brucea javanica (L.) Merr)* terhadap Profil Lipoprotein dan Triglycerida Tikus Putih (*Rattus norvegicus* L) Hiperlipidemia. *Berkala Penelitian Hayati* 6D: 25-29.
- Yokozawa, T., Cuo, E.J., Sasaki, S., Satou, A., Okamoto, T., and Sei, Y., 2006. The Protective Role of Chinese Prescription Kange-Karyu Extract on Diet-Induced Hypercholesterolemia in Rats, *Biological and Pharmaceutical Bulletin* 29:760-765.
- Zhang, L.J., Chiou, C.T., and Cheng, J.J., 2010. Cytotoxic polyisoprenyl benzophenonoids from *Garcinia subelliptica*. *Journal of Natural Product* 73: 557-562

ICI World of Journals (/search/form) / **Berkala Penelitian Hayati**

[◀ Back](#)

Berkala Penelitian Hayati



English title:

n/d

ISSN:

0852-6834, 2337-389X

GICID:

n/d

DOI:

n/d

Website:

<http://berkalahayati.org> (<http://berkalahayati.org>)

Publisher:

n/d

Country:

ID

Language of publication:

n/d

Deposited publications: 0 > Full text: 0% | Abstract: 0% | Keywords: 0% | References: 0%

[Issues and contents](#)[Journal description \(\)](#)[Details \(\)](#)[Scientific profile \(\)](#)[Editorial office \(\)](#)[Publisher \(\)](#)[Metrics \(\)](#)

JOURNAL OF BIOLOGICAL RESEARCHES is international peer-reviewed, open access journal journal of life science, formerly is Berkala Penelitian Hayati, a scientific journal using Indonesian language. The first edition of this journal is Vol 1 No 1 at Mei 1995. It was accredited by Education and Cultural Ministry number 134/DIKTI/Kep/2001. Since April 2012 the journal was change into English and the name of the journal change to Journal of Biological Researches. This journal are indexed by Google Scholar, Academia.edu, and portal garuda. This journal publishes original research, applied, review article, and educational articles in all areas of biology. Authors are encouraged to submit complete unpublished and original works, which are not under review in any other journals. The scopes of the journal include, but not limited to, the following topic areas: botany, zoology, ecology, microbiology, physiology, nano biology, coastal biology, hydrobiology, neurobiology, genetics, developmental biology, biochemistry and molecular biology, biophysics and life science. The journal is published in both printed and online versions. The online version is free access and downloads.

Non-indexed in the ICI Journals Master List 2019

Not reported for evaluation

Archival ratings ➤

Citations

Reports 2010-2019

3 - Number of journal citations

Included auto-citations - 0

Main page (<http://jml.indexcopernicus.com>)

MSHE points 2019: n/a

© Index Copernicus 2017





Year

2019

2018

2017

2016

2015

2014

2013

2012

2011

2010

2009

2008

2007

2006

2005

2004

2003

2001

2000

1999

1997

1996

1995

VOL 19, NO 2 (2014) : VOL 20, NO 1 (2014) : VOLUME VOL 20, NO 2 (2014) : VOLUME

POLYMORPHISM ANALYSIS OF IL17RA GENE TO THE IL17RA CONCENTRATION AND CHRONICITY DIFFERENCES IN NEFRITIS LUPUS NL PATIENTS

Mahrus Ismail, Hani Susanti, Widodo Widodo

Dec 09, 2014

PCR APPROACH FOR RAPID DETECTION OF *Escherichia coli* IN TEMPE USING A SPECIFIC PRIMER

Siti Harnina Bintari, Fidia Fibriana, Dewi Mustikaningtyas, Retno Sri Iswari

Dec 09, 2014

THE INFLUENCE OF Aloe vera AND XENOGRAFT XCB TOWARD OF BONE MORPHO PROTEIN 2 BMP2 EXPRESSION AND AMOUNT OF OSTEOBLAST OF ALVEOLAR BONE INDUCED INTO TOOTH EXTRACTION SOCKETS *Cavia cobaya*

Utari Kresnoadi, Retno Pudji Rahayu

Dec 11, 2014

SLAUGHTER HOUSE SOLID WASTE MANAGEMENT IN INDONESIA

Rheny Ratnawati, Yulinah Trihadiningrum

Dec 23, 2014

A STUDY OF TYPE AND INTENSITY OF DISEASE INFECTING BANANA PLANTS *Musa sp* AT TEGALAGUNG VILLAGE SEMANDING SUBDISTRICT

Supiana Dian Nurtjahyani, Devi Shyntya Agustin

Dec 31, 2014

THE EFFECT OF MUNDU FRUIT *Garcinia dulcis* ROXB KURZ METHANOL EXTRACT ON LYPHOPROTEIN PROFILE AND TRYGLISERIDE WHITE RAT *Rattus norvegicus* L

Arum Setiawan, Laila Hanum, Elvi Rusmiyanto Pancaning Wardoyo

Feb 15, 2015

BIOPHYSICS ENVIRONMENTAL CONDITIONS OF SWAMP BUFFALO *Bubalus bubalis* PAMPANGAN IN DISTRICT RAMBUTAN SOUTH SUMATERA

Yuanita Windusari, Erwin Nofyan, Mustafa Kamal, Laila Hanum, Rahmat Pratama

Jun 24, 2014

EFFECT OF GROWTH CONDITIONS ON BIOFILM FORMATION BY PHENOLDEGRADING BACTERIA ISOLATED FROM POLLUTED AND NONPOLLUTED SOURCES

Arifah Khusnuryani, Erni Martani, Tri Wibawa, Jaka Widada

Mar 27, 2015

Dendrobium spectabile (Blume) Miq. IN VITRO CULTURE AND ITS ACCLIMATIZATION ON MUS MEDIA WITH ANTIMICROBIAL AND ALCOHOLIC SUGAR SUPPLEMENTATION

Untung Santoso, Y. Sri Wulan Manohara, Kusningrum Rochiman S.

Aug 25, 2015

Berkalah Penelitian Hayati
Journal of Biological Researches

Published by The East Java Biological Society Jalan Surakarta No 5, Malang, East Java 65114, Indonesia

☎ +62 341 570631

✉ berkalahayati@yahoo.com / jbiolresearches@gmail.comThis work is licensed under a [Creative Commons Attribution-NoDerivatives 4.0 International License](https://creativecommons.org/licenses/by-nd/4.0/).**Biology Departments**

Universitas Muhammadiyah Malang

Universitas Brawijaya

UIN Maliki Malang

Universitas Islam Malang

Universitas Airlangga

Institut Teknologi Sepuluh Nopember

Universitas Gadjah Mada

IKIP PGRI Madiun

Universitas Jember

Universitas Negeri Malang

Institut Teknologi Bandung

Universitas Jenderal Soedirman

Universitas Diponegoro
Universitas Sebelas Maret
Universitas Indonesia
Insitut Pertanian Bogor
Universitas Mulawarman
Universitas Lambung Mangkurat
Konsorsium Biologi Indonesia
UIN Alauddin Makassar
Universitas Esa Unggul Jakarta
Universitas Ahmad Dahlan Yogyakarta

Current Projects

International Biodiversity Conservation (IBC Raja Ampat)
Smartbio

© 2017 Berkalahayati Developer Team

Journal of BIOLOGICAL RESEARCHES

Former of Jurnal Berkala Penelitian Hayati

JUNE 2014

Vol. 19

No. 2

CONTENT

The Influence of *Aloe vera* and Xenograft (Xcb) Toward of Bone Morpho Protein 2 (Bmp2) Expression and Amount of Osteoblast of Alveolar Bone Induced Into Tooth Extraction Sockets (Cavia Cobaya)

Utari Kresnadi and Puji Rahayu..... 51

PCR Approach for Rapid Detection of Escherichia Coli in Tempe Using a Specific Primer

Siti Harnina Bintari, Fidia Fibriana, Dewi Mustikaningtyas, Retno Sri Iswari..... 56

Polymorphism Analysis of Il-17ra Gene to The Il-17ra Concentration and Chronicity Differences in Nefritis Lupus (NI) Patients

Mahrus Ismail, Hani Susanti and Widodo..... 61

A Study of Type and Intensity of Disease Infecting Banana Plants (*Musa sp.*) at Tegalagung Village, Semanding Subdistrict

Supiana Dian, Nurtjahyani, and Devi Shyntya Agustin..... 66

Slaughter House Solid Waste Management In Indonesia

Rhenny Ratnawati and Yulinah Trihadiningrum 69

The Effect of Mundu Fruit (*Garcinia dulcis* (Roxb.) Kurz) Methanol Extract on Lyphoprotein Profile and Trygliseride White Rat (*Rattus norvegicus* L)

Arum Setiawan, Laila Hanum, and Elvi Rusmiyanto Pancaning Wardoyo74

Biophysics Environmental Conditions of Swamp Buffalo *Bubalus bubalis* Pampangan in District Rambutan South Sumatera

Yuanita Windusari, Erwin Nofyan, Mustafa Kamal, Laila Hanum and Rahmat Pratama.....78

Effect of Growth Conditions on Biofilm Formation by Phenol-Degrading Bacteria Isolated From Polluted and Non-Polluted Sources

Arifah Khusnuryani1, Erni Martani, Tri Wibawa, Jaka Widada..... 82

**PBI
East Java**

Journal of Biological Researches is a peer-reviewed, open access journal, published by Indonesia Biological Association, East Java, Indonesia. It publishes articles and short communications in biological field. Author whose work is accepted for publication is subjected to pay 1000.000 IDR (100 USD) per printed article up to 5 pages. Additional page will be charged 50.000 IDR (5 USD) per printed page without photo(s). Color photo(s) page will be charged 100.000 (10 USD) per printed page.

Information about Indonesia Biological Association membership, please contact PBI board.

Indonesia Biological Association, East Java Board:

Chairman: Prof. Sutiman B. Sumitro, M.Sc., D.Sc

Secretary : Dra. Herawati Susila, M.Sc., Ph.D

Treasurer : Dr. dr. Tjandrakirana M. Sjaifullah Noer, M.Sp. And

EDITORIAL BOARD

Editor in chief : Prof. Sutiman B. Sumitro, M.Sc., D.Sc

Member : Prof. Dr. Bambang Irawan

Prof. Dr. Hitoshi Sawada

Prof. Roger Price

Prof. Fatchiyah, Ph.D

Prof. Wolfgang Nellen

Dr. Paul Kessler

Dr. Bagyo Yanuwadi

Dr. Akira Kikuchi

Editorial Officer:

Romaidi, M.Si

Roudlotul Jannah, S.Si

Office:

Jl, Surakarta No. 5 Malang

East Java, Indonesia 65114

Telp / Fax : (0341) 570631

Website : www.berkalahayati.org

Email: berkalahayati@yahoo.com

Indexed by:



academia.edu

MANUSCRIPT PREPARATION

REGISTER

<http://www.berkalahayati.org/index.php/bph/user/register>

Full Article. A full article is expected to present an original and important major advance in bioscience having wide research scheme and deep discussion of the findings. The article should be typed in paper length of 15 pages or 3000 – 4500 words (including figures and tables).

Title. Title consist of not more than 14 words.

Author's Data. Author full name (no abbreviation) and author full address (complete with postcode).

Abstract. Abstract is not more than 400 words which has no reference and does not contain numbers, abbreviation, acronym or measurement unless essential. Key Words. Maximum 6 words, first key word being the most important.

INTRODUCTION. Write down the background of the research and mention the previous studies that had been done. State one question(s) that are needed to answer through your work. Give a description (local and scientific name) of the studied organism.

MATERIALS AND METHODS. The method section need to be detailed enough to allow reviewers to answer some or all of the following questions: (i) Is the study an experimental or an exploration?; (ii) Are the methods described in sufficient detail so that the study can be replicated?; (iii) If your research uses the previous researcher method, please describe briefly that method. If you make any modification, describe the modified part; (iv) Name the number of the samples, give courtesy to whom you obtain the samples; (v) State seasonal variation of the habitat (if applicable) or date of sampling; (vi) Human materials should be collected in conformation to standard ethics and got with informed consent.

RESULTS. Results and Discussion should be written separately. State the obtained results based on the used methods. Do not present same data in both table and graph format. Do not state references in Result section. Means should be given standard deviation. All data given in Result should be stated in the tables or graph. See Figures and Tables criteria below.

DISCUSSION. Discuss your data by comparing the current reported data with the previous results. Highlight the similarities, as well as the differences, and the uniqueness of your findings. Explain why your result(s) is coming to be. End the discussion by giving a conclusion and future research in that particular topic.

ACKNOWLEDGEMENT. State the grant source (Institution, year of the contract) and the person to whom the grant was given. Name the person(s) that help your work.

REFERENCES. References only list the published or accepted material for publication. It should be at least 80% primary references (journal articles). Accepted article(s)

but not yet published should be sent as manuscript appendix. Website, personal communications, unpublished data should be incorporated in the text in parentheses. Example: ... The deduced amino acid sequence of the gene was compared for homology with BLAST search (<http://www.ncbi.nlm.nih.gov/blast>). Personal communication need to be written with the initials surname of the source accompanied with the year of the statement, e.g., (Takahashi 2007, personal communication). Authors should provide proof(s) of the communication.

Journal article

Author AB, Author CD, Author EF. 2001. Title of article. Journal 60:128-132. Example:
Bagnara JT., and Fernandes PJ., 1993. Hormonal Influences on the Development of Amphibian Pigmentation patterns. Zoological Science 10: 733-748.

Book

Author GH, Author IJ. 2006. Title of Book. City of publication: Publisher's name. Example:
Brown TA., 1993. Genetics a Molecular Approach, 2nd ed. Chapman & Hall, London, 270, 302-303.

Chapter in a book

Author KL, Author MN. 1999. Title of a chapter: a subtitle. In: Editor MN, Editor OP (eds). Title of Book. 2nd ed. City of publication: Publisher's name. p 200-235. Example: Templeton AR., 1989. The Meaning of Species and Speciation; A Genetics Perspective dalam Otte D., dan Endler, JA. (Ed.) Speciation and its Conseuences. Sinauer Associaties, Inc., Sunderland. 1-27.

Thesis or Dissertation

Author MN. 2002. Title [Dissertation]. Location of university: Name of university. Example:
Widayanti KA. 2006. Color perception of L4M5 gene carrier female Macaca fascicularis [Thesis]. Bogor: Bogor Agricultural Univ.

FIGURE. Use graph in maximum width of 8.5 cm. Numbers and title are written in 8 point Times New Roman font.

STATISTICAL GRAPHS. Give standard deviation to every mean value. Authors that used Microsoft Excell Program need to give the raw data.

PHOTOS. Add measurement scale if needed. Arrows should be given to point certain objects.

TABLE. Give standard deviation to every mean value. Table titles should be concise. Explanatory material, notes on measurements, and other general information that applies to the whole table should be included as the first, unnumbered footnote and not in the table title. Numbers and table title are written in 8 point Times New Roman font.

The effect of mundu fruit *Garcinia dulcis* Roxb Kurz methanol extract on lyphoprotein profile and trygliseride white rat *Rattus norvegicus* L

By Arum Setiawan

THE EFFECT OF MUNDU FRUIT (*Garcinia dulcis* (ROXB.) KURZ) METHANOL EXTRACT ON LYPHOPROTEIN PROFILE AND TRYGLISERIDE WHITE RAT (*Rattus norvegicus* L)

Arum Setiawan*, Laila Hanum* and Elvi Rusmiyanto Pancaning Wardoyo**

*Departement of Biology, Faculty of Mathematic and Natural Science, Sriwijaya University,
Jl. Palembang-Prabumulih KM 32, Indralaya (Ol) South Sumatera, Indonesia, 30662

**Departement of Biology, Faculty of Mathematic and Natural Science, Tanjungpura University
Jl. Prof. Hadari Nawawi, Pontianak, West Kalimantan, Indonesia 78124

Corresponding Author email: lailahanum@gmail.com

ABSTRACT

Garcinia dulcis (Roxb.) Kurz (Guttiferae) is a tropical fruit tree native to Indonesia and its known as an Asian medicinal plant used in folk medicines. In ethno pharmacology, *Garcinia dulcis* has been used in traditional medicine, such as anti-inflammatory, anti-tumor, anti-malaria and anti-bacterial. The scientific studies of biological activity of *Garcinia dulcis* on lipide profile and triglyceride are relatively measily. Therefore, the objectives of this research were to know the effect of methanol extract of *Garcinia dulcis* fruit on total cholesterol, low density lipoprotein-cholesterol, high density lipoprotein-cholesterol and triglyceride rat hyperlipidemic. These studies using *Garcinia dulcis* fruit from Banyuwasin, South Sumatra. In this observation, 30 male rat from Laboratory and Research Institute for Integrated Testing, Gadjah Mada University, 2 months old were used and divided into five groups randomly. The first group was given high cholesterol and lipid diet, as a normal group. The second group was given high cholesterol, lipid diet and simvastatin 3,6 mg/kg bow/day dosage, as a simvastatin group. The third, fourth and fifth groups was given high cholesterol, lipid diet and methanol extract 1.8; 2.7 and 3.6 mg/kg bow/day dosage. Total cholesterol, low density lipoprotein-cholesterol, high density lipoprotein-cholesterol and triglyceride contains using spectrophotometry method, were observed 1 and 30 days. The differences among groups were analyzed using Anova followed by LSD to locate the differences. LSD analysis showed that *Garcinia dulcis* methanol extract treatment doses of 3.6 mg/kg bw and simvastatin lowers total cholesterol levels highest in the amount of 32.17 mg/dL and 24.1 mg/dL, significantly different compared with the treatment dose of 1.8 mg/kg bw. *Garcinia dulcis* fruit methanol extract treatment doses of 3.6 mg/kg bw and simvastatin lowers triglyceride levels by 7.22 mg/dL and 24.58 mg/dL, significantly different compared with the treatment dose of 1.8 mg/kg bw and controls. *Garcinia dulcis* fruit extract dose of 3.6 mg/dL lower low density lipoprotein-cholesterol highest of 16.45 mg/dL. *Garcinia dulcis* fruit methanol extract dose of 3.6 mg/kg bw was also able to increase the levels of high density lipoprotein-cholesterol by 5.62 mg/dL. The results showed that the fruit of *Garcinia dulcis* contain compounds that have potential as agents antihyperlipidemia and presumably through inhibition of the enzyme HMG-CoA reductase.

Key words: cholesterol, *Garcinia dulcis*, triglyceride

INTRODUCTION

Mangosteen is a plant that grow and spread in the lowland tropical forests of Southeast Asia, starting from the region of Thailand, Peninsular Malaysia and Indonesia and is known as the queen of fruits. In Indonesia, genus *Garcinia* are common in Sumatra, Java, Kalimantan and Maluku. The genus *Garcinia* (Familia: Guttiferae) is a tropical plant and has approximately 350 species, in-cluding *Garcinia dulcis*, *Garcinia bancana*, *Garcinia bor-neensis*, *Garcinia parvifolia*, *Garcinia rostrata*, and *Garcinia* spp (Sweeney, 2008). Besides eaten, this plant has been widely used by people for a variety of traditional tre-atments, such as limphatis, parotid, anti-inflammatory, anti-tumor and anti-bacterial activity on *Staphylococcus aureus*, *Shigella dysenteriae*, *Shigella flexneri*, *Escherichia coli*, *Vibrio cholerae* (Deachathai *et al.*, 2005).

The content of chemical compounds that have been found to vary widely in the genus *Garcinia* are flavonoids, xanton, triterpenoids, and quinone (Deachathai *et al.*, 2005). Flavonoid compounds that have been found in the genus *Garcinia* mostly biflavonoid types. *Garcinia scorte-chinii* fruit contains compounds biflavonoid (Sukpondma *et al.*, 2005) and kolaflavanon (Adaramoye & Muritala, 2013).

Research has been done on the genus *Garcinia* shows there biflavonoid other compounds, such as amantoflavon, agathisflavon, robustaflavon, hinokiflavon, volkensiflavon, rhusflavanon, succedaneflavon of *Garcinia multiflora* which shows antidiabetic and antiviral activity and hepatoprotective (Lin *et al.*, 1997; Iwu *et al.*, 1990). Five compounds xanton i.e dihidroksixanton, dihidroxydes-D-garcigerrin, metilsimpoxanton, simpoxanton and

garcinixanton apparently showed antimalarial activity (Likhawitayawuid *et al.*, 1998). Benzophenone derivative compounds, namely garcinol, isogarcinol, xantosimol, isoxantosimol and sikloxantosimol showed antibacterial activity against *Staphylococcus aureus* (Linuma *et al.*, 1996).

Several compounds of *Garcinia xanthon* showed antioxidant activity, namely dihydroxy furan and 1,4,5-trihidroksixanton (Minami *et al.*, 1995). The compound 3-hydroxy- β -20,29,30-trinorlupan-9-on garciniellipton and Garsubellin A isolated from *Garcinia subelliptica* showed cytotoxic activity, antibiotics, and increase the activity of choline asetiltransferase (Lin *et al.*, 2012; Zhang *et al.* 2010; Asano *et al.*, 1996; Fukuyama *et al.*, 1997; Kosela *et al.*, 2000).

Until now, research on the activity of the anti-hyperlipidemia methanol extract of the fruit *Garcinia dulcis* from South Sumatra on lipoprotein profile and triglycerides are still rarely found. Antihyperlipidemia activity can be known through the inhibition of HMG-CoA reductase compounds. HMG-CoA reductase is an enzyme that catalyzes the conversion of 3-hydroxy-3-methylglutaril coenzyme A (HMG-CoA) to mevalonic acid, one of the important stages in the pathway of cholesterol synthesis. Inhibition of the enzyme leads to increased density of LDL receptors in the liver cells resulting in decreased levels of cholesterol, LDL-cholesterol and triglycerides (Delgado & Remers, 1991).

Based on research conducted by Qudsiyah & Ward (2007), the methanol extract of *Garcinia dulcis* stem can reduce total cholesterol by 65% rats at levels of 3.6 mg/kg/ day. According Olaiya & Omolekan (2013), ethanol ex-tract contained in *Buchholzia coriacea* at dose

of 250 mg/ kg bw has anti-hyperlipidemia activity via decreased levels of total cholesterol, LDL-cholesterol and triglycerides hyperlipidemic mice at 45.9%, 73.5%, and 48.4 %. *Vernonia amygdalina* methanol extract dose of 200 mg/kg bw able to reduce levels of cholesterol, LDL-cholesterol and triglycerides hyperlipidemic mice by 29%, 49% and 28% (Ada-ramoye *et al.*, 2008). *Brucea javanica* seed methanol extract dose of 2.7 mg/kg bw could reduce levels of LDL-cholesterol and triglycerides were hyperlipidemic white mice by 43.8% and 32.2% (Wulandari *et al.*, 2011).

Therefore, this study aims to determine the activity of fruit extracts Mundu (*G. dulcis*) at different doses on levels of total cholesterol, triglycerides, HDL-cholesterol and LDL-cholesterol white rat (*Rattus norvegicus* L.).

MATERIAL AND METHOD

Extraction of Mundu (*Garcinia dulcis*)

Mundu (*G. dulcis*) fruit derived from Banyuasin Regency, South Sumatera Province, and has been determined by Herbarium Bogoriense LIPI, Bogor. Extraction is done by maceration using methanol pa. 250 gram samples that have been dried fruits Mundu, incorporated into maserator then add 500 mL methanol pa. Maceration is done for 3 x 24 hours. The extract obtained was concentrated by using a rotary evaporator for 3 hours.

Test Anti-hyperlipidemia

Test animals used were male Wistar rats (*Rattus norvegicus*), 2 months old with body weight of 150-250 g were obtained from preclinical Laboratory and Research Institute for Integrated Testing (LPPT) UGM. Thirty male white rats of Wistar male (*R. norvegicus*) are healthy and have normal activities were randomly divided into 5 groups with 6 replicates. The control group is the group fed a standard diet and feed high cholesterol and fat for 4 weeks; Statin group is a group fed a standard diet and feed high in fat and cholesterol and simvastatin at a dose of 3.6 mg/kg bw; Group M1, M2 and M3 are the group fed a standard diet and feed high in fat and cholesterol and methanol extract at a dose of 1.8, respectively; 2.7 and 3.6 mg/kg bw/day.

Before treatment, white male rats adapted for 1 week in the animal cages at the temperature and humidity of 50-60% 26-27°C. During the adaptation period, male rats were fed a standard 15 gr/150 gr BB mature rats and drinking water ad libitum (Gad and Chengelis, 1992). According Anonim (1993), the composition of foods to increase levels of lipid and cholesterol of the rats per day are: egg yolks 5%, 10% beef tallow, palm oil and food standard 1% to 100%.

Determination of the dose refers to Dubey *et al.* (2005), the standard drug dose conversion for human consumption simvastatin dose to rats, at 0.018. Methanol extract dose used was 1.8; 2.7; 3.6 mg/kg rat. On days 1 and 30 rat blood drawn through orbital sinus. Before blood sampling rats were fasted for 16 hours with the aim of completing the process of lipid metabolism that comes from food (Felig & Frohman, 2001). Blood is drawn and then centrifuged at 3000 rpm for 15 minutes until the serum and blood cells separate. Blood serum obtained was used to determine total cholesterol, triglycerides, HDL-

cholesterol and LDL-cholesterol using analysis kit (Azima *et al.*, 2004).

Research Parameter

Analysis of cholesterol, HDL and triglycerides performed enzymatically using the cholesterol kit, kit triglycerides, HDL precipitant kit. Absorbance measurements performed using a spectrophotometer at a 500 nm wavelength.

Total cholesterol levels were measured enzymatically using CHOD-PAP method (cholesterol oxidase-p-amino-phenozone). A total of 5 mL and 5 mL of blood serum standard, each 500 mL mixed reagent. Then, serum, standards and blanks were incubated for 20 minutes at a 25-27°C temperature and the absorbance read at a 500 nm wavelength by using a spectrophotometer. Total cholesterol level was calculated using the formula:

$$\text{Total Cholesterol } \left(\frac{\text{mg}}{\text{dL}} \right) = \frac{\Delta A \text{ sampel}}{\Delta A \text{ Standar}} \times \text{Standard concentration } \left(\frac{\text{mg}}{\text{dL}} \right)$$

Triglyceride levels were measured by the GPO-PAP method (glycerol phosphate oxidase-p-aminophenozone). A total of 5 mL of blood serum and 5 mL of each standard 500mL mixed reagent. Then, serum, standards and blanks were incubated for 20 minutes at a temperature of 25-27°C and the absorbance read at a wavelength of 500 nm. Triglyceride levels calculated by the formula:

$$\text{Triglycerida } \left(\frac{\text{mg}}{\text{dL}} \right) = \frac{\Delta A \text{ sampel}}{\Delta A \text{ Standar}} \times \text{Standard concentration } \left(\frac{\text{mg}}{\text{dL}} \right)$$

HDL-cholesterol was measured by precipitation chylomicrons, VLDL and LDL plasma by adding a precipitant reagent composed by fosfotungstat acid and magnesium chloride. A total of 100 mL of blood serum plus 250 mL precipitant and incubated for 10 min at room temperature. After that, the mixture was centrifuged at a speed of 4000 rpm for 10 minutes so that chylomicrons, VLDL and LDL forms a precipitate. Clear supernatant was separated and tested levels of HDL-cholesterol CHOD-PAP method as in the measurement of total cholesterol. LDL-cholesterol levels obtained from the Friedewald equation as follows:

$$\text{LDL-cholesterol } \left(\frac{\text{mg}}{\text{dL}} \right) = (\text{Total Cholesterol} - \text{HDL cholesterol}) - \frac{1}{5} \text{ triglycerides}$$

Data measurements of total cholesterol, HDL-cholesterol, LDL-cholesterol and triglycerides were analyzed by ANOVA, treatment showed no significant difference followed by LSD test at 5% confidence level.

RESULT

Total cholesterol and triglyceride white rats at the beginning of treatment and end of treatment is shown in Table 1. Decrease in total cholesterol levels at most Mundu fruit methanol extract treatment doses of 3.6 mg/kg bw of (32.17 mg/dL). This decrease was greater than the addition of simvastatin (24.1 mg/dL). Decreased levels of tri-glycerides treatment doses of 2.7 and 3.6 mg / kg bw same relatively large (6.36 and 7.22 mg/dL), and is lower when compared with addition of simvastatin is able to reduce levels of triglycerides by 24.58 mg/dL.

HDL-cholesterol and LDL-cholesterol mice at the beginning of treatment and end of treatment can be seen in Table 2 HDL-cholesterol is known as good cholesterol. Increased levels of HDL-cholesterol was greatest in the methanol extract of the fruit Mundu treatment dose of 3.6 mg/kg bw, which amounted to 5.62 mg/dL, compared to doses of 1.8 and 2.7 mg/kg bw which only amounted to 1.61 and 1.07 mg/dL. Increased levels of HDL-cholesterol in control is relatively small in the amount of 0.75 mg/dL. Increased levels of HDL-cholesterol following addition of

simvastatin (0.9 mg/dL) lower than the third methanol extract treatment. Neal (2005) suggest that simvastatin is a hypolipidemic drug that can increase levels of HDL-cholesterol by about 5%.

LDL-cholesterol is known as bad cholesterol. Addition of the methanol extract of the fruit Mundu treatment dose of 3.6 mg/kg bw able to reduce levels of LDL-cholesterol by 14.65 mg/dL, greater than the dose of 2.8 mg/kg bw (10.58 mg/dL) and simvastatin (8.33 mg/dL). Increased levels of LDL-cholesterol occurs in the control.

Table 1: Levels of total cholesterol and triglycerides (mg / dL) rats at the beginning of treatment and end of treatment

Treatment (mg / kg bw)	Total cholesterol (mg / dL)		Triglycerides (mg / dL)	
	Beginning treatment	End of treatment	Beginning treatment	End of treatment
Control	91.34±4.46 ^a	93.97±5.21 ^a	65.05±4.22 ^a	73.24±4.27 ^b
3.6 Statin	110.25±4.31 ^b	86.15±11.07 ^a	56.32±5.18 ^a	31.74±1.59 ^a
1.8 extract	99.22±4.67 ^a	112.38±8.99 ^b	58.36±4.28 ^a	68.71±4.76 ^b
2.7 extract	105.77±6.22 ^a	99.76±38.82 ^a	43.19±4.19 ^a	36.83±15.88 ^a
3.6 extract	112.54±5.45 ^b	80.37±29.69 ^a	39.28±4.28 ^a	32.06±16.79 ^a

Note: numbers followed by the same letter within a column indicate no significant difference ($\alpha = 5\%$)

Table 2: Levels of HDL-cholesterol and LDL-cholesterol (mg / dL) white rats at the beginning of treatment and end of treatment

Treatment (mg / kg bw)	HDL (mg/dL)		LDL (mg/dL)	
	Beginning treatment	End of treatment	Beginning treatment	End of treatment
Control	25.46±7.35 ^a	26.21±7.21 ^a	87.38±7.29 ^a	89.58±8.66 ^a
3.6 Statin	37.44±4.67 ^a	38.34±6.28 ^a	81.75±6.47 ^a	73.42±6.39 ^a
1.8 extract	38.21±6.53 ^a	39.82±6.24 ^a	78.56±8.28 ^a	81.24±4.67 ^a
2.7 extract	32.86±7.88 ^a	33.93±7.35 ^a	98.21±7.46 ^a	87.63±7.37 ^a
3.6 extract	36.29±5.78 ^a	41.91±7.43 ^a	82.47±6.43 ^a	67.82±6.43 ^a

Note: numbers followed by the same letter within a column indicate no significant difference ($\alpha = 5\%$)

DISCUSSION

Mundu fruit extraction is done by maceration using methanol solvent. The resulting methanol extract of purple yellow and is estimated to contain compounds that are polar. Based on the phytochemical analysis, the methanol extract containing flavonoids, polyphenols, quinones and alkaloids. Deachathai *et al.* (2005), the content of chemical compounds in the genus *Garcinia* i.e flavonoids, xantrons, triterpenoids, and quinones.

The composition of chemical compounds contained in extracts of methanol is still very much. The methanol extract of the Mundu fruit contains many bioactive compounds, each compound has a different biological activity. Under these conditions, the bioactive compounds contained in the methanol extract of the fruit Mundu can interact hypolipidemic produce a complex mechanism.

According Deachathai *et al.*, 2005, the Mundu fruit contains flavonoids, xantrons and quinones. This compound is expected to have an activity that can lower total cholesterol and triglycerides (Qudsiyah & Ward, 2007). Total cholesterol and triglycerides in the control treatment tended to be increased. This condition may occur due to the intake of carbohydrates that will induce the synthesis of fatty acids by fatty acid synthase enzyme into triglycerides.

Simvastatin (statin derivative) is hypolipidemic drug that inhibits cholesterol synthesis in the liver. Statins have active functional groups in the form of the lactone ring which is a dihydroxy acid as its active form. In the body of the lactone ring functional groups will be hydrolyzed to produce dihydroxy acid form of active compounds that

can inhibit the enzyme competitively with HMG-CoA reductase. According to Kawano *et al.* (2007), in addition to works by blocking the synthesis of endogenous cholesterol, statins can also increase the activity of the enzyme lipoprotein lipase (LPL). LPL enzyme is able to hydrolyze triglycerides into free fatty acids, and causes a decrease in blood triglycerides. The mechanism of reduction in triglyceride levels may be related to the inhibition of lipolysis in the liver and inhibition of endogenous cholesterol biosynthesis. Total cholesterol cannot give an exact picture of the effect of the methanol extract of the fruit Mundu.

Total cholesterol is cholesterol from food (chylomicrons), VLDL, LDL and HDL. Under these conditions, the mechanism of reduction in total cholesterol levels correlated with more specific lipoproteins, such as HDL-cholesterol and LDL-cholesterol.

Conditions are expected because of high cholesterol levels in the liver resulting in liver reduces the synthesis of LDL receptors remove excess cholesterol hepatocytes and hepatocytes in the form of VLDL into the blood stream. Catabolism of VLDL in the blood lead levels into LDL. This process causes the levels of LDL-cholesterol to rise (Felig and Frohman, 2001).

Simvastatin is a hypolipidemic drug that works by blocking an enzyme standard of HMG-CoA reductase. Inhibition of this enzyme causes a decrease in VLDL synthesis and increasing LDL receptors in the liver. Decrease in VLDL synthesis which brings triglycerides indirectly cause a decrease in blood triglyceride levels, whereas the increase in LDL receptors in the liver causes a decrease in blood levels of LDL-cholesterol (Neal, 2005). Another

consequence of the addition of simvastatin is synthesized VLDL cholesterol ester poor heart that this VLDL cholesterol esters from HDL to get. This process can lead to decreased levels of HDL-cholesterol following addition of simvastatin.

Methanol extract containing flavonoids, xanton, triterpenoids, and quinone groups are estimated to contain lactone ring. In the body, the active lactone groups will be hydrolyzed to form dihydroxy acid, the active compounds that can inhibit the enzyme competitively with HMG-CoA reductase. Barriers occur because dihydroxy acid form similar to the form of HMG-CoA reductase.

The mechanism of the biological activity of compounds of the research work by inhibiting the enzyme HMG-compete with CoA reductase. HMG-CoA reductase is an enzyme that catalyzes the conversion of 3-hydroxy-3-methyl-glutaril CoenzimA to mevalonic acid, one of the important stages in the pathway of cholesterol synthesis. Barriers of this enzyme causes an increase in the density of LDL receptors in the liver cells resulting in decreased levels of cholesterol, LDL cholesterol and increase HDL number.

From this study it can be concluded that the methanol extract of Mundu fruit (*Garcinia dulcis*) can reduce total cholesterol, triglycerides and LDL-cholesterol and the methanol extract of Mundu fruit (*Garcinia dulcis*) can increase levels of HDL-cholesterol.

ACKNOWLEDGMENTS

Acknowledgements submitted to DP2M the Directorate General of Higher Education for funding this research.

REFERENCES

- Adaramoye, O.A., Akintayo, O., Achem, J., Fafunso, M.A., 2008. Lipid-lowering effects of methanolic extract of *Vernonia amygdalina* leaves in rats fed on high cholesterol diet. *Vascular Health and Risk Management* 4: 235-241.
- Adaramoye, O.A., and Arisekola, M., 2013. Kolaviron, a biflavonoid complex from *Garcinia kola* seeds, ameliorates ethanol-induced reproductive toxicity in male wistar rats. *Nigerian Journal of Physiological Science* 28: 009-015.
- Anonim, 1993. Penapisan Farmakologi, Pengujian Fitokimia dan Pengujian Klinik, Yayasan Pengembangan Obat Bahan Alam, *Phytomedica*. Jakarta.
- Asano, J., Chiba, K., and Tada, M., 1996. Cytotoxic xanthenes from *Garcinia hanburyi*. *Journal of Phytochemistry* 41: 815-820.
- Azima, F., Muchtadi, D., Zakaria, F.R., and Priosoeryanto, B.P., 2004. Potensi Anti-Hiperkolesterolemia Ekstrak Cassia Vera (*Cinnamomum burmanni* Nees ex Blume). *Jurnal Teknologi dan Industri Pangan* 15:145-152.
- Deachathai, S.W., Mahabusarakam, Paichit, S.P., Taylor, W.C., Zhangarid, Y.J., and Yang, C.R., 2005. Phenolic Compound from The Flowers of *Garcinia dulcis*. *Journal of Phytochemistry* 66: 2368-2375.
- Delgado, J.N., and Remers, W.A., 1991. *Wilson and Gisvold's Textbook of Organic Medicinal and Pharmaceutical Chemistry*. 9th Edition. JB Lippincott Co. Philadelphia 8-9.
- Dubey, A.K., Devi, A., Kutty, G., and Shangkar, R.P., 2005. Hypolipidemic Activity of *Ginkgo biloba* Extract EGB 761 in Hypercholesterolemic Wistar Rat. *Iranian Journal of Pharmacology and Therapeutics* 4:9-12.
- Felig, P., and Frohman, L.A., 2001. *Endocrinology and Metabolism*, 4th edition, McGraw-Hill Inc.
- Fukuyama, Y., Minami, H., and Kuwayama, A., 1998. Garsubellins, poly-isoprenylated phloroglucinol derivatives from *Garcinia subelliptica*. *Journal of Phytochemistry* 49: 853-857.
- Gad, S.C., and Chengelis, C.P., 1992. *Animal Models In Toxicology*, Marcel Dekker Inc, New York.
- Iinuma, M., Tosa, H., Ito, T., Tanaka, T., and Riswan, S., 1996. Three new benzophenone-xanthenes dimers from the root of *Garcinia dulcis*. *Chemical and Pharmaceutical Bulletin* 44, 1744-1747.
- Iwu, M.M., 1986. *Biflavonones of Garcinia: Pharmacological and biological activities*. In Cody, V., Middleton, E., and Harbone, J.B., (Eds). *Plant Flavonoids in Biology and Medicine*, Alan R. Liss, New York, 485-488.
- Iwu, M.M., Igbo, O.A., Okunji, C.O., and Tempesta, M.S., 1990. Antidiabetic and aldose reductase activities of biflavonones of *Garcinia kola*. *Journal of Pharmacology* 42: 290-292.
- Kawano, H., Nagata, T., Narahara, M., Kanazawa, M. and Miyake, M., 2007. Triglycerides Accumulation by Peroxisome Proliferator In Rat, *Biological and Pharmaceutical Bulletin* 30:627-632.
- Kosela, S., Hu, L.H., Rachmatia, T., Hanafi, M., and Sim, K.Y., 2000. Dulxanthone F-H, Three New Pyranoxanthenes from the *Garcinia dulcis*. *Journal of Natural Product* 63: 406-407.
- Likhitwitayawuid, K., Chanmahasathien, W., Ruangrungsi, N., and Krungkrai, J., 1998. Xanthone with Antimalarial Activity from *Garcinia dulcis*. *Planta Medica* 64: 281-282.
- Lin, Y.M., Anderson, H., Flavin, M.T., Paik, Y.H., Mata-Greenwood, E., Pengsupar, P.T., Pezzuto, J.M., Schinazi, R.F., Hughes, S.H., and Chen, F.C., 1997. In vitro anti-HIV activity of biflavonoids isolated from *Rhus succedanea* and *Garcinia multiflora*. *Journal of Natural Product* 60: 884-888.
- Lin, K.W., Huang, A.M., and Yang, S.C., 2012. Cytotoxic and antioxidant constituents from *Garcinia subelliptica*. *Journal of Food Chemistry* 135: 851-859.
- Minami, H., Takahashi, E., Fukuyama, Y., Kodama, M., Yoshizawa, T., Nakagawa, 1995. Novel xanthenes with superoxide scavenging activity from *Garcinia subelliptica*. *Chemical and Pharmaceutical Bulletin* 43: 347-349.
- Neal, M.J., 2005. *At a Glance Farmakologi Medis*, edisi ke-5. Penerbit Erlangga. Jakarta.
- Olaiya, C.O., and Omolekan, T.O., 2013. Antihypercholesterolemic activity of ethanolic extract of in rats. *African Health Sciences* 13: 1084-1090.
- Qudsiyah, H dan Wardoyo, E.R.P., 2007. *Pemanfaatan Batang Mundu (Garcinia dulcis (Roxb.) Kurz) Sebagai Antihiperkolesterolemia*, Laporan Penelitian, Universitas Tanjungpura Pontianak.
- Sukpondma, Y., Vatcharin, R., and Phengpaich, S., 2005. Xanthone and Sesquiterpene Derivatives from the Fruits of *Garcinia scortechinii*. *Journal of Natural Product* 68: 1010-1017.
- Sweeney, P.W., 2008. Phylogeny and floral diversity in the genus *Garcinia* (Clusiaceae) and relatives. *International Journal of Plant Science* 169: 1288-1303.
- Wulandari, D., Suarsih, H., Hadatin, S., Widiyanoro, A., Kusmawati, A., dan Wardoyo, E.R.P., 2011. Efek Ekstrak Metanol Buah Makasar (*Brucea javanica* (L.) Merr) terhadap Profil Lipoprotein dan Triglicerida Tikus Putih (*Rattus norvegicus* L.) Hiperlipidemia. *Berkala Penelitian Hayati* 6D: 25-29.
- Yokozawa, T., Cuo, E.J., Sasaki, S., Satou, A., Okamoto, T., and Sei, Y., 2006. The Protective Role of Chinese Prescription Kangen-Karyu Extract on Diet-Induced Hypercholesterolemia in Rats. *Biological and Pharmaceutical Bulletin* 29:760-765.
- Zhang, L.J., Chiou, C.T., and Cheng, J.J., 2010. Cytotoxic polyisoprenyl benzophenonoids from *Garcinia subelliptica*. *Journal of Natural Product* 73: 557-562.

The effect of mundu fruit *Garcinia dulcis* Roxb Kurz methanol extract on lyphoprotein profile and trygliseride white rat *Rattus norvegicus* L

ORIGINALITY REPORT

6%

SIMILARITY INDEX

MATCH ALL SOURCES (ONLY SELECTED SOURCE PRINTED)

★F. Delgado-Vargas. "Natural Pigments: Carotenoids, Anthocyanins, and Betalains -- Characteristics, Biosynthesis, Processing, and Stability", Critical Reviews in Food Science and Nutrition, 5/1/2000	1%
--	----

Crossref

EXCLUDE QUOTES	OFF	EXCLUDE MATCHES	OFF
EXCLUDE BIBLIOGRAPHY	OFF		

FORMAT PENILAIAN (VALIDASI & PEER REVIEW)
LEMBAR
HASIL PENILAIAN SEJAWAT SEBIDANG ATAU *PEER REVIEW*
KARYA ILMIAH : JURNAL ILMIAH

Jurnal Artikel Ilmiah : The effect of munda fruit *Garcinia dulcis* Roxb Kurz methanol extract on
 lyphoprotein profile and trygliseride white rat *Rattus norvegicus* L

Penulis Artikel Ilmiah : Arum Setiawan

Identitas Jurnal Artikel Ilmiah : a. Nama Jurnal : Berkala Penelitian Hayati : Journal of
 Biological Researches
 b. Nomor/Volume/Hal : 2/19/74-77
 c. Edisi (bulan/tahun) : Juni/2014
 d. Penerbit : The East Java Biological Society
 e. Jumlah Halaman : 4

Kategori Publikasi Jurnal Ilmiah : ☒ Jurnal Ilmiah Internasional Bereputasi
 (beri \checkmark pada kategori yang tepat) ☐ Jurnal Ilmiah Internasional
☐ Jurnal Ilmiah Nasional Terakreditasi S1, S2
☐ Jurnal Ilmiah Nasional Terakreditasi S3, S4
☐ Jurnal Ilmiah Nasional Tidak Terakreditasi.

I. Hasil Penilaian Validasi :

No.	ASPEK	URAIAN/KOMENTAR PENILAIAN
1.	Indikasi Plagiasi	6 %
2.	Linearitas	Paper kurang linier dengan bidang biologi konservasi

II. Hasil Penilaian *Peer Review* :

Komponen Yang Dinilai	Nilai Maksimal Jurnal Ilmiah (isikan di kolom yang sesuai)					Nilai Akhir Yang Diperoleh
	Internasional Bereputasi (Maks 40)	Internasional (Maks 20)	Nasional Terakreditasi S1, S2 Maks 25	Nasional Terakreditasi S3, S4 Maks 20	Nasional tidak Terakreditasi (maks 10)	
Kelengkapan dan Kesesuaian unsur isi jurnal (10%)	3					3
Ruang lingkup dan kedalaman pembahasan (30%)	9					8
Kecukupan dan Kemutakhiran data/informasi dan metodologi (30%)	9					8
Kelengkapan unsur dan kualitas penerbit (30%)	9					9
Total = (100%)	30					29
Kontribusi Pengusul (Penulis Pertama /Anggota Utama)	Penulis Utama (0,6x29)=17,4					17,4

KOMENTAR/ULASAN *PEER REVIEW*

Kelengkapan dan Kesesuaian Unsur:	Paper terkait pengaruh buah munda terhadap profil liporproteoin. Isi paper sudah memenuhi kaidah-kaidah karya ilmiah, namun kurang sesuai dengan bidang biologi konservasi.
Ruang Lingkup dan Kedalaman Pembahasan:	Hasil penelitian dibahas secara komprehensif dengan penyampaian pembandingan dari temuan-temuan penelitian lainnya dan teori terkait. Referensi yang diacu dalam pembahasan sudah cukup update untuk bidang kajian ini.
Kecukupan & Kemutakhiran Data & Metodologi:	Data-data hasil penelitian cukup baik dan didukung peta lokasi sampling dan gambar yang ditampilkan menarik. Data didapatkan dengan menggunakan metode yang standard tidak terlalu mutakhir.
Kelengkapan Unsur & Kualitas Penerbit:	Penerbit The East Java Biological Society berkualitas baik, tidak termasuk predatory publisher, dan jurnal tidak terindeks di scopus.

Surabaya, 15 Mei 2020

Penilai 1

A handwritten signature in black ink, featuring a large, stylized initial 'H' and 'P' followed by a cursive name.

Prof. Hery Purnobasuki, M.Si., Ph.D.

NIP 196705071991021001

Unit Kerja : Jurusan Biologi FST Unair

Bidang Ilmu : Biologi

Jabatan/Pangkat : Guru Besar/ Pembina Utama Madya

FORMAT PENILAIAN (VALIDASI & PEER REVIEW)

LEMBAR

1.25.

HASIL PENILAIAN SEJAWAT SEBIDANG ATAU PEER REVIEW

KARYA ILMIAH : JURNAL ILMIAH

Jurnal Artikel Ilmiah : The effect of munda fruit *Garcinia dulcis* Roxb Kurz methanol extract on lyphoprotein profile and trygliseride white rat *Rattus norvegicus* L

Penulis Artikel Ilmiah : Arum Setiawan

Identitas Jurnal Artikel Ilmiah : a. Nama Jurnal : Berkala Penelitian Hayati : Journal of Biological Researchesl

b. Nomor/Volume/Hal : 2/19/74-77

c. Edisi (bulan/tahun) : Juni/2014

d. Penerbit : The East Java Biological Society

e. Jumlah Halaman : 4

Kategori Publikasi Jurnal Ilmiah : ☒ Jurnal Ilmiah Internasional Bereputasi

(beri √ pada kategori yang tepat) ☐ Jurnal Ilmiah Internasional

☐ Jurnal Ilmiah Nasional Terakreditasi S1, S2

☐ Jurnal Ilmiah Nasional Terakreditasi S3, S4

☐ Jurnal Ilmiah Nasional Tidak Terakreditasi

I. Hasil Penilaian Validasi :

No.	ASPEK	URAIAN/KOMENTAR PENILAIAN
1.	Indikasi Plagiasi	6 %
2.	Linearitas	V

II. Hasil Penilaian Peer Review :

Komponen Yang Dinilai	Nilai Maksimal Jurnal Ilmiah (isikan di kolom yang sesuai)					Nilai Akhir Yang Diperoleh
	Internasional Bereputasi (Maks 40)	Internasional (Maks 20)	Nasional Terakreditasi S1, S2 Maks 25	Nasional Terakreditasi S3, S4 Maks 20	Nasional tidak Terakreditasi (maks 10)	
Kelengkapan dan Kesesuaian unsur isi jurnal (10%)	3					3
Ruang lingkup dan kedalaman pembahasan (30%)	9					9
Kecukupan dan Kemutakhiran data/informasi dan metodologi (30%)	9					9
Kelengkapan unsur dan kualitas penerbit (30%)	9					9
Total = (100%)	30					30
Kontribusi Pengusul (Penulis Pertama /Anggota Utama)	Berkala Penelitian Hayati : Journal of Biological Researches Vol. 19(2): 74-77 Juni 2014. Terindeks Copernicus. Penulis pertama dari 3 penulis. Nilai maksimal 100%. Nilai pengusul: $0,6 \times 1 \times 30 = 18$					18

KOMENTAR/ULASAN PEER REVIEW

• Kelengkapan dan Kesesuaian Unsur:	Format lengkap dengan unsur yang ada.
• Ruang Lingkup dan Kedalaman Pembahasan:	Ruang lingkup masih dalam bidang ilmu terkait. Pembahasan baik dan mendalam.
• Kecukupan & Kemutakhiran Data & Metodologi:	Data layak sebagai hasil penelitian walau metode tidak ada yang baru.
• Kelengkapan Unsur & Kualitas Penerbit:	Penerbit The East Java Biological Society, penerbit berkualitas.

Yogyakarta, 12 Juli 2020

Penilai 2

tanda tangan

Prof. Dr. Suwarno Hadisusanto

NIP 195411161983031002

Unit Kerja : Fakultas Biologi UGM

Bidang Ilmu : Biologi

Jabatan/Pangkat : Guru Besar/ Pembina Utama Madya