

# Initial lactate levels versus lactate clearance for predicting mortality in sepsis: A prospective observational analytical study

*by* Mayang Indah Lestari

---

**Submission date:** 09-Mar-2021 12:34PM (UTC+0700)

**Submission ID:** 1528162990

**File name:** Malaysia\_Suppl--Pages--Final\_03-02-21.pdf (120.87K)

**Word count:** 3478

**Character count:** 18346

## RESEARCH ARTICLE

## Initial lactate levels versus lactate clearance for predicting mortality in sepsis: A prospective observational analytical study

Mayang Indah Lestari, Rudyanto Sedono, Zulkifli

### Abstract

**Objective:** Lactate is a useful prognostic marker, as its level increases in hypoxic tissue and/or during accelerated aerobic glycolysis due to excessive beta-adrenergic stimulation and decreased lactate clearance. The Surviving Sepsis Campaign Bundle 2018 Update suggested re-measurement of lactate within 2-4 hours so as to conduct/help/administer/introduce lactate-guided resuscitation to reduce mortality due to sepsis. The aim of this study was to compare initial lactate levels and lactate clearance at 4 h of recognition of sepsis as mortality predictors in sepsis.

**Methods:** It was a prospective study performed with ethical approval in a single tertiary care centre. Patients aged 18 years or older who were diagnosed with sepsis by the Sepsis-3 definition were included in the study while patients who were not admitted to the ICU were excluded. Dropout criteria was death of patients within 4 hours of recognition of sepsis. Baseline demographic data was obtained and subjects were treated with an hour-1 bundle and examined for initial lactate levels. At 4 hours, lactate was re-measured and patients were observed for 28 days then after. Lactate clearance was calculated by the following formula:

$$\frac{(\text{initial lactate} - \text{hour-4 lactate})}{\text{initial lactate}} \times 100.$$

**Results:** Of the 41 subjects included in the study; 27 died (28-day mortality –65.9%). Age, sex, diagnosis of the patient and Charlson's Comorbidities scores between survivors and non-survivors showed no significant differences. Non-survivors had higher Sequential (sepsis-related) Organ Function Assessment (SOFA) scores ( $11.41 \pm 3.46$  versus  $8.77 \pm 2.92$ ;  $p=0.02$ ). Initial lactate levels and lactate clearance did not differ in prognostic value (AUC 0.67 versus 0.5;  $p=0.086$ ), but initial lactate levels of  $>2$  mmol/L had the greatest sensitivity (81.5%).

**Conclusion:** Initial lactate level and lactate clearance did not differ in predicting mortality in patients with sepsis.

**Keywords:** Lactate, Sepsis, Mortality. (JPMA 71: S-25 [Suppl. 1]; 2021)

### Introduction

Lactate serves as a tissue perfusion marker in sepsis,<sup>1</sup> but the etiology of hyperlactatemia in sepsis is complex. Under normal conditions, the lactate concentration is less than 2 mmol/L,<sup>2</sup> and arterial lactate concentrations are determined by the balance between lactate production and its clearance. In sepsis, as a result of tissue hypoxia and/or high levels of aerobic glycolysis due to excessive beta-adrenergic stimulation, the lactate level increases due to lactate overproduction<sup>1</sup> and decreased clearance.<sup>3</sup>

Hypoxic conditions hamper the process of mitochondrial oxidative phosphorylation, resulting in the accumulation of pyruvate and an increase in lactate production. Gore et al. found increased levels of pyruvate and its oxidated products in patients with sepsis.<sup>4</sup> By contrast, Levraut et al. showed that hyperlactatemia in stable patients with sepsis was caused by a decreased lactate clearance rather

than overproduction.<sup>3</sup> A strong correlation was also been observed between blood lactate concentration and clearance<sup>3</sup> and studies have shown that lactate-guided resuscitation can reduce the mortality rate in patients with sepsis.<sup>5-7</sup>

Despite these differences in etiology, lactate levels can be used as a prognostic marker in patients with sepsis. The Surviving Sepsis Campaign Bundle 2018 Update suggested re-measurement of lactate levels within 2-4 hours of recognition of sepsis to reduce mortality.<sup>1</sup> A previous study by Ryoo et al. found that the initial lactate levels had a prognostic value significantly higher than that of lactate clearance at 6 hours of recognition of sepsis.<sup>8</sup> However, our current knowledge about lactate as a prognostic marker is limited since no study has yet compared initial lactate levels and lactate clearance at 4 hours of recognition of sepsis in terms of predicting mortality in patients with sepsis.

This study analysed all patients with sepsis who met the Sepsis-3 definition and our study inclusion criteria. All subjects were treated with an hour-1 bundle for sepsis.

The aim of the study was to compare the initial lactate

.....<sup>33</sup>.....  
Department of Anesthesiology and Intensive Care, Universitas Indonesia,  
Rumah Sakit Cipto Mangunkusumo, Jakarta, Indonesia.

**Correspondence:** Mayang Indah Lestari. Email: mayangindah@fk.unsri.ac.id

levels and lactate clearance at 4 hours of recognition of sepsis as predictors of mortality due to sepsis.

## Patients and Methods

It was a prospective observational analytical study conducted in a single tertiary care centre after ethical approval from Health Research Review Committee of Mohammad Hoesin Central Hospital and Faculty of Medicine, Sriwijaya University, Palembang, Indonesia, in the time duration between February till June 2019.

Patients aged 18 years or older and diagnosed with sepsis using the Sepsis-3 definition were included while patients who were not admitted to the ICU were excluded from the study. Dropout criteria was death within 4 hours of recognition of sepsis. Trained investigators, under intensivist supervision, identified all eligible patients for enrolment in this study. Sample size was calculated by purposive sampling technique.

Baseline demographic data was measured for all subjects including age, sex, diagnosis of the patient, Charlson's Comorbidity score and Sequential Organ Failure Assessment (SOFA) score.

The treatment protocol started with using the hour-1 bundle of sepsis which included 1) measuring initial/basal lactate levels; 2) obtaining blood for blood culture examination in less than one hour before antibiotics administration 3) administering intravenous broad-spectrum antibiotics; 4) rapid administration of 30 mL/kg BW crystalloid and /or lactate (4 mmol/L or more) for conditions of hypotension.; 5) and introducing vasopressors during or after fluid resuscitation if a compensatory hypotension occurs to maintain Mean Arterial Pressure (MAP) of  $\geq 65$  mmHg.<sup>1</sup> Lactate was re-measured at 4 hours. A microbiological examination was also conducted to record the source of infection. Patients were observed for 28 days, for outcomes; survival or non-survival (death). Lactate clearance was calculated using the formula  $([\text{initial lactate} - \text{hour-4 lactate}] / \text{initial lactate}) \times 100$ . Lactate was measured using a Stat Profile pHOx<sup>®</sup> Series blood gas analyser (Nova Biomedical, Waltham, U.S.).

Data was expressed as mean  $\pm$  standard deviation or median (min-max) for continuous variables (depending on the data distribution) and numbers (percentages) for categorical variables. A normality test was carried out using the Shapiro-Wilk test. subjects' characteristics were analysed using the  $\chi^2$  test to compare categorical variables, the independent t-test to compare normally distributed continuous variables and the Mann-Whitney test to compare non-normally distributed continuous

variables or categorical variables which did not meet the  $\chi^2$  test criteria. The performance was evaluated using the Receiver Operating Characteristic (ROC) curves with the Area Under the Curve (AUC). The significance level was 5%. Data was analysed using SPSS Statistics version 21.0 (IBM, New York, U.S.).

## Results

The aim of this study was to compare lactate levels and lactate clearance 4 hours after recognition of sepsis as predictors of mortality due to sepsis. A total of 44 patients met the inclusion criteria and were enrolled in our study. Three patients were dropped out due to death within 4 h of sepsis recognition, leaving a total of 41 subjects whose data were analysed in this study.

The patients in our study had a 28-day mortality rate of 65.9% (27 subjects died). No considerable differences were evident between non-survivors and survivors in terms of age, sex, diagnosis of the patient, microbiology results or comorbidity score based on Charlson classification (Table-1). The only variable that showed a clear difference between non-survivors and survivors was the SOFA score. The initial lactate levels were higher in non-survivors than in survivors (2.7 [0.5-14.7] versus 2.5 [1-8]), although the difference was not statistically significant ( $p = 0.16$ ).

We analysed Area Under the Receiver Operating

Table-1: Subject characteristics.

Variable	Total N = 41 (100%)	Non-survivor N = 27 (65.9%)	Survivor N = 14 (34.1%)	p-value
Age	50.88 $\pm$ 2.35	51.44 $\pm$ 15.28	49.79 $\pm$ 15.08	0.974
<b>Sex</b>				
Female	22 (53.7%)	13 (48.1%)	9 (64.3%)	0.51
Male	19 (46.3%)	14 (51.9%)	5 (35.7%)	
<b>Diagnosis</b>				
Medical	27 (65.9%)	17 (63%)	10 (71.4%)	0.73
Surgical	14 (34.1)	10 (37%)	4 (28.6%)	
<b>Microbiology</b>				
Gram-positive	2 (4.9%)	2 (7.4%)	0 (0%)	0.53
Gram-negative	21 (51.2%)	8 (29.6%)	13 (92.9%)	
Fungal	1 (2.4%)	0 (0%)	1 (7.1%)	
Other	17 (41.5%)	17 (63%)	0 (0%)	
<b>Charlson's Comorbidities score</b>				
0	3 (7.3%)	1 (3.7%)	2 (14.3%)	0.53
2-Jan	29 (70.7%)	20 (74.1%)	9 (64.3%)	
4-Mar	5 (12.2%)	3 (11.1%)	2 (14.3%)	
>5	4 (9.8%)	3 (11.1%)	1 (7.1%)	
SOFA score*	10.55 $\pm$ 0.55	11.41 $\pm$ 3.46	8.77 $\pm$ 2.92	0.02
Initial lactate levels	2.7 (0.5-14.7)	3.1 (0.5-14.7)	2.5 (1-8)	0.16
Lactate clearance	8.33 (-350-70.15)	8.7 (-350-70.15)	5.19(-76.47-62.5)	0.92

\*SOFA: Sequential (sepsis-related) Organ Function Assessment.

**Table-2:** Comparison of initial lactate levels and lactate clearance at 4 h for predicting mortality in septic patients.

Variable	Area under the Receiver Operating Curve (95% CI)	Confidence Interval (CI) 95%		p-value
		Lower	Upper	
Initial Lactate	0.67	0.490	0.840	0.086
Lactate clearance	0.5			

Characteristic (ROC) curves (AUC) to confirm accuracy of initial lactate levels or lactate clearance in predicting mortality in patients with sepsis (Figure-1). The initial lactate performance was poor (AUC 0.67), and lactate clearance (AUC 0.5) failed to predict mortality.

We also compared initial lactate levels and lactate clearance at 4 h for predicting mortality due to sepsis (Table-2), but the difference between these two variables was not statistically significant.

The ROC shown in Table-3 indicated that the variable with the higher sensitivity for predicting mortality due to sepsis was an initial lactate level greater than 2 mmol/L, at 81.5% which meant that an initial lactate level of more than 2 mmol/L could correctly predict the death of patients with sepsis within 28 days 81.5% of the time. The variable with the highest specificity was an initial lactate

**Table-3:** Sensitivity and specificity of initial lactate levels and lactate clearance at 4 hour for predicting mortality in septic patients.


Variable	Sensitivity	Specificity
Initial Lactate (mmol/L)		
>2	81.5%	35.7%
>3	48.1%	64.3%
>4	37%	85.7%
Lactate clearance (%)		
<10%	44.4%	57.1%
<20%	33.3%	64.3%
<30%	25.9%	78.6%

level greater than 4 mmol/L, at 85.7%. meaning that an initial lactate level of more than 4 mmol/L could correctly predict the survival of patients with sepsis within 28 days, 85.7% of the time.

**Discussion**

In critically ill patients, and especially those who suffer from shock, hyperlactatemia is normally interpreted as an anaerobic metabolism marker indicating inadequate oxygen delivery. In sepsis, the etiology of hyperlactatemia is particularly complex. For example, Ronco et al. demonstrated that increased oxygen delivery fails to decrease lactate levels in patients with sepsis having hyperlactatemia.<sup>9</sup> Similarly, Gattinoni et al. found that supra-normal oxygen delivery in sepsis, as popularised by Shoemaker et al.,<sup>10</sup> did not improve patient outcomes.<sup>11</sup> Apart from its association with hypoxia, hyperlactatemia is also related to many other metabolic processes and other clinical conditions, such as lactate clearance. Gattinoni et al. showed that a more frequent etiology of hyperlactatemia in sepsis is a diminished oxygen utilisation by tissues.<sup>12</sup>

The arterial lactate level is determined by the balance between lactate production and its clearance. Despite their different aetiologies, lactate levels remain clinically important and can be used as a prognostic marker. For example, Mahmoodpoor et al. indicated that patients with sepsis who died within 28 days had higher serum lactate levels at the time of arrival and



**Figure:** AUC of initial lactate and lactate clearance for sepsis mortality prediction performance.

required a higher dose of vasopressor.<sup>13</sup> Similarly, Lokhandwala et al. showed that lactate levels  $\geq 4$  mmol/L at 6 h after the initial examination and lactate reductions less than 20% were associated with an increased risk of in-hospital mortality due to sepsis.<sup>14</sup> Nguyen et al. found that patients with a higher lactate clearance had significantly decreased in-hospital and 28 or 60-day mortalities.<sup>15</sup> The Surviving Sepsis Campaign Bundle 2018 Update suggested re-measurement of lactate within 2-4 h of recognition of sepsis for lactate-guided resuscitation to reduce mortality.<sup>1</sup>

In our study, no significant differences were noted in age, sex, diagnosis of patient or Charlson's Comorbidities score between survivors and non-survivors, but the SOFA scores were higher in non-survivors (11.41 $\pm$ 3.46 versus 8.77 $\pm$ 2.92;  $p = 0.02$ ). Lie et al. also indicated a higher total SOFA score in non-survivors than in survivors (6.7 $\pm$ 3.8 versus 4.6 $\pm$ 2.9;  $p < 0.001$ ).<sup>16</sup>

In this study, neither lactate level nor lactate clearance differed between survivors and non-survivors. This result contrasted with a retrospective study by Ryoo et al., who found a lower median 6 h lactate level and a higher lactate clearance in survivors than in non-survivors.<sup>8</sup> This inconsistency between these study findings probably reflected the different patient populations (sepsis versus septic shock). In this study, we included all patients with sepsis diagnosed by using the Sepsis-3 definition, which included those patients who had initial lactate levels of 2 mmol/L or less.

The initial lactate levels and the lactate clearance at 4 h of recognition of sepsis did not differ in prognostic value (AUC 0.67 versus 0.5;  $p = 0.086$ ). Unlike the previous retrospective study, Ryoo et al. found that initial lactate levels had a significantly better prognostic value than lactate clearance at 6 h of recognition of sepsis (AUC 0.70 versus 0.65;  $p < 0.01$ ).<sup>8</sup> Singh et al. showed an AUC for serum lactate at 0 h of 0.734 and a lactate clearance at 48 h of 0.91.<sup>17</sup> The inconsistency among these studies probably reflects differences in the timing of the lactate measurements or in the time frame used to calculate the lactate clearance.

An initial lactate level of more than 2 mmol/L had the highest sensitivity for predicting mortality due to sepsis (81.5%). In this study, we also found that an initial lactate level of more than 4 mmol/L had a specificity of 85.7% and sensitivity of 37%. This result was consistent with the findings of Fernando et al., who concluded that an initial lactate level of 4 mmol/L or greater had a specificity of 97% and a sensitivity of 27% in predicting the deterioration of patients with sepsis.<sup>18</sup> Ryoo et al. also

found that a lactate level above 2 mmol/L had the greatest sensitivity (85.3%) and a level above 4 mmol/L had the greatest specificity (72.3%).<sup>8</sup>

## Limitations

Our study had several limitations. One limitation is that we did not examine lactate 2 h after recognition of sepsis, as suggested in the Surviving Sepsis Campaign Bundle 2018 Update, which recommends re-measurement of lactate within 2-4 h. A second limitation is that we included all patients with sepsis diagnosed by using the Sepsis-3 definition, which included those who had lactate levels of 2 mmol/L or less. A third limitation was that this was a single-centre study, which cannot be generalized to other institutions and populations.

## Conclusion

The SOFA score, which serves as a life-threatening organ dysfunction indicator in sepsis, was higher in non-survivor patients. Initial lactate levels and lactate clearance after 4 hrs of recognition of sepsis did not differ as predictors of mortality in patients with sepsis. Both initial lactate levels and lactate clearance were found to be poor predictors of mortality in patients with sepsis. Few inconsistencies among previous studies and our findings probably reflected differences in patient populations (sepsis versus septic shock) and in the lactate measurement times used to calculate lactate clearance. Our study limitations necessitate further investigations.

**Acknowledgement:** We thank Mohammad Hoesin Hospital, Palembang for allowing us to conduct this research.

**Disclaimer:** The abstract has been previously presented in 11th Malaysian Indonesian Brunei Medical Sciences Conference 2019, and the manuscript was a part of a research paper

**Conflict of Interest:** This article has no conflict of interest.

**Funding Disclosure:** There is no funding disclosure for this article.

## References

1. Levy MM, Evans LE, Rhodes A. The Surviving Sepsis Campaign Bundle: 2018 update. *Intensive Care Med* 2018; 44:925-928. doi: 10.1007/s00134-018-5085-0.
2. Levy B. Lactate and shock state: the metabolic view. *Curr Opin Crit Care* 2006; 12:315-21. doi: 10.1097/01.ccx.0000235208.77450.15.
3. Levraut J, Ciebiera JP, Chave S, Rabary O, Jambou P, Carles M, et al. Mild hyperlactatemia in stable septic patients is due to impaired lactate clearance rather than overproduction. *Am J Respir Crit Care Med* 1998; 157:1021-6. doi: 10.1164/ajrccm.157.4.9705037.
4. Gore DC, Jahoor F, Hibbert JM, DeMaria EJ. Lactic acidosis during sepsis is related to increased pyruvate production, not deficits in

- tissue oxygen availability. *Ann Surg* 1996; 224:97-102. doi: 10.1097/0000658-199607000-00015.
5. Jansen TC, van Bommel J, Schoonderbeek FJ, Sleswijk Visser SJ, van der Klooster JM, Lima AP, et al. Early lactate-guided therapy in intensive care unit patients: a multicenter, open-label, randomized controlled trial. *Am J Respir Crit Care Med* 2010; 182:752-61. doi: 10.1164/rccm.200912-1918OC.
  6. Jones AE, Shapiro NI, Trzeciak S, Arnold RC, Claremont HA, Kline JA. Lactate clearance vs central venous oxygen saturation as goals of early sepsis therapy: a randomized clinical trial. *JAMA* 2010; 303:739-46. doi: 10.1001/jama.2010.158.
  7. Ryou SM, Lee J, Lee YS, Lee JH, Lim KS, Huh JW, et al. Lactate Level Versus Lactate Clearance for Predicting Mortality in Patients With Septic Shock Defined by Sepsis-3. *Crit Care Med* 2018; 46:e489-95. doi: 10.1097/CCM.0000000000003030.
  8. Ronco JJ, Fenwick JC, Wiggs BR, Phang PT, Russell JA, Tweeddale MG. Oxygen consumption is independent of increases in oxygen delivery by dobutamine in septic patients who have normal or increased plasma lactate. *Am Rev Respir Dis* 1993; 147:25-31. doi: 10.1164/ajrccm/147.1.25.
  9. Shoemaker WC, Appel PL, Waxman K, Schwartz S, Chang P. Clinical trial of survivors' cardiorespiratory patterns as therapeutic goals in critically ill postoperative patients. *Crit Care Med* 1982; 10:398-403. doi: 10.1097/00003246-198206000-00015.
  10. Gattinoni L, Brazzi L, Pelosi P, Latini R, Tognoni G, Pesenti A, et al. A trial of goal-oriented hemodynamic therapy in critically ill patients. SvO2 Collaborative Group. *N Engl J Med* 1995; 333:1025-32. doi: 10.1056/NEJM199510193331601.
  11. Marik PE. Lactate guided resuscitation-nothing is more dangerous than conscientious foolishness. *J Thorac Dis* 2019; 11 (Suppl 15): s1969-72. doi: 10.21037/jtd.2019.07.6.
  12. Mahmoodpoor A, Shadvar K, Saghaleini SH, Koleini E, Hamishehkar H, Ostadi Z, et al. Which one is a better predictor of ICU mortality in septic patients? Comparison between serial serum lactate concentrations and its removal rate. *J Crit Care* 2018; 44:51-6. doi: 10.1016/j.jccr.2017.10.019.
  13. Lokhandwala S, Andersen LW, Nair S, Patel P, Cocchi MN, Donnino MW. Absolute lactate value vs relative reduction as a predictor of mortality in severe sepsis and septic shock. *J Crit Care* 2017; 37:179-84. doi: 10.1016/j.jccr.2016.09.023.
  14. Nguyen HB, Loomba M, Yang JJ, Jacobsen G, Shah K, Otero RM, et al. Early lactate clearance is associated with biomarkers of inflammation, coagulation, apoptosis, organ dysfunction and mortality in severe sepsis and septic shock. *J Inflamm (Lond)* 2010; 7:e6. doi: 10.1186/1476-9255-7-6.
  15. Lie KC, Lau CY, Van Vinh Chau N, West TE, Limmathurotsakul D. Utility of SOFA score, management and outcomes of sepsis in Southeast Asia: a multinational multicenter prospective observational study. *J Intensive Care* 2018; 6:9. doi: 10.1186/s40560-018-0279-7.
  16. Singh S, Bhardawaj A, Shukla R, Jhadav T, Sharma A, Basannar D. The handheld blood lactate analyser versus the blood gas based analyser for measurement of serum lactate and its prognostic significance in severe sepsis. *Med J Armed Forces India* 2016; 72:325-31. doi: 10.1016/j.mjafi.2016.05.007.
  17. Fernando SM, Bamaby DP, Herry CL, Gallagher EJ, Shapiro NI, Seely AJE. Helpful Only When Elevated: Initial Serum Lactate in Stable Emergency Department Patients with Sepsis Is Specific, but Not Sensitive for Future Deterioration. *J Emerg Med* 2018; 54:766-73. doi: 10.1016/j.jemermed.2018.01.040.

# Initial lactate levels versus lactate clearance for predicting mortality in sepsis: A prospective observational analytical study

## ORIGINALITY REPORT

15%

SIMILARITY INDEX

7%

INTERNET SOURCES

12%

PUBLICATIONS

2%

STUDENT PAPERS

## PRIMARY SOURCES

- 1 Ata Mahmoodpoor, Kamran Shadvar, Seied Hadi Saghaleini, Evin Koleini, Hadi Hamishehkar, Zohre Ostadi, Nader D. Nader. "Which one is a better predictor of ICU mortality in septic patients? Comparison between serial serum lactate concentrations and its removal rate", Journal of Critical Care, 2018  
Publication 1%
- 2 [www.psmid.org](http://www.psmid.org)  
Internet Source 1%
- 3 Lee, Tae Rim, MunJu Kang, Won Chul Cha, Tae Gun Shin, Min Seob Sim, IkJoon Jo, KeunJeong Song, Yeon Kwon Jeong, and Jun Hwi Cho. "Better lactate clearance associated with good neurologic outcome in survivors who treated with therapeutic hypothermia after out-of-hospital cardiac arrest", Critical Care, 2013.  
Publication 1%
- 4 Submitted to Manchester Metropolitan University 1%

5 [www.freepatentsonline.com](http://www.freepatentsonline.com) 1%

Internet Source

---

6 Young A Kim, Eun-Ju Ha, Won Kyoung Jhang, Seong Jong Park. "Early blood lactate area as a prognostic marker in pediatric septic shock", Intensive Care Medicine, 2013 1%

Publication

---

7 [nrronline.org](http://nrronline.org) 1%

Internet Source

---

8 [pbbmi.org](http://pbbmi.org) 1%

Internet Source

---

9 Klein Klouwenberg, Peter MC, Maaik SM van Mourik, David SY Ong, Janneke Horn, Marcus J Schultz, Olaf L Cremer, and Marc JM Bonten. "Validation of a novel surveillance paradigm for ventilator-associated events", Critical Care, 2013. <1%

Publication

---

10 Qin Yin, Cai-jun Wu, Jun Yang, Chen-chen Hang, Chun-sheng Li. "Effects of Shenfu Injection (参附注射液) on cerebral metabolism in a porcine model of cardiac arrest", Chinese Journal of Integrative Medicine, 2016 <1%

Publication

---

11 [test.dovepress.com](http://test.dovepress.com)

<1%

12

Mohammed Badr Salim, Hesham Elaasr, Mervat El Damarawy, Ashraf Wadee, Alaa Ashour, Fatma Mohammad Nasr. "Atrial ejection force and brain natriuretic peptide as markers for mortality in sepsis", The Egyptian Journal of Critical Care Medicine, 2015

Publication

<1%

13

Xianshi Zhou, Ye Ye, Fang Tian, Fanwei Wu. "Lactate levels in arterial and venous blood may be correlated but not equivalent", Journal of Critical Care, 2017

Publication

<1%

14

Bruno Levy. "Lactate and shock state: the metabolic view", Current Opinion in Critical Care, 08/2006

Publication

<1%

15

Cai, J.-Y., J. Sun, and Z.-Q. Yu. "Serum mannose-binding lectin levels after aneurysmal subarachnoid hemorrhage", Acta Neurologica Scandinavica, 2016.

Publication

<1%

16

"The relationship between oxygen delivery and consumption", Disease-a-Month, 199407

Publication

<1%

17

Khie Chen Lie, Chuen-Yen Lau, Nguyen Van Vinh Chau, T. Eoin West, Direk Limmathurotsakul. "Utility of SOFA score, management and outcomes of sepsis in Southeast Asia: a multinational multicenter prospective observational study", Journal of Intensive Care, 2018

Publication

&lt;1%

18

W Abbink. "Parathyroid hormone-related protein and calcium regulation in vitamin D-deficient sea bream (*Sparus auratus*)", Journal of Endocrinology, 06/01/2007

Publication

&lt;1%

19

[www.medsci.org](http://www.medsci.org)

Internet Source

&lt;1%

20

Anthony M Napoli. "The role of lactate clearance in the resuscitation bundle", Critical Care, 2011

Publication

&lt;1%

21

John McNelis, Corrado P Marini, Antoni Jurkiewicz, Samuel Szomstein, H.Hank Simms, Garry Ritter, Ira M Nathan. "Prolonged lactate clearance is associated with increased mortality in the surgical intensive care unit", The American Journal of Surgery, 2001

Publication

&lt;1%

22

Paul N. Severin. "Endotoxin-induced hyperlactatemia results from decreased lactate

&lt;1%

clearance in hemodynamically stable rats",  
Critical Care Medicine, 11/2002

Publication

23

[emedicine.medscape.com](http://emedicine.medscape.com)

Internet Source

<1%

24

[pdfs.semanticscholar.org](http://pdfs.semanticscholar.org)

Internet Source

<1%

25

[www.labdeurgencias.com.ar](http://www.labdeurgencias.com.ar)

Internet Source

<1%

26

[www.nejm.org](http://www.nejm.org)

Internet Source

<1%

27

Bannert, Erik, Tanja Tesch, Jeannette Kluess, Jana Frahm, Susanne Kersten, Stefan Kahlert, Lydia Renner, Hermann-Josef Rothkötter, and Sven Dänicke. "Metabolic and Hematological Consequences of Dietary Deoxynivalenol Interacting with Systemic Escherichia coli Lipopolysaccharide", Toxins, 2015.

Publication

<1%

28

Chang Hwan Oh, Jong Won Kim, Geon Ha Kim, Kyeong Ryong Lee, Dae Young Hong, Sang O Park, Kwang Je Baek, Sin Young Kim. "Serum Lactate Could Predict Mortality in Patients With Spontaneous Subarachnoid Hemorrhage in the Emergency Department", Frontiers in Neurology, 2020

<1%

29

Daniel Backer. "Lactic acidosis", Applied Physiology in Intensive Care Medicine, 2006

Publication

---

<1%

30

Ruud G. Nijman, Rikke Jorgensen, Michael Levin, Jethro Herberg, Ian K. Maconochie. "Management of Children With Fever at Risk for Pediatric Sepsis: A Prospective Study in Pediatric Emergency Care", Frontiers in Pediatrics, 2020

Publication

---

<1%

31

Sanna Lahtinen, Juho Nurkkala, Samuli Hannula, Pasi Ohtonen, Petri Koivunen, Janne H. Liisanantti. "Perioperative risk factors for one-year mortality in patients with free flap reconstruction due to cancer of the head and neck", Journal of Oral and Maxillofacial Surgery, 2021

Publication

---

<1%

32

Vered Molina-Hazan, Gideon Paret. "Chapter 8 The Physiology and the Clinical Significance of Postoperative Hyperlactatemia After Pediatric Cardiac Surgery", IntechOpen, 2012

Publication

---

<1%

33

saripediatri.org

Internet Source

---

<1%

34

Internet Source

<1%

---

35

[www.ccjm.org](http://www.ccjm.org)

Internet Source

<1%

---

36

Etienne Macedo, Isac Castro, Luis Yu, Regina R.C. Abdulkader, José M. Vieira. "Impact of Mild Acute Kidney Injury (AKI) on Outcome after Open Repair of Aortic Aneurysms", Renal Failure, 2009

Publication

<1%

---

37

[ecommons.aku.edu](http://ecommons.aku.edu)

Internet Source

<1%

---

38

María F. Escobar, María P. Echavarría, María A. Zambrano, Isabella Ramos, Juan P. Kusanovic. "MATERNAL SEPSIS", American Journal of Obstetrics & Gynecology MFM, 2020

Publication

<1%

---

39

Won Seok Yang, Hui Dong Kang, Sang Ku Jung, You Jin Lee, Se Hyun Oh, Youn-Jung Kim, Chang Hwan Sohn, Won Young Kim. "A Mortality Analysis of Septic Shock, Vasoplegic Shock, and Cryptic Shock Classified by the Third International Consensus Definitions (Sepsis-3)", The Clinical Respiratory Journal, 2020

Publication

<1%

---

40

N/A. "Practice parameters for hemodynamic support of sepsis in adult patients in sepsis",  
Critical Care Medicine, 03/1999

Publication

<1%

---

Exclude quotes      On

Exclude matches      Off

Exclude bibliography      On