

Relationship of Food Leftover with Covid-19 Inpatients Recovery Speed in Palembang Indonesi

by Ramzi Amin

Submission date: 16-Nov-2021 07:13PM (UTC-0600)

Submission ID: 1705088699

File name: ith_Covid-19_Inpatients_Recovery_Speed_in_Palembang_Indonesi.pdf (315.75K)

Word count: 2140

Character count: 11554



Bioscientia Medicina: Journal of Biomedicine & Translational Research

Journal Homepage: www.bioscmed.com

Relationship of Food Leftover with Covid-19 Inpatients Recovery Speed in Palembang Indonesia

Devi Eryanti¹, Chairil Anwar^{2,5*}, Yuanita Windusari^{3,5}, Ramzi Amin², Ahmad Ghiffari^{4,5}

¹Master Program of Public Health, Faculty of Public Health, Universitas Sriwijaya, Palembang, Indonesia

²Faculty of Medicine, Universitas Sriwijaya, Palembang, Indonesia

³Faculty of Public Health, Universitas Sriwijaya, Palembang, Indonesia

⁴Faculty of Medicine, Universitas Muhammadiyah Palembang, Palembang, Indonesia

⁵Faculty of Environment, Universitas Sriwijaya, Palembang, Indonesia

ARTICLE INFO

Keywords:

COVID-19

Food waste

Feeding

Hospitalization

Recovery

*Corresponding author:

Chairil Anwar

E-mail address:

chairil53@fk.unsri.ac.id

15

All authors have reviewed and approved the final version of the manuscript.

<https://doi.org/10.32539/bsm.v5i12.377>

ABSTRACT

Background: Hospital nutrition services support the patient's recovery process, which the activity is feeding. One indicator of quality in food service is food leftover. The high food waste can affect the patient's health. The high food waste will pose a risk of malnutrition in hospitalized patients. The study aimed to analyze the food leftovers in COVID-19 inpatients at the Palembang referral hospital. **Methods:** This cross-sectional quantitative descriptive study was with a sample of 112 patients selected by purposive sampling in February-May 2021. Data were taken using the Comstock form and then translated in the form of percent leftover food. According to the Minister of Health, the minimum leftover food is less than 25 percent. **Results:** The results showed that more than 25 percent of the leftover food were 51 respondents; Thirty-three respondents (54.1%) have fully recovered and 28 respondents (45.9%) still in improvement. On the other hand, the leftovers food of less than 25 percent were 51 respondents; Forty respondents (78.4%) have fully cured, and 11 people (21.6%) still in treatments. The analysis of Chi-square test results shows a p-value of 0.013, showing a significant relationship between food waste and the recovery of COVID-19 patients. **Conclusion:** The food waste of COVID-19 patients in referral hospitals was still relatively high. It is necessary to modify the diet menu to prevent leftovers and speed up the patient's recovery.

1. Introduction

Hospital nutrition services patients adapted to clinical conditions, nutritional status or body metabolism¹. Hospital dining is an activity that starts from menu planning to evaluation to provide quality food according to nutritional needs, costs, safe and acceptable to consumers to achieve optimal nutritional status and accelerate patient recovery². The management of food administration in hospitals is also strongly influenced by the food waste factor as an indicator of a hospital's quality of minimum service

standards³. The food served in the hospital serves to maintain the immune system and help the healing process so that the food served must meet the needs of both quality, quantity, adequacy of nutrition, and the accuracy of the diet⁴.

Measurement of food quantity is by weighing or using a visual estimate. Visual estimates, such as the Comstock method², are easy to do, inexpensive and widely used to assess food waste in large amounts of food administration⁵. The visual methods

demonstrated that the hospital average food waste could easily compare to the weighting scheme⁶. Weighing estimates, such as the waste audit, calculate plate waste for the average meal⁷. It is crucial for assessing food waste in inpatients, especially COVID-19 patients, to evaluate food service activities in hospitals. There are currently no publications regarding food waste at the COVID-19 referral hospital in Palembang City.

2. Methods

This research was conducted at the COVID-19 referral hospital in Palembang city, i.e. Siti Fatimah Regional General Hospital, Ernaldi Bahar Hospital, RK Charitas Hospital, Palembang BARI Hospital, and Moehammad Hoesin Hospital Palembang. Data collection was carried out three times a day at mealtimes (breakfast, lunch and dinner) from February to May 2021. The Lameshow sample size was 112 hospitalized patients by equal number to the regular and special diets. The respondents carried out data collection by filling out a Google form distributed by nurses and the COVID-19 room dietician.

Food leftovers for COVID-19 patients were considered infectious objects and directly disposed of

the infectious waste^{8,9}. The types of food consisted of rice/staple, animal protein, plant protein, fruits and vegetables, water and milk. The classification were; 0% (no leftovers), 25% (quarter portions of food leftovers), 50% (half portions of leftovers), 75% (three-quarter portion leftovers) and 100% (there are whole food leftovers). The Universitas Sriwijaya Faculty of Public Health issued a Certificate of Research Ethics with 43/EC/KBHKI/FK-UMP/XI/2019.

3. Results

The one hundred and twelve respondents who met the research inclusion criteria were aged 18-70 years, cooperative, receiving regular or special diets, and did not use a ventilator. Table 1 shows the characteristics of responders. There were more respondents over the age of 40, more men than women, and more higher education than low education than SMA. Hospital food has various processed variations (Figure 1). The results showed that staple foods had the most leftovers by patients. The association between food waste and the respondent's recovery was shown in Table 2. The p-value of the test was 0.013, less than the significant level of 0.05.

Table 1. Characteristics of respondents based on age, gender and education (n=112)

No	Characteristics	n	%
1	Age		
	<40 years	50	44.5
	40 years	62	55.4
2	Gender		
	Male	59	52.7
	Female	53	47.3
3	Education		
	Lower than senior high school (SMA)	11	9.8
	Higher or equal to SMA	101	90.2




Figure 1. Food leftovers by percentage of food types

Table 2. Relationship between food waste and healing process (n=112)

Leftovers	Recovery				Total		p-value	OR	95%CI
	Full		Repair		n	%			
	n	%	n	%					
Not good (>25%)	33	54.1	28	45.9	61	100	0.013	0.324	0.140-0.748
Good (≤25%)	40	78.4	11	21.6	51	100			
Total	73	65.2	39	34.8	112	100			

4. Discussion

The staple food was left mainly by inpatients. Purwakarta Holistic Hospital uses the standard portion of "little" for the patient's meal menu to deal with the leftover staple food¹⁰. At RSUD dr. Soeratto Gemolong showed that the types and preparations of plant protein affected the patient's assessment. Plant protein was the most left by the patients because some were not fond of tofu and tempeh. The types of plant protein lack plating, which causes patients to be lazy to eat¹¹. Nutrition installations should develop the creativity of processing and cooking foodstuffs such as tofu and tempeh foodstuffs can be spent by patients¹². Others have discovered similar tendencies, with over 40% of serving veggies going uneaten compared to only 18% of entrees. This could be due to hospitalized patients' decreased appetite for veggies due to inadequate cooking practices¹³. Most studies show less leftover food at breakfast (morning meal) than other main meals, although this was not a general finding¹⁴.

The odds ratio in this study was 0.013, which

implies there was a 0.324 times greater probability of healing faster. In 2017 showed that changing rice portions in the patient's diet can reduce food waste ($r=0.804$; $p<0.05$)¹⁰. At the Universitas Muhammadiyah Malang Hospital, the average patient leftover food in 3 meals was 57%¹⁵. The factors influencing the patient to leave food are internal factors (clinical conditions, eating habits, gender) and external factors (food taste, food temperature, texture, food colour, portion and variety of food ingredients, environmental factors/out-of-hospital food)¹⁶. Inpatients experienced changes that can affect food intake or absorption, metabolism, and excretion of nutrients; changes in appetite, senses of taste and smell¹⁷. The COVID-19 patients suffered from sensory disorders called anosmia, either momentary or sedentary, caused by a mechanical blockage inhibiting the smell reaching the nerves and eventually losing the sense of smell¹⁸. The hospitalized COVID-19 patients admitted to the ICU are suffered from anorexia secondary to an infection, dyspnea, dysosmia, dysgeusia, stress, confinement, and organizational problems limiting the desire to eat¹⁹. These issues

substantially impact the COVID-19 patients being treated, resulting in insufficient food intake, wasted food waste, and a slowdown of the patient's recovery ²⁰.

This study had several limitations. Respondent bias can be caused by the demographic of respondents above the age of 40. Respondents may be frustrated by the usage of a Google form to collect data. Hospital policies changes during a pandemic. Due to direct infectious waste management, leftover food bypass the nutrition installation for waste inspection can result in a misjudgment of exact leftover food.

5. Conclusion

The food waste of COVID-19 patients in referral hospitals was still relatively high. While high food waste will affect the time it takes for patients to recover, better food service delivery and diet modification quality are demanded.

6. Acknowledgment

We would like to thank all the referral hospital employees for cooperating in this study, especially the nutritionists, education and training department (Diklat), and COVID-19 treatment rooms nurses. The study was ¹¹ founded by Unggulan Profesi Research Grant ¹¹ Universitas Sriwijaya (SP DIPA-023.17.2.677515/2021), with the Rector's Decree Number of 0014/UN9/SK.LP2M.PT/2021.

7. References

1. Kementerian Kesehatan Republik Indonesia. ¹³ Pedoman pelayanan gizi dan dietetik di rumah sakit darurat dalam penanganan pandemi COVID-19. Jakarta; 2020.
2. Kementerian Kesehatan Republik Indonesia. ¹³ Pedoman pelayanan gizi rumah sakit (PGRS). Jakarta; 2013.
3. Kementerian Kesehatan Republik Indonesia. Keputusan Menteri Kesehatan Republik Indonesia no. 129/Menkes/SK/II/2008 tentang Standar Pelayanan Minimal Rumah Sakit. Jakarta; 2008.
4. Mardianingsih N, Utami FA, Palupi IR. Capaian standar pelayanan minimal gizi di rumah sakit umum daerah (RSUD) Manokwari Papua Barat. ⁸ J Gizi Klin Indones. 2020;16(4):152–67.
5. Williams PG, Walton K. Plate waste in hospitals and strategies for change. ⁸ Eur e-journal Clin Nutr Metab. 2011;6(6):e235–41.
6. Saskia R, Primadona S, Mahmudiono T. ¹⁴ Hubungan tingkat kematangan dan suhu dengan sisa makanan lauk nabati pada pasien anak di ruang inap Rumkital dr. Ramelan Surabaya. ¹⁴ Media Gizi Indones. 2018;13(2):100–7.
7. ¹⁰ Dias-ferreira C, Santos T, Oliveira V. Hospital food waste and environmental and economic indicators – A Portuguese case study. ¹⁰ Waste Manag. 2015;46:146–54.
8. Maalouf A, Maalouf H. Impact of COVID-19 pandemic on medical waste management in Lebanon. ⁵ Waste Manag Res. 2021;
9. ⁵ Das AK, Islam MN, Billah MM, Sarker A. COVID-19 pandemic and healthcare solid waste management strategy – A mini-review. ⁵ Sci Total Environ [Internet]. 2021;778:146220. Tersedia pada: ¹⁸ <https://doi.org/10.1016/j.scitotenv.2021.146220>
10. ¹⁸ Fatkhurohman F, Lestari YN, Torina DT. The relationship of the changing in standard portion toward food waste among holistic hospital patients in 2016 (food waste study of rice on lunch menu in holistic hospital). ¹⁸ Gizi Indones. 2017;40(1):1.
11. ⁷ Ariyanti V, Widyarningsih EN, Rauf R. Hubungan antara karakteristik sensorik makanan dengan sisa makanan biasa pada pasien rawat inap RSUD Dr. Soeratno, Gemolong, kabupaten Sragen. ⁷ J Kesehat. 2017;10(1):17–25.
12. ⁷ Muliani U. Faktor-faktor yang berhubungan dengan sisa makanan saring pasien rawat inap. ⁷ J Keperawatan. 2013;IX(1):31–6.
13. ⁴ Puspita WL, Prawiningdyah Y, Nisa FZ. Penerapan Hazard Analysis Critical Control Point (HACCP) terhadap penurunan bahaya

- mikrobiologis pada makanan khusus anak berbasis hewani di rumah sakit umum daerah dr. Soedarsono Pontianak. *J Gizi Klin Indones.* 2010;7(1):8.
14. Thibault R, Coëf M, Joly F, Bohé J, Schneider SM, Déchelotte P. How the COVID-19 epidemic is challenging our practice in clinical nutrition — feedback from the field. *Eur J Clin Nutr.* 2021;75:407–16.
 15. Karunia Tanuwijaya L, Gresari Sembiring L, Yanuar Dini C, Putri Arfiani E, Arimba Wani Y. Sisa Makanan Pasien Rawat Inap: Analisis Kualitatif. *Indones J Hum Nutr.* 2018;5(1):51–61.
 16. Fadilla C, Rachmah Q, Juwariyah J. Inpatients food waste description at Sidoarjo general hospital. *Amerta Nutr.* 2020;4(3):198–204.
 17. Ronitawati P, Puspita M, Citra K. Faktor-faktor yang berhubungan dengan sisa makanan di rumah sakit umum daerah Koja Jakarta Utara tahun 2017. *Heal Sci Growth.* 2018;3(2):57–76.
 18. Butowt R, Bartheld CS Von. Anosmia in COVID-19: Underlying mechanisms and assessment of an olfactory route to brain infection. *Neurosci.* 2020;00(0):1–22.
 19. Verbeek JH, Rajamali B, Ijaz S, Sauni R, Toomey E, Blackwood B, et al. Personal protective equipment for preventing highly infectious diseases due to exposure to contaminated body fluids in healthcare staff. *Cochrane Database Syst Rev.* 2019;7(7):1–106.
 20. Lironika A, Suryadi MY. The distribution schedule and food taste was correlated with patient's plate waste at obstetric and surgical room in dr. Soebandi hospital, Jember. *Amerta Nutr.* 2019;3(3):194–200.

Relationship of Food Leftover with Covid-19 Inpatients Recovery Speed in Palembang Indonesi

ORIGINALITY REPORT

16%

SIMILARITY INDEX

0%

INTERNET SOURCES

11%

PUBLICATIONS

10%

STUDENT PAPERS

PRIMARY SOURCES

- 1 Nassim Essabah Haraj, Siham El Aziz, Asma Chadli, Asma Dafir et al. "Nutritional status assessment in patients with Covid-19 after discharge from the intensive care unit", *Clinical Nutrition ESPEN*, 2021
Publication 1%
- 2 Renata Karina Reis, Mayra Gonçalves Meneguetti, Silmara Elaine Malaguti-Toffano, Laelson Rochelle Milanês Sousa et al. "Knowledge, Behaviors, and Perceptions of Risk of COVID-19 Among Brazilian Nursing Students", *Nurse Educator*, 2021
Publication 1%
- 3 Rostika Flora, Mohammad Zulkarnain, Sukirno. " β -endorphin response to aerobic and anaerobic exercises in Wistar male rats", *Medical Journal of Indonesia*, 2020
Publication 1%
- 4 Submitted to Udayana University
Student Paper 1%

5	Submitted to Xiamen University Student Paper	1 %
6	Zidni Ilma Nafi'a. "Faktor Kepuasan Pasien Terhadap Pelayanan Makanan di Rumah Sakit: Literature Review", Jurnal Manajemen Kesehatan Yayasan RS.Dr. Soetomo, 2021 Publication	1 %
7	Arina Zulfa Azizah, Joyeti Darni, Amilia Yuni Damayanti, Mira Dian Naufalani. "Perbedaan Tingkat Kepuasan Makan Pasien Di Rumah Sakit Bersertifikat Halal Dengan yang Belum Bersertifikat Halal", Ghidza: Jurnal Gizi dan Kesehatan, 2019 Publication	1 %
8	Dimuthu Rathnayake, Shanti Dalpatadu. "A systematic approach to reduce hospital food waste based on patient experience", British Journal of Healthcare Management, 2020 Publication	1 %
9	Submitted to Shelton State Community College Student Paper	1 %
10	Submitted to University of Dundee Student Paper	1 %
11	Suci Dwijayanti, Hermawati -, Sahirah Inas Taqiyah, Hera Hikmarika, Bhakti Yudho Suprpto. "Indonesia Sign Language	1 %

Recognition using Convolutional Neural Network", International Journal of Advanced Computer Science and Applications, 2021

Publication

12

Alfred B. Addison, Billy Wong, Tanzime Ahmed, Alberto Macchi et al. "Clinical Olfactory Working Group Consensus Statement on the Treatment of Post Infectious Olfactory Dysfunction", Journal of Allergy and Clinical Immunology, 2021

Publication

1 %

13

Fauzatun Hadiyati, Ratna Supriastuti, Arifah Mujiharti. "The Effect of Dietary Modification and Nutritional Education on Nutrition Care Process (NCP) for Increasing Dietary Intake, Body Weight, and Nutritional Status of Cancer Patients at Risk of Malnutrition and Malnourished in the Inpatient of Dharmais Cancer", Indonesian Journal of Cancer, 2021

Publication

1 %

14

Submitted to Universitas Respati Indonesia

Student Paper

1 %

15

E. García-Escobar, S. Valdés, F. Soriguer, J. Vendrell et al. "Fatty liver index as a predictor for type 2 diabetes in subjects with normoglycemia in a nationwide cohort study", Scientific Reports, 2021

Publication

1 %

16 Submitted to Universitas Brawijaya 1 %
Student Paper

17 Submitted to Liverpool John Moores University <1 %
Student Paper

18 Submitted to Southern Luzon State University <1 %
Student Paper

Exclude quotes Off

Exclude matches Off

Exclude bibliography Off